arXiv:0712.0401v3 [math-ph] 26 Jun 2008

Universidade de São Paulo
Instituto de Física

Structural and Dynamical Aspects of the AdS/CFT Correspondence: a Rigorous Approach

Pedro Lauridsen Ribeiro

Advisor: João Carlos Alves Barata

Doctorate Thesis submitted to the Institute of Physics of the University of São Paulo for achievement of the title of Doctor in Sciences

Examination Board:
Prof. Dr. João Carlos Alves Barata (IF-USP)
Prof. Dr. Élcio Abdalla (IF-USP)
Prof. Dr. Frank Michael Forger (IME-USP)
Prof. Dr. Severino Toscano do Rêgo Melo (IME-USP)
Prof. Dr. Ricardo Schwarz Schör (ICEx-UFMG)

São Paulo
2007

This is a (hopefully, apart from some corrections throughout the text) faithful English translation of my Ph.D. thesis, originally written in Portuguese, aiming at submission to arXiv's and LQP Crossroads' preprint servers for public use under these servers' terms and conditions – I intend to add some extra chapters in the near future with material which has been excluded from the original Portuguese text due to the lack of spacetime.

I apologise the readers for not being able to provide the English translation of all epigraphs, more specifically regarding the texts of Guimarães Rosa and Edith Pimentel Pinto – not only because I couldn't find them (for the second author, it probably doesn't exist) but also because undertaking the translation task myself would go far beyond my literary capabilities (of course, if someone knows about an English translation, I'd be glad to be informed about it). Should you notice any typographic and/or language errors, I'll also be glad to know about them. Please contact me at `pribeiro@fma.if.usp.br` and/or `pedro.lauridsen.ribeiro@desy.de` regarding these matters.

The Author
Hamburg, December 2007

*This thesis is dedicated to the memory of my grandmother, Maria de Lourdes Pimentel Pinto (*1922 – †1996) and my grand-aunt, Edith Pimentel Pinto (*1921 – †1992), for I know that they watched over me (and still do!) all this time...*

Ἀυτὰ τὰ πράγματα οἷά ἐστιν, διαιροῦντα εἰς ὕλην, αἰτιον, ἀναφοράν.[i]

Marcus Aurelius
Τὰ εἰς εαὐτόν [ii], 12.10

Man hört nur die Fragen, auf welche man imstande ist, eine Antwort zu finden.[iii]

Friedrich Wilhelm Nietzsche
Die fröhliche Wissenschaft[iv], §196

Abstract.

We elaborate a detailed study of certain aspects of (a version of) the *AdS/CFT correspondence*, conjectured by Maldacena [Mal98] and Witten [Wit98], between quantum field theories in a gravitational background given by an asymptotically anti-de Sitter (AAdS) spacetime, and conformally covariant quantum field theories in the latter's conformal infinity (in the sense of Penrose), aspects such that: (a) are independent from (the pair of) specific models in Quantum Field Theory, and (b) susceptible to a recast in a mathematically rigorous mould. We adopt as a starting point the theorem demonstrated by Rehren [Reh00] in the context of Local Quantum Physics (also known as Algebraic Quantum Field Theory) in anti-de Sitter (AdS) spacetimes, called *algebraic holography* or *Rehren duality*.

The main body of the present work consists in extending Rehren's result to a reasonably general class of d-dimensional AAdS spacetimes ($d > 3$), scrutinizing how the properties of such an extension are weakened and/or modified as compared to AdS spacetime, and probing how non-trivial gravitational effects manifest themselves in the conformal infinity's quantum theory.

Among the obtained results, we quote: not only does the imposition of reasonably general conditions on bulk null geodesics (whose plausibility we justify through geometrical rigidity techniques) guarantee that our generalization is geometrically consistent with causality, but it also allows a "holographic" reconstruction of the bulk topology in the absence of horizons and singularities; the implementation of conformal symmetries in the boundary, which we explicitly associate to an intrinsically constructed family of bulk asymptotic isometries, have a purely asymptotic character and is dynamically attained through a process of *return to equilibrium*, given suitable boundary conditions at infinity; gravitational effects may cause obstructions to the reconstruction of the bulk quantum theory, either by making the latter trivial in sufficiently small regions or due to the existence of multiple inequivalent vacua, which on their turn lead to the existence of solitonic excitations localized around domain walls, similar to D-branes. The language employed for the quantum theories relevant for our generalization of Rehren duality follows the functorial formulation of Local Quantum Physics due to Brunetti, Fredenhagen and Verch [BFV03], extended afterwards by Sommer [Som06] in order to incorporate boundary conditions.

Resumo.

Elaboramos um estudo detalhado de alguns aspectos d(e uma versão d)a *correspondência AdS/CFT*, conjecturada por Maldacena [Mal98] e Witten [Wit98], entre teorias quânticas de campo num fundo gravitacional dado por um espaço-tempo assintoticamente anti-de Sitter (AAdS), e teorias quânticas de campos conformalmente covariantes no infinito conforme (no sentido de Penrose) deste espaço-tempo, aspectos estes: (a) independentes d(o par d)e modelos específicos em Teoria Quântica de Campos, e (b) suscetíveis a uma reformulação em moldes matematicamente rigorosos. Adotamos como ponto de partida o teorema demonstrado por Rehren [Reh00] no contexto da Física Quântica Local (também conhecida como Teoria Quântica de Campos Algébrica) em espaços-tempos anti-de Sitter (AdS), denominado *holografia algébrica* ou *dualidade de Rehren*.

O corpo do presente trabalho consiste em estender o resultado de Rehren para uma classe razoavelmente geral de espaços-tempos AAdS d-dimensionais ($d > 3$), escrutinar como as propriedades desta extensão são enfraquecidas e/ou modificadas em relação ao espaço-tempo AdS, e como efeitos gravitacionais não-triviais se manifestam na teoria quântica no infinito conforme.

Dentre os resultados obtidos, citamos: condições razoavelmente gerais sobre geodésicas nulas no interior (cuja plausibilidade justificamos por meio de resultados de rigidez geométrica) não só garantem que a nossa generalização é geometricamente consistente com causalidade, como também permite uma reconstrção "holográfica" da topologia do interior na ausência de horizontes e singularidades; a implementação das simetrias conformes na fronteira, que associamos explicitamente a uma família de isometrias assintóticas do interior construída de maneira intrínseca, ocorre num caráter puramente assintótico e é atingida dinamicamente por um processo de *retorno ao equilíbrio*, mediante condições de contorno adequadas no infinito; efeitos gravitacionais podem eventualmente causar obstruções à reconstrução da teoria quântica no interior, ou por torná-la trivial em regiões suficientemente pequenas ou devido à existência de múltiplos vácuos inequivalentes, que por sua vez levam à existência de excitações solitônicas localizadas ao redor de paredes de domínio no interior, similares a D-branas. As demonstrações fazem uso extensivo de geometria Lorentziana global. A linguagem empregada para as teorias quânticas relevantes para nossa generalização da dualidade de Rehren segue a formulação funtorial de Brunetti, Fredenhagen e Verch para a Física Quântica Local [BFV03], estendida posteriormente por

Sommer [Som06] para incorporar condições de contorno.

Acknowledgements

My sincere thanks to Prof. João Barata, for taking me in at the Department of Mathematical Physics, for proposing the present theme of investigation and for believing me even at the most difficult moments of the development of this project, bearing thus a solid and essential support.

To my former, undergraduate research group, friends up to day I hold dearly: Profs. Alinka Lépine-Szily, Rubens Lichtenthäler Filho and Valdir Guimarães, for the support and enticement when I decided to migrate from experimental/computational Nuclear Physics to Mathematical Physics, and above all for teaching me the first steps of scientific research and a righteous attitude towards it.

To the (ex-)colleagues of the Department of Mathematical Physics (IFUSP) Alessandro Marques, Alex Dias, Amílcar Queiroz, Brenno Vallilo, Davi Giugno, Ivan Pontual, Karlucio Castello Branco, Leandro Beviláqua, Marcelo Pires, "digníssimo" Leonardo Sioufi and Patrícia Teles, for the priceless conversations (in both senses!) and for the company along these more than four years.

To the masters and companions of the Institute of Mathematics and Statistics, for being open to the enriching exchanges of ideas which contributed so much to my learning, in particular the Operator Algebras group (Cíntia da Silva, David Dias, Marcela Merklen, Patrícia Hess and Profs. Cristina Cerri, Martha Salerno, Ricardo Bianconi and Severino Toscano), Prof. Michael Forger's group (Bruno Soares, Eder Annibale, Fernando Antoneli, Getúlio Bulhões, Leandro Gomes, Mário Salles, Paola Gaviria, Sandra Yepes and Sandro Romero) and as well to Profs. Helena Ávila, Paulo Cordaro to the colleagues Alexandre Kawano, Arthur Julião, Antônio Ronaldo Garcia and Nestor Centurión.

I'd also like to thank Profs. Bert Schroer, Christian Jäkel, Daniel Vanzella, Fernando Auil, Jens Mund and, once more, to Prof. Michael Forger for the patience and enthusiasm during their participation at the series of informal seminars along the years 2004–2005 dedicated to the study of the reference [BFV03], which constitutes a found-

ing stone of the conceptual framework underlying the present Thesis. I thank likewise these and Profs. Aiyalam Balachandran, Detlev Buchholz, Henning Rehren, Klaus Fredenhagen, Kostas Skenderis, Sumati Surya e Walter Wreszinski, for the inspiring dialogs and/or email exchanges at crucial moments of the development of the present work, that, hence, constituted an invaluable aid with their suggestions and constructive criticism.

To the secretaries of the Department of Mathematical Physics: Amélia Genova, Elizabeth Vargas e Simone Shinomiya, for the solicitude and tireless sympathy.

To FAPESP, for the financial support (this project was graced with a Direct Doctorate scholarship relative to the process 01/14360-1).

Well, it's a lot of people, and most certainly I forgot someone above... Thanks and my apologies to those!

And, lastly (and first), the more than special thanks:

To my parents, for this journey was also theirs, of us all!

To my brother Francisco, for the spiritual counterpoint.

To Clau, for keeping (enduring!) my mental (in)sanity, and for teaching me the importance of the tings that must be said.

To God, Almighty Eternal Father, for showing me a path that fascinated me and giving me strength to follow it. Let's hope it grants good fruits to His honour!

Contents

Preface xvii

List of Figures xxv

Frequently used notation and conventions xxvii

I The AdS/CFT correspondence and its geometrical manifestations 1

1 The geometry of asymptotically anti-de Sitter spacetimes 3
1.1 Generalities on space forms . . . 3
1.2 Geometry of AdS_d: boundary conditions at infinity . . . 6
1.2.1 Structure of conformal infinity . . . 7
1.2.2 Wedges in AdS and diamonds in the boundary . . . 10
1.3 AAdS spacetimes . . . 14
1.3.1 Global geometry and null geodesics . . . 16
1.3.1.1 Causality (bulk to boundary) . . . 16
1.3.1.2 Simple connectedness (bulk *versus* boundary) . . . 21
1.3.1.3 Localization (boundary to bulk) . . . 23

2 Dynamics and time evolution 31
2.1 Gravitation . . . 31
2.2 The Fefferman-Graham expansion . . . 33
2.3 Gravitational "thermodynamics" in AAdS wedges . . . 40
2.3.1 Geometrical time evolution in diamonds . . . 40
2.3.2 Construction of asymptotic isometries . . . 46
2.3.3 "Return to equilibrium" in wedges . . . 49

II Holographic avatars of Local Quantum Physics 55

3 Locally Covariant Quantum Theories 57
3.1 Basic definitions . . . 58
3.2 Relation with the usual axioms . . . 59
3.3 Quantum states in curved spacetimes . . . 62
3.4 Locally covariant quantum fields . . . 62

4 Implementing the Holographic Principle in AAdS spacetimes 67
4.1 Local covariance with boundary conditions . . . 67
4.2 Locally covariant quantum theories in AdS . . . 72
4.2.1 Boundary values as a dynamical condition . . . 73
4.3 Algebraic holography in AAdS spacetimes . . . 81
4.4 Holographic analysis of superselection sectors . . . 85
4.4.1 A digression . . . 86
4.4.2 (Essential) Haag duality in (A)AdS spacetimes . . . 89
4.4.3 Obstructions to the inversion of Rehren duality . . . 94

III A synthesis 97

Coda 99

Appendices 102

A Elements of Lorentzian geometry 105
A.1 Local theory . . . 105
A.2 Global theory . . . 111
A.2.1 Causal structure, geodesic completeness . . . 111
A.2.2 Conformal infinity . . . 117

B Elements of operator algebras 121
B.1 *-Algebras . . . 122
B.2 C*-algebras and von Neumann algebras . . . 127
B.2.1 C*-algebras . . . 127
B.2.2 von Neumann algebras . . . 129
B.2.3 Tomita-Takesaki modular theory . . . 134
B.2.4 C*- e W*-dynamical systems. KMS condition . . . 138
B.3 Locally convex *-algebras and Borchers-Uhlmann algebras . . . 141

C Categories and functors 145

C.1 Rudiments 145
C.2 Functors and natural transformations 148

D Rudiments of homotopy **153**
D.1 Homotopic maps 153
D.2 The fundamental group and covering spaces 154

Bibliography **157**

Notes and translations of (some) epigraphs **169**

Erratum **171**

Preface

Semântica

Eu sei lá o que querem dizer as palavras.

Sou caminho
e se imanto as limalhas a meu jeito
o desenho não é figurativo.

E se empreendo, gaguejando,
a leitura do meu campo de esforço
o feito
infante ainda
é mensagem cifrada.

Edith Pimentel Pinto
"Normativa" (*Sinais e Conhecenças*[v], 1986)

Two of the greatest open problems in contemporary theoretical physics are the lack of quantitative comprehension of the nonperturbative aspects of non Abelian gauge theories (quark confinement, etc.) and the quantization of the gravitational field, which makes itself phenomenologically necessary due to the limits of physical consistency of General Relativity, which describes classical gravity up to what we've been able so far to test experimentally. One of the most tantalizing ideas in current literature is that both problems are somehow related by a *holographic* correspondence, in which gravity in a certain region of spacetime would admit a "dual" description at the boundary of such a region by means of a gauge theory, in the same way that a bidimensional hologram reproduces a tridimensional image. Moreover, the energy regimes of the physical processes associated to both theories are of such nature that *nonperturbative* aspects of the gauge theory would manifest themselves at the *(semi)classical* gravitational regime.

The history of how this idea arose goes back to the seventies. The four laws of black hole dynamics, established by Bardeen, Carter and Hawking [BCH73], possess a

strong analogy with the 4 laws of thermodynamics:

ZEROTH LAW [BCH73]: The surface gravity κ is constant throughout the event horizon of a stationary black hole [1] $\leftrightarrow$ a thermal bath in equilibrium possesses constant temperature T throughout all its extension.

FIRST LAW [BCH73]: $\delta M = \Omega_H \delta J + \frac{1}{8\pi}\kappa\delta A$, where M is the mass of the black hole, Ω_H is the magnitude of the axial component of the timelike KILLING field tangent to the horizon, J the angular momentum of the black hole and A the area of (a spatial section of) the event horizon. The variations of the metric implicit in the formula are also (quasi)stationary [1] $\leftrightarrow \delta Q = P\delta V + T\delta S$, where Q is the amount of heat, P the pressure, V the volume and S the entropy.

SECOND LAW [Haw71]: *Area Theorem* – the area of (spatial sections of) the event horizon cannot decrease throughout the CAUCHY evolution of a nonstationary black hole satisfying the null energy condition (NEC) [2] $\leftrightarrow$ the entropy of a closed system cannot decrease by means of any physical process.

THIRD LAW [Isr86]: A nonstationary, sufficiently regular black hole satisfying the weak energy condition cannot become *extremal*, i.e., lose its trapped surfaces (i.e., acausal, codimension-two surfaces such that both of their normal null geodesic congruences possess negative expansion) external to its event horizon in finite (retarded) time, which would imply $\kappa \to 0$ [3] $\leftrightarrow$ it's impossible to reach $T = 0$ in finite time by means of any physical process.

In particular, the *area* of the event horizon of a black hole corresponds to the *entropy*. BEKENSTEIN [Bek74] suggested that such an identification was more than just a mere analogy – the entropy of an black hole in equilibrium would, indeed, be given by the

[1] Although the zeroth, first and second laws are stated in [BCH73], only the zeroth and first laws are proved in this reference, for perturbations stationary black holes (the third law is only suggested in a rudimentary form).

[2] The statement and proof of the second law were previously elaborated by HAWKING [Haw71] in a nonstationary context.

[3] The third law was formulated in the present precise form and demonstrated by ISRAEL [Isr86]. One must recall that NERNST's formulation for the third law, i.e., $S = 0$ at $T = 0$, is neither universally valid (a typical example is the ice, which possesses a residual entropy at $T = 0$ due to the hydrogen bridges. Such a fact was established in an exact way in two dimensions by LIEB [Lie67a, Lie67b]. Notice that the free BOSE gas, which goes under BOSE-EINSTEIN condensation at $T = 0$ and, hence, possesses degeneracy at the ground state, *doesn't* constitute another (counter)example, for the asymptotic behaviour of such a degeneracy in the partition function as $T \searrow 0$ is just polynomial, and not exponential as in the case of ice, resulting in $S = 0$ if we adopt BOLTZMANN's definition for the entropy. I thank Prof. Walter Wreszinski for calling my attention to this point), nor is its "natural" analog for black holes [Wal97].

formula

$$S = \frac{A_{hor}}{4},$$

and a generalization of the second law of thermodynamics, taking into account the matter entropy (outside the black hole) as well as the entropy of the black hole, should hold. This scenario, suggested by HAWKING's Area Theorem [Haw71], is not necessarily valid classically; however, the work of HAWKING [Haw75] showed [1] that a *quantum* field coupled to the gravitational field of a black hole *thermalizes*, after a sufficiently long time interval, *precisely at temperature* $T = \frac{\kappa}{2\pi}$, *suggested by the first law of black hole dynamics!* This result is surprising, for the latter is a purely geometrical quantity, which indicates an *universal* (i.e., independent of the quantum field model we couple to the classical gravitational field) character for the laws of black hole thermodynamics.[2] Motivated by such a result, 'T HOOFT [Hoo93] proposed taking BEKENSTEIN's idea to the last consequences: *the microscopic degrees of freedom of a quantum theory of gravity are completely encoded in the boundary of the spacetime's volume where such theory is defined.* This principle, formalized later by SUSSKIND [Sus95], is named *holographic principle*.

Within the integrated description of gauge theories and gravity proposed by string theory, two results suggest that this theory may provide a realization of the holographic principle: the microscopic calculation of BEKENSTEIN entropy made by STROMINGER and VAFA [SV96], and, above all, the work of MALDACENA [Mal98], which, by means of a(n effective limit of a) string model on the background spacetime $AdS_5 \times S^5$, where the AdS_5 factor is the anti-DE SITTER (AdS) spacetime in five dimensions, obtained the field multiplet of a supersymmetric, conformally invariant gauge theory with structural group $SU(N)$ in the $N \to \infty$ limit, living at the conformal boundary of AdS_5 (the S^5 factor becomes a global group of internal (R-)symmetries $SU(4) \sim SO(6)$). MALDACENA conjectured that this relation establishes a one-to-one correspondence between both theories, even outside the effective limit. The formulation of this correspondence in terms of k-point functions was obtained by GUBSER, KLEBANOV and POLYAKOV [GKP98] in a particular case, and more in general by WITTEN [Wit98]. In these two works, it was shown that the (SCHWINGER) k-point functions of the gauge theory in the above limit could be obtained from variations of the *classical* (Euclidean) supergravity

[1] The geometrical optics approximation for the two point function near the horizon of a SCHWARZSCHILD, black hole, adopted by HAWKING in [Haw75], is difficult to justify, for the behaviour of the metric in this region implies an index of refraction that varies very fast along the relevant time coordinate. A conceptually precise calculation of HAWKING radiation was obtained by FREDENHAGEN and HAAG [FH90].

[2] From the viewpoint of the related UNRUH *effect*, this is not so surprising, if one has in mind the results of BISOGNANO and WICHMANN in the context of WIGHTMAN quantum fields [BW75, BW76, Sew82]. However, such a comparison demands care, as the circumstances of the HAWKING effect are different in several aspects [Wal94].

action around its value at the background geometry under variation of the boundary conditions at conformal infinity. Moreover, WITTEN's prescription for the generating functional of the conformal field theory allowed one to rephrase MALDACENA's statement uniquely in terms of usual field theories, without reference to strings. Such a correspondence became known in literature as *AdS/CFT correspondence*. By means of this formulation, the MALDACENA conjecture was successfully tested in a myriad of particular cases.[3]

WITTEN's formulation for the AdS/CFT correspondence opened way to the following question: might it be possible to establish a correspondence between quantum field theories (QFT's) in AdS and conformally covariant QFT's in AdS's conformal infinity, *uniquely from the fundamental principles of QFT?* The question indeed happens to have a *positive* answer: within a version of WIGHTMAN's axiomatic framework [SW00] for quantum fields in AdS, a correspondence between k-point functions in the (latter's) bulk conformally invariant k-point functions in the conformal boundary was rigorously established by BERTOLA, BROS, MOSCHELLA and SCHAEFFER [BBMS00]. Moreover, a correspondence between bulk and boundary algebras of local observables was demonstrated by REHREN [Reh00], within the algebraic formalism of HAAG and KASTLER [HK64, Haa96] for QFT (*Local Quantum Physics*) – the latter correspondence is named *algebraic holography* or REHREN *duality*. Although the relation between these two formulations and WITTEN's prescription has been investigated in several aspects [DR02, DR03, Reh05], a fundamental question remained unanswered: *how gravitational effects are encoded in these formulations?*

As both versions refer themselves uniquely to (pure) AdS spacetimes (for they depend critically on its causal structure), and not to more general geometries possessing the same conformal infinity of AdS (i.e., *asymptotically AdS* (AAdS) spacetimes), even the formulation *per se* of such rigorous counterparts in a more general geometrical context becomes elusive. There are, inclusively, evidences [AS02] that, for nontrivial AAdS geometries, REHREN duality becomes physically incompatible with known tests of the MALDACENA conjecture, and, as such, needs to be reformulated. Which takes us to a second question, intimately linked to the former: *How much from the underlying framework of* REHREN *duality and its properties does survive in "generic" AAdS spacetimes?*

[3]Unfortunately, the phenomenal research activity aroused by the fundamental works about the AdS/CFT correspondence, cited above, made practically impossible to elaborate even a pretensely representative bibliography of its different directions (the work [Mal98] alone possesses more than 4.700 citations by 30/07/2007, according to the SPIRES-HEP online database). Thus, we shall content ourselves in citing only works of direct relevance to our development, as needed, and forward the interested reader SPIRES-HEP's homepage (`http://www.slac.stanford.edu/spires/`) for other directions.

The body of the present Thesis proposes a natural generalization of REHREN duality to AAdS spacetimes and investigates how the properties of the original version are modified by gravitational effects in the bulk, seeking to elucidate both questions elaborated above.

One of the fundamental lessons of Local Quantum Physics is that the physical information of a quantum field theory is contained not in the individual fields / observables, but in the relative inclusions of local algebras. In particular, such information is highly sensitive to the causal structure of spacetime. Hence, it's to be expected that the distortion of the causal structure of the bulk of an AAdS spacetime due to gravitational effects be "felt" by the "dual" theory at the boundary. We'll see, after a qualitative, global study of this deformation within a reasonably general set of hypotheses about the bulk geometry, that this indeed occurs. We shall show effects of two kinds:

1. The implementation of asymptotic isometries, far from being an immediate fact as it is for AdS, is dynamically attained as a process of *return to equilibrium*, which has as a consequence the potential breaking of conformal symmetry. The states over the boundary quantum theory obtained by this process possess a rather rich structure, of which we've only scratched the surface, and whose thermal properties deviating from equilibrium encode nontrivial details of the bulk geometry.

2. The structure of superselection sectors is radically modified by gravitational effects. Given the hypothesis that arbitrarily small bulk regions possess nontrivial observables, plus other natural ones within the context of the algebraic theory of superselection sectors [Ara99, Haa96], we'll show that the bulk theory acquires soliton excitations localized around domain walls, similar to the D-branes that occur in the original formulation of the AdS/CFT correspondence, signaling spontaneous breaking of internal symmetries. The nontrivial increase of the local algebras at the boundary needed to the incorporation of the intertwiners among different partial vacua makes these algebras to acquire *nonlocal* (i.e., extended) elements implementing internal symmetries in the region where each partial vacuum differs from the total vacuum. And, precisely because of these elements, it becomes fundamentally impossible, unlike in pure AdS, to completely reconstruct the bulk quantum theory from the boundary theory alone (Subsection 4.4.3, page 94).

We've put emphasis, in this work, in robust, mathematically rigorous results which are independent from a specific quantum field model, by virtue of the universal character of several aspects of the AdS/CFT correspondence. We shall not investigate in detail, though, the relation of the formalism proposed here to WITTEN's prescription – such a task lies beyond the scope of the present work.

On the structure of this Thesis

This Thesis has three parts. In Part I, which comprises Chapters 1 and 2, we develop the geometrical part of the work. In Part II, which comprises Chapters 3 and 4, we present our proposal of generalization of REHREN duality within a general formalism for QFT in curved spacetime which incorporates covariance and locality in functorial manner, and study universal (i.e., independent of specific models) consequences from the dynamical and structural viewpoints. Part III consists only of the Coda, which concludes the work.

In Chapter 1, we'll start a systematic treatment of anti-DE SITTER (AdS) and asymptotically anti-DE SITTER (AAdS) spacetimes. We shall investigate, within a set of hypotheses about the global behaviour of null geodesics, how the causal structure is globally modified by gravitational effects consistent with the boundary conditions at infinity (Proposition1.8, page 20) – the effect is mainly due to gravitational time delay of null geodesics traversing the bulk (Theorem 1.3, page 16). Some of these results are not new [Woo94, GW00], but the technique of gravitational time delay is fundamental to most of the geometrical developments to follow. We shall define a simple generalization of *wedge* regions, employed by REHREN in [Reh00], to AAdS spacetimes (Definition 1.5, page 15) – such regions generalize, in a certain sense, the RINDLER wedge $\{x^1 > |x^0|\}$ in MINKOWSKI spacetime, and constitute the prototype of an event horizon of an *asymptotically* stationary black hole in a background with negative cosmological constant. We'll show how it's possible to reconstruct the bulk topology by means of intersections of wedges enveloping sufficiently small diamonds (Subsubsection 1.3.1.3, page 23). This result is original [Rib07], and the proof involves an interesting application of the method employed by PENROSE [HE73, Wal84] to prove the existence of singularities as result of gravitational collapse, together with a compactness argument, to obtain a geometric maximum principle that guarantees the precise envelope (Theorem 1.14, page 27). The presentation of this Chapter has several improvements with respect to [Rib07].

In Chapter 2, we'll investigate the dynamics of the EINSTEIN equations in AAdS spacetimes (Section 2.2, page 33). Such a discussion aims at estimating in a more quantitative way nontrivial gravitational effects in terms of geometrical quantities defined in the conformal boundary. We'll also construct in an intrinsic manner families of asymptotic isometries (boosts) naturally associated to each AAdS wedge (Proposition 2.5, page 46), and, more in general, families of asymptotically conformal diffeomorphisms associated to any relatively compact diamond in causally simple spacetimes (formulae (2.37)–(2.39), page 45). To such families, it's possible to associate a *surface gravity* in the horizon of each wedge or diamond (Definition 2.2, page 51) and, hence, formulate an asymptotic analog of the zeroth law of black hole (thermo)dynamics (Theorem 2.6, page

51), and corresponding results for the second law (contained in Proposition 1.8, page 20, and properly reinterpreted in the context of this Chapter) and the characterization of reversible processes (Theorem 2.7, page 54, which follows directly from Proposition 1.8, *ibid.* and Remark 1.7, page 21). The presented construction is sufficiently robust to be implemented in situations not necessarily linked to the geometrical context of the AdS/CFT correspondence.

In Chapter 3, we'll introduce a generalization of the formalism of HAAG and KASTLER to curved spacetimes, proposed by BRUNETTI, FREDENHAGEN and VERCH [BFV03] and formulated in terms of the language of *categories* and *functors*.

In Chapter 4, we'll present in a precise way our proposal of generalization for REHREN duality. For such, it's necessary to adapt the constructions of Chapter 3 to our present geometrical context, which demands the imposition of *boundary conditions* in a covariant manner. Thus, we shall initiate our discussion with the algebraic treatment of this issue proposed by SOMMER [Som06], which investigates how it's possible to extend a locally covariant quantum theory in the sense of BRUNETTI ET AL. to non globally hyperbolic spacetimes (Definition 4.1, page 70). One follows the example of local quantum theories in anti-DE SITTER spacetimes – here, we sought to insert the formulation proposed by BUCHHOLZ, FLORIG and SUMMERS [BFS00] within a locally covariant context. The advantage of this formulation, besides the minimal number of premises, is the imposition of boundary conditions at conformal infinity on the "elementary" states in the form of a condition of thermodynamic stability (formulae (4.6) and (4.7), page 73) – which, in the case of AAdS spacetimes, is generalized in the form of a condition of "return to equilibrium" (condition (d), page 84). Employing the techniques of BORCHERS and YNGVASON [BY99] (stated in Theorems 4.1, 4.2 and 4.3, page 75ff.), we shall show that such a condition can be understood as a prescription for the scaling behaviour of observables in a neighborhood of infinity (Proposition 4.4, page 77), which can be generalized to asymptotically anti-DE SITTER spacetimes. With this framework at hand, the generalization of REHREN duality (Definition 4.3, page 73) we seek emerges naturally (Definition 4.8, page 82). We'll investigate the structure of states over the obtained boundary theory by means of this generalization, and we'll end up with an outline of a correspondence between superselection sectors of the bulk quantum theory and its boundary dual, linking sectors localized in diamonds (i.e., in the sense of DOPLICHER, HAAG and ROBERTS [Haa96]) at the boundary and soliton sectors in the bulk, localized around codimension-two domain walls, quite similar to D-branes.

In the Coda, we'll present our conclusions and list possible future directions of investigation suggests by the present work.

A pedagogical obstacle to the conception of the present Thesis was the broad spectrum of employed mathematical techniques – global Lorentzian geometry, operator algebras, categories and functors. Hence, we've collected most of the needed mathematical concepts in four Appendices, in order not to deviate the reader's attention from the central ideas of the present work and make the text mathematically self contained.

Appendix A summarizes the necessary notions of Lorentzian geometry and causal structure.

Appendix B presents basic concepts of Operator Algebras (C*-algebras, VON NEUMANN algebras and BORCHERS-UHLMANN algebras), including a minimal introduction to TOMITA-TAKESAKI theory, employed in Chapter 4. This Appendix begins with a detailed treatment of *-algebras, seeking to unify the presentation of the results and concepts of operator algebras that depend only on the algebraic structure.

Appendix C condensates the basic concepts of categories and functors employed in Chapters 3 and 4.

Appendix D draws quick strokes of the concepts of homotopy needed in Chapter 1.

I wish you all a nice reading!

The Author
São Paulo and Hamburg, resp. September and December 2007

List of Figures

1.1 **(a)** Fundamental domain of AdS_d (red), given by the quotient modulo $\mathbb{Z}$, where the spatial infinity is represented in blue and the (fundamental) Poincaré domain is the upper part of the dotted cut; **(b)** Image of the conformal embedding of AdS_d into ESU_d (red). The infinity $\mathscr{I}$ of AdS_d is represented in blue and the Poincaré domain $\mathscr{P}oi(p)$ is delimited by the dotted curves; **(c)** $\mathscr{P}oi(p)$ (side view) delimited by the dotted lines in red, with $\mathscr{I}$ represented in blue; **(d)** Visualization of $\mathscr{I}$ as ESU_{d-1} (red) with the Minkowski domain $\mathscr{M}in(p)$, corresponding to the image of the conformal embedding of $\mathbb{R}^{1,d-2}$ into ESU_{d-1}, in blue. 9

1.2 **(a)** Correspondence between bulk wedges and boundary diamonds. **(b)** A Minkowski domain and the corresponding Poincaré domain. 12

1.3 Gravitational time delay. The null geodesics which emanate from $p \in \mathscr{I}$ and remain in $\mathscr{I}$ (full lines) focus at the antipodal $\bar{p}$. Whereas in AdS_d a null geodesic emanating from p and traversing the bulk would also have $\bar{p}$ as its future endpoint (dotted line), in an AAdS spacetime satisfying the hypothesis of global focusing of Theorem 1.3 such a geodesic has future endpoint $q \in I^+(p, \mathscr{I})$ (dashed/dotted line). 17

1.4 Effect of gravitational time delay (whose effect on null geodesics is exemplified by the dotted lines in the longitudinal cut **(b)** of **(a)**) on wedges and their causal complements in AAdS spacetimes. Whereas in AdS_d the causal complement of the wedge $\mathscr{W}_{p,\bar{q}}$ is the wedge $\mathscr{W}_{q,\bar{p}}$ (red), the causal complements of both wedges (respectively green and blue) are no longer wedges in AAdS spacetimes satisfying the conditions of Proposition 1.8 and, hence, of Theorem 1.3, and in such a way that the open region $\mathscr{W}'_{p,\bar{q}} \cap \mathscr{W}'_{q,\bar{p}} \neq \varnothing$ is *nonvoid*, which cannot happen in AdS_d. 22

Frequently used notation and conventions

- $\mathbb{Z}$, $\mathbb{R}$, $\mathbb{C}$ – respectively, the ring of integers and the fields of real and complex numbers. $\mathbb{Z}_{+/-}$, $\mathbb{R}_{+/-}$ – respectively, the positive (> 0) / negative (< 0) integers and the positive / negative reals. $\bar{\mathbb{Z}}_{+/-}$, $\bar{\mathbb{R}}_{+/-}$ – respectively, the non negative (≥ 0) / non positive (≤ 0) integers and the non negative / non positive reals.

- The real and imaginary parts of $z = x + iy \in \mathbb{C}$ are respectively denoted by $\Re z \doteq x$ and $\Im z \doteq y$, and the complex conjugate of z, by $\bar{z} \doteq x - iy$.

- The power set of X is denoted by $P(X) \doteq \{U \subset X\}$.

- Given any sets A, S, $A^S \doteq \{a : S \to A\}$ denotes the *Cartesian power* of A to S. In particular, if $S = \{1, \ldots, n\}$, we write $A^S \doteq A^n = \{(a_1, \ldots, a_n) : a_i \in A, \forall i = 1, \ldots, n\}$.

- Given an open set $\mathscr{O}$ in a topological space X, $K \Subset \mathscr{O}$ denotes that K possesses compact closure $\bar{K}$ and $\bar{K} \subset \mathscr{O}$.

- Given a symmetric non degenerate bilinear form g over $\mathbb{R}^d$, the *signature* of g is given by the pair of integers (p, q) or by their difference $p - q$, where $p + q = d$ and p is the number of *negative* eigenvalues of g, named *index* of g. We associate to g the symmetric non degenerate bilinear form $g^{-1} \doteq [g_{ij}]^{-1}$, of same index, over the dual of $\mathbb{R}^d$, and the volume d-form given by $\sqrt{|g|} \doteq \sqrt{((-1)^p \det[g_{ij}])^{\frac{1}{2}}}$, where $[g_{ij}]$ is the matrix associated to g in some orthonormal basis.

- Manifolds and their subsets are denotes by capital calligraphic letters (e.g. $\mathscr{M}$, $\mathscr{O}$, $\mathscr{I}$, etc.). All manifolds employed in the text are $\mathscr{C}^\infty$, σ-compact (and, thus, paracompact), connected and orientable, unless otherwise indicated (regarding the notions of topology involved, see [Dug66]).

- Given a manifold $\mathscr{M}$, we denote the tangent and cotangent spaces at $p \in \mathscr{M}$ respectively by $T_p\mathscr{M}$ and $T_p^*\mathscr{M}$, and the tangent and cotangent bundles, by $T\mathscr{M}$ and $T^*\mathscr{M}$.

- More generally, we write any vector bundle over a d-dimensional $\mathscr{M}$ as $\mathscr{E} \xrightarrow{p} \mathscr{M}$, where p denotes the projection (surjective submersion) map, $\mathscr{E}$ the total $(D+d)$-dimensional space and $\mathscr{M}$ the basis. The typical fibre is usually denoted by $E_x \doteq p^{-1}(x) \cong \mathbb{R}^D$, $\forall x \in \mathscr{M}$.

- Given a vector field X, we denote the LIE derivative of functions / sections along X by $\pounds_X$.

- Tensor indices are given by small Greek letters μ, ν, etc. when written in terms of local coordinates, and by small Latin letters a, b, etc. when they only indicate the rank of the tensor, without reference to coordinates (PENROSE–RINDLER abstract index notation). In both cases, the contraction of indices follows EINSTEIN's convention.

- We denote by δ^a_b the KRONECKER delta (= identity matrix as a linear operator in $T^{(*)}\mathscr{M}$).

- Given a rank-(r,s) tensor $T^{\mu_1\cdots\mu_r}_{\nu_1\ldots\nu_s}$, we denote respectively the symmetric antisymmetric parts of T w.r.t. contravariant indices $\mu_j, \ldots, \mu_k$, $1 \leq j < k \leq r$ by

$$T^{\mu_1\cdots(\mu_j\cdots\mu_k)\cdots\mu_r}_{\nu_1\cdots\nu_s} \doteq \frac{1}{(k-j+1)!} \sum_{\pi\in S_{k-j+1}} T^{\mu_1\cdots\mu_{\pi(j)}\cdots\mu_{\pi(k)}\cdots\mu_r}_{\nu_1\cdots\nu_s}$$

and

$$T^{\mu_1\cdots[\mu_j\cdots\mu_k]\cdots\mu_r}_{\nu_1\cdots\nu_s} \doteq \frac{1}{(k-j+1)!} \sum_{\pi\in S_{k-j+1}} \mathrm{sgn}(\pi) T^{\mu_1\cdots\mu_{\pi(j)}\cdots\mu_{\pi(k)}\cdots\mu_r}_{\nu_1\cdots\nu_s},$$

where S_l is the group of permutations of l elements and $\mathrm{sgn}(\pi) = 1$ in case π involves an even number of 2-element switchings, and -1 if it involves an odd number of them. Analogously, we denote respectively the symmetric and antisymmetric parts of T w.r.t. covariant indices $\nu_m, \ldots, \nu_n$, $1 \leq m < n \leq s$, by $T^{\mu_1\cdots\mu_r}_{\nu_1\cdots(\nu_m\cdots\nu_n)\cdots\nu_s}$ e $T^{\mu_1\cdots\mu_r}_{\nu_1\cdots[\nu_m\cdots\nu_n]\cdots\nu_s}$. If we want to exclude the interval $\mu_j, \ldots, \mu_k$ from the symmetrization or antisymmetrization of the indices $\mu_1, \ldots, \mu_l$, we'll write respectively $(\mu_1\cdots|\mu_j\cdots\mu_k|\cdots\mu_l)$ or $[\mu_1\cdots|\mu_j\cdots\mu_k|\cdots\mu_l]$. The corresponding formulas for (anti)symmetrization of a tensor shall have the combinatorial factor $\frac{1}{l!}$ substituted by $\frac{1}{(l-k+j-1)!}$. All these conventions hold without modification for abstract indices.

- $\mathbb{R}^{p,q}$ denotes the semi-Riemannian manifold $(\mathbb{R}^{p+q}, \eta)$ (see Appendix A), where

the metric η is given by the symmetric non degenerate bilinear form of index p

$$\eta = \mathrm{diag}(-\overset{p}{\cdots}-+\overset{q}{\cdots}+) \doteq \left[\begin{array}{ccc|ccc} -1 & & 0 & & & \\ & \ddots & & & 0 & \\ 0 & & -1 & & & \\ \hline & & & +1 & & 0 \\ & 0 & & & \ddots & \\ & & & 0 & & +1 \end{array}\right] .$$

In particular, d-dimensional MINKOWSKI spacetime is given by $\mathbb{R}^{1,d-1}$ – that is, we adopt the signature convention $(-+\cdots+)$ (index 1) for Lorentzian metrics.

- Given $x, y \in \mathbb{R}^d$, the Euclidean norm of x is denoted by $|x|$, and the corresponding scalar product, by $\langle x, y\rangle$. The LEBESGUE measure of a BOREL subset A of $\mathbb{R}^d$ is denoted by $|A|$ (for the relevant notions of measure theory, see [Mun71, Rud87]).

- S^d denotes the d-dimensional sphere $\{x \in \mathbb{R}^{d+1} : |x| = 1\}$, whose induced Euclidean metric is denoted by $d\Omega_d^2 = (d\theta^1)^2 + \sin^2\theta^1((d\theta^2)^2 + \sin^2\theta^2(\cdots((d\theta^{d-1})^2 + \sin^2\theta^{d-1}(d\theta^d)^2)\cdots)) = (d\theta^1)^2 + \sin^2\theta^1 d\Omega_{d-1}^2$, where $\theta^I = (\theta^1, \ldots, \theta^d)$ are the angular (spherical) coordinates of S^d, i.e., $0 \leq \theta^1, \ldots, \theta^{d-1} \leq \pi$ and $0 \leq \theta^d < 2\pi$.

- The norm of a normed vector space is denoted by $\|.\|$, with additional indication when needed.

- The scalar product of a pre-Hilbert space (real or complex) is denoted by $\langle ., . \rangle$. We set the convention that, in the complex case, the sesquilinear form $\langle ., . \rangle$ is linear in the *second* variable and anti-linear in the *first*.

- *-algebras, in particular C*- and VON NEUMANN, are denoted by Gothic capital letters (e.g. $\mathfrak{F}$, $\mathfrak{A}$, $\mathfrak{R}$, etc.). Such algebras, if unital, have their identity element denoted by $\mathbb{1}$ (see Appendix B).

- Functors are also denoted by Gothic capital letters.

- Asymptotic behaviour – we say that $f(x) = O(g(x))$ as $x \to x_0$ (x_0 can be $\pm\infty$) if there exists a neighbourhood $\mathscr{U} \ni x_0$ and $C > 0$ such that $|f(x)| \leq C|g(x)|$, $\forall x \in \mathscr{U}$ and $f(x) = o(g(x))$ as $x \to x_0$ if for all $\epsilon > 0$ there exists a neighbourhood $\mathscr{U} \ni x_0$ such that $|f(x)| < \epsilon|g(x)|$, $\forall x \in \mathscr{U}$; f and g are said to be *equivalent* (notation: $f(x) \sim g(x)$) as $x \to x_0$ if there exists a neighbourhood $\mathscr{U} \ni x_0$ and $C > 0$ such that $C^{-1}g(x) \leq f(x) \leq Cg(x)$, $\forall x \in \mathscr{U}$. Given a sequence of functions $\{g_k\}_{k\in\mathbb{Z}_+}$, we say that f is an *asymptotic sum* of g_k, $k \in \mathbb{Z}_+$ as $x \to x_0$ (notation: $f \sim_{x\to x_0} \sum_{k=1}^{\infty} g_k$) if $|f - \sum_{k=1}^{K} g_k| = O(g_{K+1})$, for all K.

- $\mathscr{D}(\mathscr{O})$ denotes the space of $\mathscr{C}^\infty$ functions of compact support in $\mathscr{O}$ (*test functions*), and $\mathscr{S}(\mathbb{R}^d)$, the space of $\mathscr{C}^\infty$ functions whose derivatives of order ≥ 0 go to 0 faster tan any polynomial in x as $|x| \to \infty$ (*tempered* or SCHWARTZ test functions). We also write, following L. SCHWARTZ, $\mathscr{E}(\mathscr{O}) = \mathscr{C}^\infty(\mathscr{O})$. We shall employ these different notations alternately along the text. Respectively, the (general) distributions, tempered e compactly supported distributions are given by the topological duals $\mathscr{D}'(\mathscr{O})$, $\mathscr{S}'(\mathbb{R}^d)$, $\mathscr{E}'(\mathscr{O})$.

- When $\mathscr{O} \subset \mathscr{M}$, where $(\mathscr{M}, g)$ is a d-dimensional, orientable semi-Riemannian manifold, we use invariant volume element $\sqrt{|g|}$ to identify test function spaces with the corresponding test densities (= d-forms, since we've assumed $\mathscr{M}$ orientable). Hence, the topological duals are distributions and not distributional densities (i.e., by the *"smearing"* $u(f)$ of a distribution u with a test function f, one understands the expression $u(f\sqrt{|g|})$, for the metric will always be clear in context).

- Given a vector bundle $\mathscr{E} \xrightarrow{p} \mathscr{M}$, the space of *sections* $\mathscr{C}^\infty$ of $\mathscr{E}$ is the $\mathscr{C}^\infty(\mathscr{M})$-module denoted by $\Gamma^\infty(\mathscr{M}, \mathscr{E}) \doteq \{\phi : \mathscr{M} \xrightarrow{\mathscr{C}^\infty} \mathscr{E} : p \circ \phi = \mathrm{id}_{\mathscr{M}}\}$. Denoting the *zero section* of $\mathscr{E}$ (i.e., which takes the value of the origin at each fibre) simply by 0, we define the *support* $\mathrm{supp}\phi$ of a section ϕ as the complement of the largest open set in $\mathscr{M}$ where $\phi = 0$. We denote, then, space of compactly supported $\mathscr{C}^\infty$ sections of $\mathscr{E}$ by $\Gamma^\infty_c(\mathscr{M}, \mathscr{E})$.

En efeto, rematado ya su juicio, vino a dar en el más estraño pensamiento que jamás dio loco en el mundo, y fue que le pareció convenible y necesario, así para el aumento de su honra como para el servicio de su república, hacerse caballero andante y irse por todo el mundo con sus armas y caballo a buscar las aventuras y a ejercitarse en todo aquello que él había leído que los caballeros andantes se ejercitaban, deshaciendo todo género de agravio y poniéndose en ocasiones y peligros donde, acabándolos, cobrase eterno nombre y fama. Imaginábase el pobre ya coronado por el valor de su brazo, por lo menos del imperio de Trapisonda; y así, con estos tan agradables pensamientos, llevado del estraño gusto que en ellos sentía, se dio priesa a poner en efeto lo que deseaba. Y lo primero que hizo fue limpiar unas armas que habían sido de sus bisabuelos, que, tomadas de orín y llenas de moho, luengos siglos había que estaban puestas y olvidadas en un rincón.[vi]

Miguel de Cervantes Saavedra
El ingenioso hidalgo don Quijote de la Mancha[vii], Primero Libro, Cap. I

Part I

The AdS/CFT correspondence and its geometrical manifestations

Chapter 1

The geometry of asymptotically anti-de Sitter spacetimes

Devia ou não devia contar-lhe, por motivos de talvez. Do que digo, descubro, deduzo. Será, se? Apalpo o evidente? Trebusco. Será este nosso desengonço e mundo o plano – intersecção de planos – onde se completam de fazer as almas?

João Guimarães Rosa
"O espelho" (*Primeiras Estórias*[viii])

1.1 Generalities on space forms

Among the d-dimensional Lorentzian metrics g which solve the Einstein equations without matter and with cosmological constant Λ

$$\mathrm{Ric}(g) - \frac{1}{2}R(g)g + \Lambda g = 0, \tag{1.1}$$

the simplest ones are those with constant sectional curvature.[1] More precisely, if g is such that its sectional curvature

$$K(g)(X,Y) \doteq \frac{g(\mathrm{Riem}(g)(X,Y)X,Y)}{g(X,X)g(Y,Y) - g(X,Y)^2} = C \tag{1.2}$$

[1] For $d < 4$, these are the *only* solutions of (1.1)! See Appendix A, formula (A.2), page 106 discussion that follows.

for all pairs X, Y of tangent vectors at a point p generating a plane in which g is a nondegenerate bilinear form [2] (C may, em principle, depend on p), then g satisfies (1.1) if and only if $C = \frac{2}{(d-1)(d-2)}\Lambda$. In this case, we have:

$$\mathrm{Riem}(g)_{abcd} = \frac{2\Lambda}{(d-1)(d-2)}(g_{ac}g_{bd} - g_{ad}g_{bc}) \tag{1.3}$$

($g_{ac}g_{bd} - g_{ad}g_{bc}$ can be seen as the metric induced by g in the tangent space of 2-vectors; $\mathrm{Riem}(g)_{abcd}$, on its turn, is a symmetric bilinear form in this space. The identity (1.3) hence follows from (1.2) by polarization). Conversely, any d-dimensional Lorentzian manifold with constant sectional curvature C satisfies (1.1) with $\Lambda = \frac{(d-1)(d-2)}{2}C$. In these circumstances, we can invoke two results (see [O'N83] for the proof of both):

Theorem 1.1 ([O'N83]) *Let $(\mathscr{M}, g)$ and $(\mathscr{M}', g')$ be two d-dimensional Lorentzian manifolds of constant sectional curvature $K(g) = K(g') = C$. Then any points $p \in \mathscr{M}$, $p' \in \mathscr{M}'$ possess isometric neighbourhoods. If $\mathscr{M}'$ is geodesically complete, the isometry in question extends uniquely to a* local isometry *$\phi : \mathscr{M} \to \mathscr{M}'$ (i.e., the tangent map $d\phi(p) : T_p\mathscr{M} \to T_{p'}\mathscr{M}'$ is a isometry, for all $p \in \mathscr{M}$).* □

Theorem 1.2 ([O'N83]) *Let $(\mathscr{M}, g)$ and $(\mathscr{M}', g')$ be two d-dimensional, geodesically complete and connected Lorentzian manifolds with constant sectional curvature $K(g) = K(g') = C$. Then, for any points $p \in \mathscr{M}$, $p' \in \mathscr{M}'$ and any isometry $L : T_p\mathscr{M} \to T_{p'}\mathscr{M}'$, there exists a unique isometric covering map (see Definition D.3, page 155) $\phi : \mathscr{M} \to \mathscr{M}'$ such that $d\phi(p) = L$.* □

Definition 1.1 *A geodesically complete Lorentzian manifold with constant sectional curvature is called a* space form.

It follows from Theorem 1.2 that any simply connected space form is uniquely determined by Λ. We list below such solutions.

$\Lambda > 0$ – de Sitter spacetimes (notation: $dS_d(\Lambda)$), given by the spatially compact hyperboloid

$$dS_d(\Lambda) \doteq \{X \in \mathbb{R}^{1,d} : \eta(X, X) = R^2\},\ R = \sqrt{\frac{(d-1)(d-2)}{2\Lambda}}. \tag{1.4}$$

Its isometry group is the de Sitter *group* $O(1, d)$, with component connected to identity [3] $SO_e(1, d)$.

[2] The denominator in (1.2) denotes the square of the Lorentzian area of the parallelogram with sides X and Y.

[3] For orientable and time orientable Lorentzian manifolds, this is given by the *proper* (preserving the orientation of the volume element, i.e., whose tangent map has determinant one) and *orthochronous* (preserving time orientation) isometries.

$\Lambda = 0$ – MINKOWSKI spacetime $\mathbb{R}^{1,d-1}$. Its isometry group is the POINCARÉ *group* $O(1,d-1) \ltimes \mathbb{R}^d$, with component connected to identity $SO_e(1,d-1) \ltimes \mathbb{R}^d \doteq \mathscr{P}^{\uparrow}_{+,d}$.

$\Lambda < 0$ – *anti*-DE SITTER spacetimes (notation: $AdS_d(\Lambda)$), given by the universal covering (see Appendix D, Definition D.3, page 155) of the timelike compact hyperboloid

$$(1.5) AdS_d(\Lambda) \doteq \{X \in \mathbb{R}^{2,d-1} : (X^0)^2 - (X^1)^2 - \cdots - (X^{d-1})^2 + (X^d)^2 = R^2\},$$
$$R = \sqrt{-\tfrac{(d-1)(d-2)}{2\Lambda}}.$$

Its isometry group is the *anti*-DE SITTER *group*, given in the fundamental domain (1.5) by $O(2,d-1)$, with component connected to identity $SO_e(2,d-1)$.

REMARK 1.2 *In a slight abuse of notation, we shall omit in general the cosmological constant when employing a notation above for the different simply connected space forms (i.e., we'll write $(A)dS_d$ instead of $(A)dS_d(\Lambda)$), whenever such a practice doesn't cause confusion. Unlike a recurrent practice in literature, we don't identify AdS_d with the fundamental domain (1.5), for the latter has a minimal role throughout this work, besides causing unnecessary technical complications.*

All solutions above are *maximally symmetric:* the LIE algebra of KILLING fields of each space form has maximal dimension $\frac{d(d+1)}{2}$ (conversely, any connected Lorentzian manifold whose LIE algebra of KILLING fields has maximal dimension possesses constant sectional curvature). By a standard result [O'N83], every KILLING field uniquely extends to a one-parameter group of isometries, whence it follows that the LIE group of isometries of a space form has also dimension $\frac{d(d+1)}{2}$. Moreover: let p, q be two points of a space form $\mathscr{M}$. We can connect them by a finite sequence of geodesic segments $\gamma_i, i = 1, \ldots, k$ with affine parameters $\lambda_i \in [0,1]$ – in this case, $\gamma_i(1) = \gamma_{i+1}(0)$, $\gamma_1(0) = p$ and $\gamma_k(1) = q$ – by covering any curve segment linking p to q with a finite number of normal neighbourhoods. Consider the middle point $\bar{p}_i$ of γ_i, and the isometry $L_i : T_{\bar{p}_i}\mathscr{M} \to T_{\bar{p}_i}\mathscr{M}$ given by $-\mathrm{id}_{T_{\bar{p}_i}\mathscr{M}}$. By Theorem 1.1, each L_i determines an isometry $\phi_i : \mathscr{M} \to \mathscr{M}$, in this case satisfying $\phi_i(\gamma_i(0)) = \gamma_i(1)$. Finally, $\phi = \phi_k \circ \cdots \circ \phi_1$ is an isometry linking p to q. That is, any two points p, q are connected by an isometry, i.e., *any space form is a homogeneous space* (i.e., the action of the isometry group is *transitive*). In particular, the isometry group acts in space forms *without fixed points*, i.e., given any point p, there exists an isometry ϕ such that $\phi(p) \neq p$.

1.2 Geometry of AdS_d: boundary conditions at infinity

We can write a global system of spherically symmetric spatial coordinates for AdS_d. Consider $(\tau, r, \mathbf{e}) \in \mathbb{R} \times \bar{\mathbb{R}}_+ \times S^{d-2}$. Writing

$$\left\{ \begin{array}{ll} X^0 = \sqrt{R^2 + r^2} \sin t & \\ \mathbf{X} \ = r\mathbf{e} & \\ X^d = \sqrt{R^2 + r^2} \cos t & \end{array} \right. , \ R \text{ given by (1.5)}, \tag{1.6}$$

the AdS_d metric becomes

$$ds^2 = -R^2 \left(1 + \frac{r^2}{R^2}\right) dt^2 + \left(1 + \frac{r^2}{R^2}\right)^{-1} dr^2 + r^2 d\Omega^2_{d-2}. \tag{1.7}$$

The fundamental domain is obtained by the restriction $-\pi < t \leq \pi$. As the covering is, thus, obtained by "unwrapping" the time coordinate τ, we shall denominate the global chart (1.6–1.7) the *the covering chart*. A coordinate system defined only in the domain $\{X : X^{d-1} + X^d > 0\}$ (called Poincaré *(fundamental) domain*), but far more convenient for the analysis of several geometrical aspects of AdS_d, is given by $(x^\mu, z) \in \mathbb{R}^{1,d-2} \times \mathbb{R}_+$, if we write

$$\left\{ \begin{array}{ll} X^\mu & = \frac{R}{z} x^\mu \ (\mu = 0, \ldots, d-2) \\ X^{d-1} & = R \left(\frac{1-z^2}{2z} + \frac{1}{2z} x_\mu x^\mu \right) \\ X^d & = R \left(\frac{1+z^2}{2z} - \frac{1}{2z} x_\mu x^\mu \right) \end{array} \right. \tag{1.8}$$

In this case, (1.7) acquires the following form:

$$ds^2 = \frac{R^2}{z^2} \left(\eta_{\mu\nu} dx^\mu dx^\nu + dz^2 \right). \tag{1.9}$$

This chart is denominated *horocyclic* or a Poincaré *chart*. One can see by formula (1.9) that each timelike hypersurface in the Poincaré domain given by $z = const.$ is conformal to $\mathbb{R}^{1,d-2}$ by a factor

$$(X^{d-1} + X^d)^2 = \frac{R^2}{z^2}. \tag{1.10}$$

Hence, the Poincaré domain corresponds to the half $z > 0$ of $\mathbb{R}^{1,d-1} \ni (x^\mu, x^d - 1 \doteq z)$, up to the conformal factor (1.10). We can obtain the action of several subgroups of $SO_e(2, d-1)$ on AdS_d from the subgroup of (the component connected to identity of) the conformal group $SO_e(2, d)$ of $\mathbb{R}^{1,d-1}$ that individually preserves each half $\{(x^\mu, x^{d-1}) : x^{d-1} \gtrless 0\}$ by using this parametrization, as follows:

- POINCARÉ *subgroup:*

$$(z, x^\mu) \mapsto (z, \Lambda^\mu_\nu x^\nu + a^\mu),\ \Lambda \in SO_e(1, d-2),\ a \in \mathbb{R}_d. \tag{1.11}$$

  This subgroup preserves the z =constant hypersurfaces, and acts on each one as the POINCARÉ group acts in $\mathbb{R}^{1,d-2}$.

- *Dilation subgroup:*

$$(z, x^\mu) \mapsto (\lambda z, \lambda x^\mu),\ \lambda \in \mathbb{R}_+. \tag{1.12}$$

The remaining isometries, which do not preserve the fundamental POINCARÉ domain, will be presented in Subsection 1.2.1.

1.2.1 Structure of conformal infinity

Let us consider the covering parametrization (1.6). We'll show now that AdS_d possesses a conformal infinity in the sense of Definition A.3. Applying the change of variables

$$\begin{gathered}\tau = t,\ u = 2\left(\sqrt{1+\tfrac{r^2}{R^2}}\right) - \tfrac{r}{R},\ dr = -R\left(\tfrac{1}{u^2} + \tfrac{1}{4}\right) du \\ r \in [0, +\infty) \leftrightarrow u \in (0, 2],\end{gathered} \tag{1.13}$$

the metric (1.7) becomes

$$ds^2 = \frac{R^2}{u^2}\left[-\left(1+\frac{u^2}{4}\right)^2 dt^2 + du^2 + \left(1-\frac{u^2}{4}\right)^2 d\Omega^2_{d-2}\right]. \tag{1.14}$$

The spatial infinity of AdS_d is obtained by taking $r \to \infty$, which corresponds to the limit $u \searrow 0$. The metric of the conformal completion $d\bar{s}^2 = u^2 ds^2$, in this limit, becomes

$$d\bar{s}^2 = R^2(-dt^2 + d\Omega^2_{d-2} + du^2) = R^2(du^2 + ds^2_0), \tag{1.15}$$

where ds^2_0 is the metric of EINSTEIN*'s static universe* in $(d-1)$ dimensions (ESU_{d-1}), which is isometric to $\mathbb{R} \times S^{d-2}$. In order to see the conformal boundary of the fundamental domain, let us return to coordinates τ, r; one notices that the boundary is then conformal to $\mathbb{R}^{1,d-2}$ by a factor $(X^{d-1} + X^d)^{-2}$. Taking projective (DIRAC-WEYL) coordinates, we have:

$$x^\mu = \frac{X^\mu}{X^{d-1} + X^d} \overset{r\to\infty}{\longrightarrow} \begin{cases} x^0 = \frac{\sin\tau}{\cos\tau + \mathrm{e}^d} \\ \mathbf{x} = \frac{\mathbf{e}}{\cos\tau + \mathrm{e}^d} \end{cases}. \tag{1.16}$$

More precisely, the conformal boundary of the fundamental domain of AdS_d corresponds to the conformal compactification of $\mathbb{R}^{1,d-2}$ (notation: $c\mathbb{R}^{1,d-2}$).

ESU_{d-1}, contrary to $c\mathbb{R}^{1,d-2}$, doesn't suffer from causality conflicts due to the action of the conformal group of $\mathbb{R}^{1,d-2}$, which extends to the universal covering [4] $\widetilde{SO}_e(2,d-1)$. Its global causal ordering is given by [LM75]

$$(\tau,\mathrm{e}) \ll_{\mathscr{I}} (\tau',\mathrm{e}') \text{ if and only if } \tau-\tau' > 2\mathrm{Arccos}(\mathrm{e}.\mathrm{e}'), \tag{1.17}$$

where $\mathrm{Arccos}(a)$ corresponds to the principal branch of $\cos^{-1}(a)$ $(-\pi < a \leq \pi)$. In terms of the Poincaré parametrization (1.8), each z =const. hypersurface corresponds, in projective coordinates, exactly to $\mathbb{R}^{1,d-2}$, a result which extends to $z=0$. Thus, conversely to (1.16), we can obtain once more the global chart of ESU_{d-1} by explicitly writing the conformal embedding of $\mathbb{R}^{1,d-2}$ into the latter. Namely, rewriting η in "spatially" spherical coordinates

$$\eta = -(dx^0)^2 + d\mathbf{x}.d\mathbf{x} = -(dx^0)^2 + dr^2 + r^2 d\Omega^2_{d-3}$$

and moving to advanced (v) and retarded (u) radial coordinates in the light cone

$$\left.\begin{array}{l} v \doteq t+r \\ u \doteq t-r \end{array}\right\} \Rightarrow \eta = -du\,dv + \frac{1}{4}(v-u)^2 d\Omega^2_{d-3},$$

we see that, under the choice of conformal factor $\Omega^2 = \frac{4}{(1+u^2)(1+v^2)}$ and the coordinate change $(u,v) \mapsto (U,V)$ (given by $U = 2\mathrm{arctan}u$ e $V = 2\mathrm{arctan}v$) and $(U,V) \mapsto (T \doteq \frac{V+U}{2}, R \doteq \frac{V-U}{2})$, it follows that

$$ds^2_0 = \Omega^2\eta = -dT^2 + dR^2 + \sin^2 R d\Omega^2_{d-3} = -dT^2 + d\Omega^2_{d-2},$$

according to (1.15). That is, the coordinate change $(x^\mu) \mapsto (T, \theta^1 \doteq R, \theta^2, \ldots, \theta^{d-2})$ corresponds from the passive viewpoint to what, from the active viewpoint, is the conformal embedding $\mathbb{R}^{1,d-2}$ into ESU_{d-1} (U and V are the light-cone radial coordinates in ESU_{d-1}). One should also notice, in the light of these considerations, that AdS_d, expressed in terms of the projective coordinates (1.16) properly extended to the covering, corresponds to half of ESU_d. The Poincaré domain, on its turn, corresponds to the image of the conformal embedding of the half $x^{d-1} > 0$ of $\mathbb{R}^{1,d-1}$ into ESU_d, and its infinity $(x^{d-1}=0)$, to the image of the conformal embedding of $\mathbb{R}^{1,d-2}$ into ESU_{d-1}. The above analysis is pictorially synthesized in Figure 1.1.

Starting from what has been said above, it's immediate to identify the remaining subgroup of $\widetilde{SO}_e(2,d-1)$. The conjugation of the subgroup of translations along x^μ (see (1.11), page 7) by the *relativistic ray inversion* map in $\mathbb{R}^{1,d-1}$

$$I : (x^\mu, x^{d-1}) = (x^\mu, z) \mapsto -\frac{1}{z^2 + x_\nu x^\nu}(x^\mu, z),\ x_\nu x^\nu = \eta_{\rho\sigma}x^\rho x^\sigma$$

[4] *Modulo* the group $\mathbb{Z}_2$ of reflections across the spatial infinity of $c\mathbb{R}^{1,d-2}$.



Figure 1.1: **(a)** Fundamental domain of AdS_d (red), given by the quotient modulo $\mathbb{Z}$, where the spatial infinity is represented in blue and the (fundamental) POINCARÉ domain is the upper part of the dotted cut; **(b)** Image of the conformal embedding of AdS_d into ESU_d (red). The infinity $\mathscr{I}$ of AdS_d is represented in blue and the POINCARÉ domain $\mathscr{P}oi(p)$ is delimited by the dotted curves; **(c)** $\mathscr{P}oi(p)$ (side view) delimited by the dotted lines in red, with $\mathscr{I}$ represented in blue; **(d)** Visualization of $\mathscr{I}$ as ESU_{d-1} (red) with the MINKOWSKI domain $\mathscr{M}in(p)$, corresponding to the image of the conformal embedding of $\mathbb{R}^{1,d-2}$ into ESU_{d-1}, in blue.

results in the *special conformal transformations* (also called *conformal translations*)

$$(x^\mu, z) \mapsto I \circ (. + b^\mu) \circ I(x^\mu, z) = \frac{1}{1 - 2b_\mu x^\mu + b^2(x^2 + z^2)}. \tag{1.18}$$
$$.(x^\mu - b^\mu(x^2 + z^2), z),\ b \in \mathbb{R}^{1,d-2} \ni a^\mu,\ a^2 := a_\mu a^\mu.$$

Such transformations, if conjugated by adequate elements of the subgroups (1.11–1.12) and extended to the parameter space of $\widetilde{SO}_e(2, d-1)$, no longer preserve the Poincaré domain, but preserve the infinity of AdS_d as a whole. For $z = 0$, (1.18) reduces to the special conformal transformations in $\mathbb{R}^{1,d-2}$. The subgroups given by (1.11) and (1.12) in the projective coordinates x^μ correspond, respectively, to the actions of the Poincaré and dilation subgroups on $\mathbb{R}^{1,d-2}$. Summing up, the isometry group of the fundamental domain of AdS_d (resp. AdS_d) corresponds precisely to the conformal group of $(c)\mathbb{R}^{1,d-2}$ (resp. ESU_{d-1}).

1.2.2 Wedges in AdS and diamonds in the boundary

Let us consider, in AdS_d, the following causally complete region, called *(standard) wedge* (notation: $\mathscr{W}_0$):

$$\mathscr{W}_0 \;=\; \{X \in \mathbb{R}^{2,d-1} : \eta(X,X) = -R^2 \text{ and } X^{d-1} > |X^0|\}, \tag{1.19}$$

or, in Poincaré coordinates (z, x^μ):

$$\mathscr{W}_0 = \{(z, x^\mu) : \sqrt{z^2 + \mathbf{x}\cdot\mathbf{x}} < 1 - |x^0|, z < 1\}. \tag{1.20}$$

Taking the limit $z \to 0$, one notices that the intersection of $\mathscr{W}_0$ with the conformal boundary (notation: $\mathscr{K}_0$) is given by

$$\mathscr{K}_0 \;=\; \{x^\mu : |\mathbf{x}| < 1 - |x^0|\}, \tag{1.21}$$

that is, $\mathscr{K}_0$ corresponds to a *diamond* (with spatial radius equal to 1) in $\mathbb{R}^{1,d-2}$.

A wedge in AdS_d is the causal completion of the orbit of a uniformly accelerating observer. By formula 1.19, we see that such orbits are given by the restriction to the fundamental domain of AdS_d of the boosts in the $X^0 - X^{d-1}$ plane:

$$\begin{aligned} X^0 &\mapsto X^0 \cosh\lambda + X^{d-1}\sinh\lambda; \\ X^{d-1} &\mapsto X^0 \sinh\lambda + X^{d-1}\cosh\lambda; \\ \mathbf{X} &\mapsto \mathbf{X};\ X^d \mapsto X^d,\ \lambda \in \mathbb{R}. \end{aligned} \tag{1.22}$$

In Poincaré coordinates, it's easier to visualize the transformation in the conformal embedding of AdS_d into ESU_d. $\mathscr{W}_0$ is the $x^{d-1} > 0$ half of a diamond contained in the conformal embedding of $\mathbb{R}^{1,d-1}$ into ESU_d, with vertices in $\mathscr{I}$. (1.22) corresponds to

the unique one-parameter subgroup of $\widetilde{SO}_e(2,d-1)$ which preserves this diamond. The former's action is symmetric under rotations around the axis determined by the vertices, and, restricted to the $x^0 - z$ plane, becomes

$$z_\pm \mapsto z_\pm(\lambda) \doteq \frac{(1+z_\pm) - e^{-\lambda}(1-z_\pm)}{(1+z_\pm) + e^{-\lambda}(1-z_\pm)},\ \lambda \in \mathbb{R}, \tag{1.23}$$

where we've employed the light-cone coordinates $z_\pm \doteq x^0 \pm z$. The corresponding isotropy subgroup for $\mathscr{D}_0$ is obtained, say, in the $x_0 - x_1$ plane by substituting x^1 for z and the light-cone coordinates $x_\pm \doteq x^0 \pm x^1$ for $z_\pm$ in (1.23). The one-parameter subgroup of $SO_e(2,d-1)$ given by (1.23) corresponds to the conjugation $(x,z) \mapsto (K^{-1}\circ(e^\lambda .)\circ K)(x,z)$ of the dilation $(x,z) \mapsto (e^\lambda x, e^\lambda z)$ by the conformal transformation K in $\mathbb{R}^{1,d-1}$ given by the composition

$$\mathscr{W}_0 \ni (x,z) \doteq (x^0, \mathbf{x}, z) \mapsto K(x,z) \doteq I(x^0 - 1, \mathbf{x}, z) - \left(\frac{1}{2}, \mathbf{0}, 0\right),$$

where I is the relativistic ray inversion map. In this form, the rotational symmetry of the formula (1.23) around the x^0 axis, invoked above, becomes evident.

Finally, as a prelude to Section 1.3, we shall pass to proceed in a coordinate-free manner. Denoting the vertices of $\mathscr{D}_0$ by $p_0 = (-1, \mathbf{0})$ and $q_0 = (1, \mathbf{0})$, we see that, if $\overline{AdS_d}$ is the conformal closure of AdS_d, then $(I^-(p_0, \overline{AdS_d}) \cap I^+(q_0, \overline{AdS_d})) \cap AdS_d = \mathscr{W}_0$ and $(I^-(p_0, \overline{AdS_d}) \cap I^+(q_0, \overline{AdS_d})) \cap \mathscr{I} = (I^-(p_0, \mathscr{I}) \cap I^+(q_0, \mathscr{I})) = \mathscr{D}_0$. More in general, let us define for $p, q \in \mathbb{R}^{1,d-2}, p \ll_{\mathscr{I}} q$ the *wedge*

$$\mathscr{W}_{p,q} \doteq (I^-(p, \overline{AdS_d}) \cap I^+(q, \overline{AdS_d})) \cap AdS_d \tag{1.24}$$

and the *diamond*

$$\mathscr{D}_{p,q} \doteq (I^-(p, \overline{AdS_d}) \cap I^+(q, \overline{AdS_d})) \cap \mathscr{I} = I^-(p, \mathscr{I}) \cap I^+(q, \mathscr{I}) \tag{1.25}$$

associated to p, q. In particular, $\mathscr{W}_0 = \mathscr{W}_{p_0,q_0}$ and $\mathscr{D}_0 = \mathscr{D}_{p_0,q_0}$. Let us denote by $u^\lambda_{p,q}$ the one-parameter subgroup of isometries of AdS_d which preserve $\mathscr{W}_{p,q}$ and $\mathscr{D}_{p,q}$, given by (1.23) in the case $p = p_0$, $q = q_0$. This subgroup is given by

$$\begin{aligned} u^\lambda_{p,q}(x,z) &= K^{-1}_{p,q}(e^\lambda K_{p,q}(x,z)), \text{ where} \\ K_{p,q}(x,z) &\doteq K\left(\Lambda_{p,q}\left(x - \frac{x(p)+x(q)}{2}\right), z\right), \end{aligned} \tag{1.26}$$

where $\Lambda_{p,q}$ is the LORENTZ boost around the origin which makes the direction $\overrightarrow{x(p)x(q)}$ parallel to the x^0 axis, and K is the map used to define the isotropy subgroup (1.23)

of $\mathscr{W}_0$ above. These one-parameter subgroups will play a key role in Chapters 2 and 4.

Let us make now the description of the image of the conformal embedding of $\mathbb{R}^{1,d-2}$ into ESU_{d-1} independent from the point chosen to represent the spatial infinity of the former. Let $p \in \mathscr{M}in(r)$. All null geodesics emanating from p will focus at a single point of $\mathscr{I}$, which constitutes the future endpoint of all achronal null generators of $\partial I^+(p, \mathscr{I})$. This point is denominated *antipodal* of p, denoted by $\bar{p}$. The antipodal of p has the following properties:

$$\partial I^+(p, \mathscr{I}) \smallsetminus \{p\} = \partial I^-(\bar{p}, \mathscr{I}) \smallsetminus \{\bar{p}\}; \tag{1.27}$$

$$\partial I^+(p, \overline{AdS_d}) \smallsetminus \{p\} = \partial I^-(\bar{p}, \overline{AdS_d}) \smallsetminus \{\bar{p}\}. \tag{1.28}$$



Figure 1.2: **(a)** Correspondence between bulk wedges and boundary diamonds. **(b)** A Minkowski domain and the corresponding Poincaré domain.

Let us define $\mathscr{M}in(p) \doteq \mathscr{D}_{p,\bar{\bar{p}}}$, the Minkowski *domain* to the future of $p \in \mathscr{I}$. This region corresponds to the conformal embedding of $\mathbb{R}^{1,d-2}$ into $\mathscr{I}$ such that p is the future timelike infinity of $\mathbb{R}^{1,d-2}$. $\mathscr{P}oi(p) \doteq \mathscr{W}_{p,\bar{\bar{p}}}$ corresponds to the domain of a Poincaré chart in AdS_d, hence called the Poincaré *domain* to the future of p. We shall extend the definitions (1.24) of a wedge and (1.25) of a diamond to all pairs of points $p \ll_{\mathscr{I}} q$ such that $p, q \in \overline{\mathscr{M}in(r)}$ for some $r \in \mathscr{I}$, with the same notation.

An equivalent way of characterizing wedges $\mathscr{W}_{p,q}$ and diamonds $\mathscr{D}_{p,q}$ such that $p, q \in \mathscr{M}in(r)$ for some $r \in \mathscr{I}$ is the following: as ESU_{d-1} is globally hyperbolic, it follows that $\mathscr{D}_{p,q}$ is relatively compact and, thus, globally hyperbolic as well. It's possible, then, to distinguish three possible situations:

1. *There is no $r \in \mathscr{I}$ such that $p, q \in \overline{\mathscr{M}in(r)}$: in this case, let us take $r = p$; it follows that $\bar{r} = \bar{p} \ll_{\mathscr{I}} q$ and, hence, $\mathscr{D}_{p,q}$ contains a* Cauchy *surface for $\mathscr{I}$. In particular, the* Cauchy *surfaces of $\mathscr{D}_{p,q}$ are* compact *and, more important,* non contractible *(for the definition of contractibility, see Section D.1, page 153).*

2. *$p, q \in \mathscr{M}in(r)$ for some $r \in \mathscr{I}$: as here $\mathscr{D}_{p,q}$ can be understood as a diamond in $\mathscr{M}in(r)$, the* Cauchy *surfaces of $\mathscr{D}_{p,q}$ are* non compact *and* contractible.

3. *$p, q \in \partial\mathscr{M}in(r)$ for some $r \in \mathscr{I}$: this borderline case includes the* Minkowski *domains themselves, and, by definition, $\mathscr{D}_{p,q}$ can then be obtained as the limit $\mathscr{D}_{p,q} = \lim_{n\to\infty} \mathscr{D}_{p,q_n} = \bigcup_{n=1}^{\infty} \mathscr{D}_{p,q_n}$, where once more we choose $r = p$ and $\{q_n\}_{n\in\mathbb{Z}_+} \subset \mathscr{M}in(r)$ is a sequence of points such that $q_n \overset{n\to\infty}{\longrightarrow} q$ and $q_n \ll_{\mathscr{I}} q_{n+1} \ll_{\mathscr{I}} q$ for all n. In particular, $\mathscr{D}_{p,q_n} \in \mathscr{M}in(r_n)$, where $r_n \ll_{\mathscr{I}} r$ is sufficiently close to $r = p$. The* Cauchy *surfaces of $\mathscr{D}_{p,q}$ are also noncompact and contractible.*

We emphasize that by no means can case 1 be incorporated by any limit of an increasing sequence of diamonds lying in case 2, as in case 3. Summing up, the collection of diamonds that interest us is the one composed of diamonds with contractible Cauchy surface*. The diamonds within case 2 are, in a certain sense, "dense" in this collection – given any $\mathscr{D}_{p,q}$ with contractible* Cauchy *surface and any $q' \ll_{\mathscr{I}} q$, no matter how close to q, we have that $\mathscr{D}_{p,q'}$ is a diamond within case 2.*

The structure of ESU_{d-1} yet implies the following facts of great importance to us:

1. From (1.27) and (1.28), it follows that
   (1.29)
   $$\mathscr{W}_{q,\bar{p}} = (\mathscr{W}_{p,\bar{q}})'_{AdS_d} \text{ and } \mathscr{D}_{q,\bar{p}} = (\mathscr{D}_{p,\bar{q}})'_{\mathscr{I}},\ \forall p, q \in \mathscr{M}(r) \text{ for some } r \text{ and } p \ll_{\mathscr{I}} \bar{q},$$
   and, as a consequence,

2. $$\mathscr{M}in(r) = (\{\bar{r}\})'_{\mathscr{I}} = (\{\bar{r}\})'_{\overline{AdS_d}} \cap \mathscr{I},\ \mathscr{P}oi(r) = (\{\bar{r}\})'_{\overline{AdS_d}} \cap AdS_d. \tag{1.30}$$

3. The bijection
   (1.31)
   $$\rho_{AdS_d} : \mathscr{W}_{p,q} \mapsto \rho_{AdS_d}(\mathscr{W}_{p,q}) \doteq \mathscr{D}_{p,q},\ p, q \in \mathscr{M}in(r) \text{ for some } r \text{ and } p \ll_{\mathscr{I}} q,$$
   denominated Rehren *bijection*, preserves causal complements.

4. The action of $\widetilde{SO}_e(2,d-1)$ on $\mathscr{I}$ is transitive and preserves the causal structure. Hence, it acts transitively on the set of diamonds. As $SO_e(2,d-1)$ is the conformal group of Minkowski spacetime, it follows that the former acts transitively on the collection of the diamonds contained in some Minkowski domain.

5. As $\widetilde{SO}_e(2,d-1)$ also preserves the causal structure of AdS_d, it also follows that this group acts transitively on the set of wedges.

6. Every element of $\widetilde{SO}_e(2,d-1)$ can be obtained from a finite sequence of isometries of the form (1.26) for a suitable family of diamonds / wedges within cases 2 and 3 of Remark 1.3, page 13.

The above construction, which makes use only of the causal relations in the conformal closure, constitutes the geometrical framework of Rehren duality [Reh00], and was employed in this coordinate-free form by Bousso and Randall [BR02] for studying qualitative aspects of the AdS/CFT correspondence.

1.3 AAdS spacetimes

Having scrutinized with sufficient detail the structure of AdS_d, we shall now proceed with the general case of our interest. Following the nomenclature of Appendix A, consider stably causal spacetimes $(\mathscr{M},g)$ endowed with conformal completion $(\overline{\mathscr{M}},\bar{g})$, conformal infinity $(\mathscr{I},\bar{g}^{(0)})$ and conformal factor z satisfying the conditions of Definition A.3 (page 117).

Definition 1.4 *We say that $(\mathscr{M},g)$ is an* AdS-type *spacetime id $(\mathscr{I},\bar{g}^{(0)})$ is a $(d-1)$-dimensional Lorentzian manifold (or, equivalently, the covector normal to $\mathscr{I}$, $dz\restriction_{z=0}$, is spacelike with respect to $\bar{g}$) and diffeomorphic to $\mathbb{R}\times S^{d-2}$. If, beyond that, $\bar{g}^{(0)}$ belongs to the conformal class of the metric of ESU_{d-1}, $(\mathscr{M},g)$ is said to be* asymptotically AdS *(AAdS).*

Finally, a spacetime $(\mathscr{M},g)$ endowed with conformal infinity $(\mathscr{I},\bar{g}^{(0)})$ is said to be locally AdS-type *(resp.* locally AAdS*) if any $p\in\mathscr{I}$ has an open neighbourhood $\overline{\mathscr{U}}$ in the conformal completion $(\overline{\mathscr{M}},\bar{g})$ such that $(\overline{\mathscr{U}},\bar{g}\restriction_{\overline{\mathscr{U}}})$ is isometric to a neighbourhood $\overline{\mathscr{V}}$ of $q\in\mathscr{J}$, where $\mathscr{J}$ is the conformal infinity of an AdS-type (resp. AAdS) spacetime.*

The definition of locally AdS-type ad locally AAdS spacetimes comprises the examples of black holes negative cosmological constant (Schwarzschild-AdS, Reissner-Nordström-AdS, Kerr-Newman-AdS, etc.), not contemplated by the definition of AdS-type and AAdS spacetimes.

With Definition 1.4 at hand, we define in AAdS spacetimes the antipodal $\bar{p}$ and the MINKOWSKI domain $\mathscr{M}in(p)$ to the future of $p \in \mathscr{I}$ exactly as in AdS, for these definitions don't refer to the bulk geometry. More in general, we have the

DEFINITION 1.5 *Let $(\mathscr{M}, g)$ be a locally AdS-type spacetime with conformal infinity $(\mathscr{I}, \bar{g}^{(0)})$, and $p \ll_{\mathscr{I}} q \in \mathscr{I}$ such that $I^+(p, \overline{\mathscr{M}}) \cap I^-(q, \overline{\mathscr{M}})$ is relatively compact (hence, globally hyperbolic) and $I^+(p, \mathscr{I}) \cap I^-(q, \mathscr{I})$ possesses contractible* CAUCHY *surfaces. Let us denote the collection of pairs $(p, q) \in \mathscr{I}^2$ satisfying these conditions by $\mathscr{D}(\mathscr{I})$.*

The (bulk) wedge *associated to p, q is the region*

$$\mathscr{W}_{p,q} = I^+(p, \overline{\mathscr{M}}) \cap I^-(q, \overline{\mathscr{M}}) \cap \mathscr{M},$$

and the (boundary) diamond *associated to p, q, the region*

$$\mathscr{D}_{p,q} = I^+(p, \mathscr{I}) \cap I^-(q, \mathscr{I}).$$

The past horizon $\partial_- \mathscr{W}_{p,q}$ *(resp.* future horizon $\partial_+ \mathscr{W}_{p,q}$*) of $\mathscr{W}_{p,q}$ is given by*

$$\partial_- \mathscr{W}_{p,q} \doteq (\partial I^+(p, \overline{\mathscr{M}}) \setminus \{p\}) \cap I^-(q, \overline{\mathscr{M}}) \cap \mathscr{M},$$

$$\partial_+ \mathscr{W}_{p,q} \doteq (\partial I^-(q, \overline{\mathscr{M}}) \setminus \{q\}) \cap I^+(p, \overline{\mathscr{M}}) \cap \mathscr{M},$$

and the past horizon $\partial_- \mathscr{D}_{p,q}$ *(resp.* future horizon $\partial_+ \mathscr{D}_{p,q}$*) of $\mathscr{D}_{p,q}$, by*

$$\partial_- \mathscr{D}_{p,q} \doteq (\partial I^+(p, \mathscr{I}) \setminus \{p\}) \cap I^-(q, \mathscr{I}),\ \partial_+ \mathscr{D}_{p,q} \doteq (\partial I^-(q, \mathscr{I}) \setminus \{q\}) \cap I^+(p, \mathscr{I}).$$

We denote by $\mathscr{W}(\mathscr{M}, g) \doteq \{\mathscr{W}_{p,q} : (p, q) \in \mathscr{D}(\mathscr{I})\}$ the collection of wedges in $(\mathscr{M}, g)$, and $\mathscr{D}(\mathscr{M}, g) \doteq \{\mathscr{D}_{p,q} : (p, q) \in \mathscr{D}(\mathscr{I})\}$, the collection of diamonds in $(\mathscr{I}, \bar{g}^{(0)})$. The REHREN bijection *associated to $(\mathscr{M}, g)$ is given by*

$$\begin{aligned} \rho_{(\mathscr{M},g)} : \mathscr{W}(\mathscr{M}, g) &\longrightarrow \mathscr{D}(\mathscr{M}, g) \\ \mathscr{W}_{p,q} &\mapsto \rho_{(\mathscr{M},g)}(\mathscr{W}_{p,q}) \doteq \mathscr{D}_{p,q}. \end{aligned} \tag{1.32}$$

Notice that it's Remark 1.3 which allowed us to naturally extend the definition of wedges and diamonds to locally AdS-type spacetimes. If $(\mathscr{M}, g)$ is an AAdS spacetime, the POINCARÉ domain to the future of $r \in \mathscr{I}$ coincides, as in the case of MINKOWSKI domains, with the definition in the AdS case: $\mathscr{P}oi(r) \doteq \mathscr{W}_{r,\bar{r}}$.

The wedges in a locally AdS-type spacetime cover precisely the region

$$I^-(\mathscr{I}, \overline{\mathscr{M}}) \cap I^+(\mathscr{I}, \overline{\mathscr{M}}) \cap \mathscr{M} \doteq \mathscr{W}(\mathscr{I}),$$

called *domain of outer communications* of $(\mathscr{M}, g)$, which is identified in several examples with the exterior of AAdS black holes.

In Subsection 1.3.1 below, we shall study the causal, topological and localization properties of $\rho_{(\mathscr{M},g)}$, in terms of the null geodesics of AAdS spacetimes.

1.3.1 Global geometry and null geodesics

1.3.1.1 Causality (bulk to boundary)

Einstein's static universe $\mathscr{I}$ is globally hyperbolic, and the orbits generated by the subgroup of $\widetilde{SO}_e(2, d-1)$ corresponding to time translations in AdS_d (rotation of the points of the hyperquadric (1.5), page 5, in the $X^0 - X^d$ plane) correspond to the lines parallel to the cylinder [5] $\mathbb{R} \times S^{d-2}$, constituting timelike geodesics which maximize the proper time between its points. Moreover, any point in $\mathscr{I}$ belong to a unique geodesic of this family, and we can choose a common parametrization to all these geodesics (denominated *time generators* of $\mathscr{I}$) such that the union of the points corresponding to the same value of the affine parameter constitute a Cauchy surface.

With this, we can define the *(gravitational) time delay* of a complete null geodesic γ in $\mathscr{M}$, with endpoints in $\mathscr{I}$: *grosso modo*, it represents how much the future endpoint of γ moves away from the antipodal of the past endpoint of γ, which, in the AdS case, *is* the future endpoint.

First, we'll prove that, given certain extra conditions, the gravitational time delay of inextendible null geodesics is always *positive* (see Figure 1.3 for an illustration of the result):

Theorem 1.3 (positive gravitational time delay) *Let $(\mathscr{M}, g)$ be an AAdS spacetime satisfying the following* global focusing *hypothesis: any inextendible null geodesic possesses a pair of conjugate points (see in Appendix A the discussion that follows equation (A.14), page 110). Then, given $p \in \mathscr{I}$, every inextendible null geodesic in $\mathscr{M}$ with past endpoint p has, if any, future endpoint in $I^+(p, \mathscr{I})$. A similar result is valid if we exchange future with past, and $I^+(p, \mathscr{I})$ with $I^-(p, \mathscr{I})$.*

Proof. Initially, we shall prove two Lemmata:

Lemma 1.4 (Absence of causal shortcuts) *Let $p, p' \in \mathscr{I}$. If $p \perp_{\mathscr{I}} p'$, then there cannot exist a causal curve in $(\overline{\mathscr{M}}, \bar{g})$ linking p to p'.*

Proof. Suppose that $p' >_{\overline{\mathscr{M}}} p$ (the opposite case is treated in an analogous way). We'll prove that the gravitational time delay implied by the hypotheses of the Theorem contradicts the causal disjointness of p and p' with respect to $\mathscr{I}$. Denote by $T(p')$ the unique time generator of $(\mathscr{I}, \bar{g}^{(0)})$ containing p'.

Notice that $\partial I^+(p, \mathscr{I}) \doteq \Sigma$ is a closed, achronal hypersurface, which cuts $(\mathscr{I}, \bar{g}^{(0)})$ in two disjoint open sets $I^+(p, \mathscr{I}) \doteq A$ and $\mathscr{I} \setminus \overline{I^+(p, \mathscr{I})}$

[5] The generator of this subgroup corresponds to $1/2(P^0 + K^0)$ in a Minkowski domain, where P^0 is the generator of translations along the direction x^0, and K^0 the generator of special conformal transformations along the direction x^0.



Figure 1.3: Gravitational time delay. The null geodesics which emanate from $p \in \mathscr{I}$ and remain in $\mathscr{I}$ (full lines) focus at the antipodal $\bar{p}$. Whereas in AdS_d a null geodesic emanating from p and traversing the bulk would also have $\bar{p}$ as its future endpoint (dotted line), in an AAdS spacetime satisfying the hypothesis of global focusing of Theorem 1.3 such a geodesic has future endpoint $q \in I^+(p, \mathscr{I})$ (dashed/dotted line).

$\doteq B$, and intersects each time generator of $\mathscr{I}$ precisely once, for any time generator has points in $I^+(p, \mathscr{I})$ and $I^-(\bar{p}, \mathscr{I})$. By hypothesis, $p' \in B$. Moreover, $T(p')$ must cross Σ at some instant. Hence, there exists a $p'' \in T(p')$ such that $p'' \gg_{\mathscr{I}} p'$ and $p'' \in \Sigma$. Let η be the null generator of Σ which contains p''. As the segment of η which links p to p'' is null and achronal, η is necessarily the fastest curve $(\mathscr{I}, \bar{g}^{(0)})$ linking p to $T(p')$.

Now, consider the achronal boundary $\partial I^+(p, \overline{\mathscr{M}}) = \partial J^+(p, \overline{\mathscr{M}}) \doteq \overline{\Sigma}$. $\overline{\Sigma} \cap \mathscr{I}$ is closed, achronal and intersects each time generator of $\mathscr{I}$ at precisely one point, for every time generator possesses points in $I^+(p, \overline{\mathscr{M}})$ and $I^-(p, \overline{\mathscr{M}})$, and $\overline{\Sigma}$ splits $\overline{\mathscr{M}}$ in two disjoint open sets (from the viewpoint of a manifold with boundary, of course) $I^+(p, \overline{\mathscr{M}}) \doteq \dot{A}$ and $\overline{\mathscr{M}} \setminus \overline{I^+(p, \overline{\mathscr{M}})} \doteq \dot{B}$. Thus, $T(p')$ needs to cross $\overline{\Sigma}$ at, say, p'''. Once that $p < p'$ from the viewpoint of $(\overline{\mathscr{M}}, \bar{g})$, we must have $p''' \ll_{\mathscr{I}} p'$ or $p''' = p'$. In both cases, we have $p''' \ll_{\mathscr{I}} p''$, which implies that the null generator $\bar{\eta}$ of $\overline{\Sigma}$ containing p''' is strictly faster than η. As η was

the fastest curve in $(\mathscr{I}, \bar{g}^{(0)})$ linking p to $T(p')$, $\bar{\eta}$ necessarily traverses $\mathscr{M}$, which contradicts the hypotheses of the Theorem, for $\bar{\eta}$ must be necessarily achronal and, hence, cannot have a pair of conjugate points [BEE96, HE73, Wal84]. □

Lemma 1.5 *Let $p, p' \in \mathscr{I}$. If $p' \in \partial I^+(p, \mathscr{I})$ and $p' \neq \bar{p}$, then there cannot exist a null geodesic segment in $(\overline{\mathscr{M}}, \bar{g})$ which doesn't belong to $\mathscr{I}$ linking p to p'.*

Proof. Let η the (necessarily unique) null generator of $\partial I^+(p, \mathscr{I})$ linking p to p'. Suppose that there exists another null geodesic $\bar{\eta}$ traversing *mathscrM* and liking p to p'. As $(\mathscr{I}, \bar{g}^{(0)})$ is totally geodesic by construction, it follows that $\bar{\eta}$ necessarily incides at p' transversally to $\mathscr{I}$. Hence, if we choose any point p'' in η after p', then there exists a *broken* future directed null geodesic segment linking p to p'', which implies, on its turn, the existence of a timelike curve in $(\overline{\mathscr{M}}, \bar{g})$ linking p to p'' [HE73].

Let $T(p'')$ the time generator of $(\mathscr{I}, \bar{g}^{(0)})$ containing p''. Now, consider $\overline{\Sigma}$ as in the previous Lemma. Once again, $T(p'')$ must cross $\overline{\Sigma}$, say, at p'''. Thus, necessarily $p''' \ll_{\mathscr{I}} p''$. But this implies that, given that $(\overline{\mathscr{M}}, \bar{g})$ is (strongly) causal, $p \perp_{\mathscr{I}} p'$. This contradicts the result of Lemma 2.1. □

Going back to the proof of the Theorem, let γ a null geodesic segment emanating from p, traversing $\mathscr{M}$ and with future endpoint $p' \in \mathscr{I}$. As we've assumed that $(\mathscr{M}, g)$ is (strongly) causal, one sees that $p' \notin \overline{I^-(p, \overline{\mathscr{M}})}$. The Lemmata 1.4 and 1.5 imply that, if p' is not dragged into $I^+(p, \mathscr{I})$ by gravitational time delay, then $p' = \bar{p}$, exactly as in the case of AdS_d. However, even in this case, the presence of a pair of conjugate points in γ implies that there exists a timelike curve traversing the bulk and linking p to $p' = \bar{p}$. Repeating the argument of Lemma 1.5, the result follows. The proof for the case with reverse time orientation is the same. □

Remark 1.6 *The method used in the proof is similar to the one of the positive mass theorem for asymptotically flat spacetimes due to* Penrose, Sorkin *and* Woolgar *[PSW93], and for AAdS spacetimes by* Woolgar *[Woo94] and* Page, Surya *and* Woolgar *[PSW02]. A proof of a result similar to Theorem 1.3, employing a somewhat different strategy, was proposed by* Gao *and* Wald *[GW00].*

The condition stated in Theorem 1.3 about the global focusing of null geodesics exerts a crucial role in the latter's proof, for the omnipresence of pairs of conjugate points exclude presence of *null lines*, i.e., complete *achronal* null geodesics. This comes from the fact that, given a pair of conjugate points $p < q$ in a null geodesic γ, any point $r > q$, resp. $r < p$ satisfies $p \ll r$, resp. $q \gg r$. More precisely, the proof of Theorem 1.3 consists in showing that, in all possible instances of violation of the assertion, it's

possible to construct a null line traversing the bulk. Such a situation sharply contrasts with the AdS_d case, once that in the latter *all* null geodesics emanating from p and traversing the bulk meet at $\overline{p}$.

This focusing condition is valid if, for instance, the EINSTEIN equations without matter (1.1) are satisfied everywhere, for these imply the validity of the *null energy condition (NEC)* $\mathrm{Ric}(g)_{ab}k^a k^b \geq 0$ for all lightlike k^a, *and* if the so-called *null generic condition* is satisfied by all inextendible null geodesics (see [BEE96] for a discussion on this latter condition). Roughly, this last condition is necessary so that the null expansion of $\partial I^+(p, \overline{\mathscr{M}}) \cap \mathscr{M}$ eventually becomes negative at each one of its generators – the RAYCHAUDHURI equation (A.7) and NEC, then, take care of showing that there exist two conjugate points in the maximal extension of each generator (see in Appendix A the discussion following the formulae (A.11), page 110, and (A.23), page 113).

The violation of the null generic condition suggests a certain geometrical rigidity for null hypersurfaces having such a property. That this intuition is justified at global level, it can be seen in the following results:

THEOREM 1.6 (GEOMETRICAL MAXIMUM THEOREM [AGH98a, Gal00]) *Let $\mathscr{S}_1$ be a* future *null hypersurface (i.e., whose (future) null generators are future inextendible) and $\mathscr{S}_2$ a* past *null hypersurface (i.e., whose (past) null generators are past inextendible). Suppose that:*

1. *$\mathscr{S}_1$ and $\mathscr{S}_2$ possess a common point $p \in \mathscr{M}$ and $\mathscr{S}_2$ finds itself in the future side of $\mathscr{S}_1$ in a neighbourhood of p;*

2. *$\mathscr{S}_1$ has null expansion $\theta_1 > 0$ (see in Appendix A the discussion immediately preceding formula (A.9), page 109) in the sense of supports [6] such that the set of second null fundamental forms of the support hypersurfaces $\mathscr{S}_{,\epsilon}$ is bounded below;*

3. *$\mathscr{S}_2$ possesses expansion $\theta_2 \leq 0$ in the sense of supports,*

then $\mathscr{S}_1$ and $\mathscr{S}_2$ coincide in a neighbourhood $\mathscr{O}$ of p. Moreover, $\mathscr{S}_1 \cap \mathscr{O} = \mathscr{S}_2 \cap \mathscr{O}$ is a null $\mathscr{C}^\infty$ hypersurface with expansion $\theta = 0$. □

[6] Given a future null $\mathscr{C}^0$ hypersurface $\mathscr{S}$ and $p \in \mathscr{S}$, a *future (resp. past) support hypersurface* for $\mathscr{S}$ at p is a future (resp. past) null $\mathscr{C}^\infty$ hypersurface $\mathscr{S}_p$ possessing a null geodesic segment, containing p in its interior and localized in the future (resp. past) side of $\mathscr{S}$ in a neighbourhood of p. We analogously define support hypersurfaces for past null $\mathscr{C}^0$ hypersurfaces. We say that a future (resp. past) null $\mathscr{C}^0$ hypersurface $\mathscr{S}$ possesses *expansion* $\theta \geq 0$ (resp. ≤ 0) *in the sense of supports* if, for any $p \in \mathscr{S}$, $\epsilon > 0$, there exists a past (resp. future) support hypersurface $\mathscr{S}_{p,\epsilon}$ for $\mathscr{S}$ at p such that the expansion $\theta_{p,\epsilon}$ of $\mathscr{S}_{p,\epsilon}$ satisfies $\theta_{p,\epsilon}(p) \geq -\epsilon$ (resp. $\leq +\epsilon$). Support hypersurfaces constitute a way of studying the expansion of non differentiable null hypersurfaces due to the presence of caustics, as in for instance achronal boundaries, black hole horizons, etc..

The proof of Theorem 1.6 consists in denoting $\mathscr{S}_1$ and $\mathscr{S}_2$ locally by the graph of a Lipschitz function, and, in these coordinates, rewriting the null expansion as a quasilinear elliptic operator of second order, similar to the mean curvature operator when the former is prescribed in a codimension-one Riemannian submanifold, and for which there are maximum principles (see, for instance, [AGH98a, GT98]). From Theorem 1.6, it follows the

Theorem 1.7 (Null Splitting Theorem [Gal00]) *Let $(\mathscr{M}, g)$ be a spacetime endowed with null geodesic completeness and whose* Ricci *tensor satisfies NEC everywhere. If $\mathscr{M}$ has a null line γ, then γ belongs to an achronal, edgeless and* totally geodesic *null hypersurface, i.e., any geodesic tangent to this hypersurface at any point belongs to the latter in a neighbourhood of this point.* □

Returning to our study, another fundamental difference between AdS spacetimes and "generic" AAdS spacetimes is expressed by the following Proposition, which in particular implies that the collection of wedges in an AAdS spacetimes may not be closed under causal complements, albeit the Rehren bijection still preserves causality in this context. More precisely, we have $\mathscr{W}'_{p,\bar{q}} \cap \mathscr{W}'_{q,\bar{p}} \neq \varnothing$ (see Figure 1.4), due to the

Proposition 1.8 (Second Law of Dynamics of AAdS Wedges) *Let $(\mathscr{M}, g)$ be an AAdS spacetime satisfying the NEC $Ric(g)k^a k^b \geq 0$ for any lightlike vector k^a and the conditions of Theorem 1.3. Define $\Xi_p^+ \doteq \partial I^+(p, \overline{\mathscr{M}}) \setminus \{p, \bar{p}\}$ and $\Xi_{\bar{p}}^- \doteq \partial I^-(\bar{p}, \overline{\mathscr{M}}) \setminus \{p, \bar{p}\}$. Then:*

(i) $\Xi_p^+ \cap I^-(\bar{p}, \overline{\mathscr{M}}) = \Xi_{\bar{p}}^- \cap I^+(p, \overline{\mathscr{M}}) = \varnothing$.

(ii) $\Xi_p^+ \cap \Xi_{\bar{p}}^- \cap \mathscr{M} = \varnothing$.

Proof. (i) We know that $\Xi_p^+ \cap \mathscr{I} = \Xi_{\bar{p}}^- \cap \mathscr{I}$, then let's concentrate only at the bulk. Namely, suppose that there exists $q \in \mathscr{M}$ such that $q \in \Xi_{\bar{p}}^-$ and $q \notin \Xi_p^+$. If $q \gg p$, this contradicts the fact that there is no timelike curve linking p to $\bar{p}$. Repeat the argument exchanging past with future, and the roles of p and $\bar{p}$.

(ii) Suppose that Ξ_p^+ and $\Xi_{\bar{p}}^-$ coincide at some point $q \in \mathscr{M}$. We know that (a) *$\mathscr{S}_1 \doteq \Xi_{\bar{p}}^- \cap \mathscr{M}$ is a future null $\mathscr{C}^0$ hypersurface, and $\mathscr{S}_2 \doteq \Xi_p^+$ is a past null $\mathscr{C}^0$ hypersurface.* Taking any $q_2 \leq q \leq q_1$ with $q_i \in \mathscr{S}_i$, $i = 1, 2$, we see that $\partial I^-(q_1, \mathscr{M})$ contains a past support hypersurface $\mathscr{S}_-$ for $\mathscr{S}_1$ at q and $\partial I^+(q_2, \mathscr{M})$, a future support hypersurface $\mathscr{S}_+$ for $\mathscr{S}_2$ at q. Using the hypothesis that $\mathrm{Ric}(g)_{ab} k^a k^b \geq 0$ for any null vector k^a, we can invoke the Raychaudhuri equation (A.7) (page 108) and, recalling that the null geodesic congruences given by $\mathscr{S}_-$ and $\mathscr{S}_+$ have torsion $\omega_{+ab} = \omega_{-ab} = 0$, show that $-\frac{1}{\theta_-^2}\frac{d\theta_-}{d\lambda_-} \leq -\frac{1}{d-2}$ and $-\frac{1}{\theta_+^2}\frac{d\theta_+}{d\lambda_+} \geq +\frac{1}{d-2}$, where $\lambda_\pm$ is an affine parametrization of the null geodesics $\gamma_\pm$

respectively in $\mathscr{S}_2$ and $\mathscr{S}_1$ such that $\gamma_\pm(0) = q$, $\gamma_-(\lambda) = q_2$ and $\gamma_+(\lambda) = q_1$, for some $\lambda > 0$. Hence it follows that $\theta_-(p) \geq -\frac{d-2}{\lambda}$ and $\theta_+(p) \leq \frac{d-2}{\lambda}$. Taking q_2 sufficiently close to p and q_1 sufficiently close to $\bar{p}$,we can take λ arbitrarily large. Thus, (b) *$\mathscr{S}_1$ and $\mathscr{S}_2$ satisfy $\theta_2 \leq 0 \leq \theta_1$ in the sense of supports.* Hence, we conclude from (a), (i) and (b) that $\mathscr{S}_1$ and $\mathscr{S}_2$ satisfy the hypotheses of Theorem 1.6 and, as a consequence, both coincide in a neighbourhood of q, where they form a null $\mathscr{C}^\infty$ hypersurface with zero expansion. This implies that Ξ_p^+ and $\Xi_{\bar{p}}^-$ must coincide along a null geodesic containing q and hence traversing the bulk. In this case, γ must possess p as past endpoint and $\bar{p}$ as future endpoint. But this geodesic belongs to an achronal boundary, and thus it must be achronal, contradicting the hypotheses of Theorem 1.3. □

REMARK 1.7 *Notice that, before we invoke the hypothesis of global focusing of Theorem 1.3 in the proof of (ii), we've seen that, once more, one produced a null line γ contained in Ξ_p^+ and $\Xi_{\bar{p}}^-$. In the moment that we invoked Theorem 1.6, if $(\mathscr{M}, g)$ were* asymptotically simple*, we could have gone even further: by the Null Splitting Theorem 1.7, the null hypersurface which contains the null line in $\mathscr{M}$ linking p to $\bar{p}$ has no boundary, which would be constituted by the endpoints of $\partial I^+(p, \mathscr{M})$ and $\partial I^-(\bar{p}, \mathscr{M})$ if it existed. Thus, it follows from null geodesic completeness of $(\mathscr{M}, g)$ that $\partial I^+(p, \mathscr{M}) = \partial I^-(\bar{p}, \mathscr{M})$ is $\mathscr{C}^\infty$ and totally geodesic with respect to g.*

An important consequence of Proposition 1.8 is that the focusing condition causes a nontrivial "shrinking" of AAdS wedges towards conformal infinity, as we can see in Figure 1.4 – see the discussion at the end of Chapter 2 for a dynamical interpretation of this phenomenon, which explains the why of our denominating this Proposition *Second Law of Dynamics of AAdS Wedges.*

1.3.1.2 Simple connectedness (bulk *versus* boundary)

We shall cite now, for completeness, two results which show that every wedge $\mathscr{W}_{p,q} \in \mathscr{W}(\mathscr{M}, g)$ in a locally AdS-type spacetime $(\mathscr{M}, g)$, satisfying certain additional conditions, is *simply connected* if $\mathscr{D}_{p,q}$ is (for instance, wedges contained in a POINCARÉ domain of an AAdS spacetime). More precisely, any causal curve in the bulk with endpoints p and q is homotopic to a causal curve *in* $\mathscr{I}$ linking p to q. Summing up, the topology of $\mathscr{W}_{p,q}$ is determined by the topology of $\mathscr{D}_{p,q}$, showing that *any wedge $\mathscr{W}_{p,q} \in \mathscr{W}(\mathscr{M}, g)$ is contractible.*

This is a particular case of *topological censorship* – the topology of the region exterior to the event horizon of a black hole is determined by the topology of $\mathscr{I}$ – proven in the present context by GALLOWAY, SCHLEICH, WITT and WOOLGAR [GSWW99].



Figure 1.4: Effect of gravitational time delay (whose effect on null geodesics is exemplified by the dotted lines in the longitudinal cut **(b)** of **(a)**) on wedges and their causal complements in AAdS spacetimes. Whereas in AdS_d the causal complement of the wedge $\mathscr{W}_{p,\bar{q}}$ is the wedge $\mathscr{W}_{q,\bar{p}}$ (red), the causal complements of both wedges (respectively green and blue) are no longer wedges in AAdS spacetimes satisfying the conditions of Proposition 1.8 and, hence, of Theorem 1.3, and in such a way that the open region $\mathscr{W}'_{p,\bar{q}} \cap \mathscr{W}'_{q,\bar{p}} \neq \varnothing$ is *nonvoid*, which cannot happen in AdS_d.

Theorem 1.9 (Galloway, Schleich, Witt and Woolgar [GSWW99]) *Let $(\mathscr{M}, g)$ be a locally AdS-type spacetime such that $(\overline{\mathscr{M}}, \bar{g})$ is globally hyperbolic and $\mathscr{I}$ possesses a connected component $\mathscr{I}_0$ with a compact spacelike cut.*[7] *Suppose that any future complete null geodesic γ in $(\overline{\mathscr{M}}, \bar{g})$ emanating from some $p = \gamma(0)$ in a neigh-*

[7] Given a (connected) manifold $\mathscr{M}$, a *cut* is a hypersurface without boundary Σ such that $\mathscr{M} \setminus \Sigma$ has two connected components.

bourhood of $\mathscr{I}_0$ satisfies $\int_0^{+\infty} \mathrm{Ric}(\bar{g}(\gamma(\lambda)))_{ab}\dot{\gamma}^a(\lambda)\dot{\gamma}^b(\lambda)d\lambda \geq 0$. Then, there is no future causal communication of $\mathscr{I}_0$ through $\mathscr{M}$ with any other component of $\mathscr{I}$, i.e., $J^+(\mathscr{I}_0, \overline{\mathscr{M}}) \cap (\mathscr{I} \setminus \mathscr{I}_0) = \varnothing$. □

Theorem 1.9 splendidly complements Lemma 1.4, employed in the proof of Theorem 1.3. Using Theorem 1.9 and covering space arguments [GH81, O'N83], cone can finally prove the following *topological censorship* theorem:

Theorem 1.10 (Galloway, Schleich, Witt and Woolgar [GSWW99]) *Let $(\mathscr{M}, g)$ be a locally AdS-type spacetime such that $(\mathscr{W}(\mathscr{I}) \cup \mathscr{I}, \bar{g}\restriction_{\mathscr{W}(\mathscr{I})\cup\mathscr{I}})$ is globally hyperbolic and $\mathscr{I}$ possesses a compact spacelike cut. Suppose that any future complete null geodesic γ in $(\overline{\mathscr{M}}, \bar{g})$ emanating from some $p = \gamma(0)$ in a neighbourhood of $\mathscr{I}$ satisfies $\int_0^{+\infty} \mathrm{Ric}(\bar{g}(\gamma(\lambda)))_{ab}\dot{\gamma}^a(\lambda)\dot{\gamma}^b(\lambda)d\lambda \geq 0$. Then any causal curve $\gamma : (0,1) \to \mathscr{W}(\mathscr{I})$ with endpoints $\gamma(0), \gamma(1) \in \mathscr{I}$ is fixed endpoint homotopic to a curve $\gamma_0 : [0,1] \to \mathscr{I}$ (see AppendixD).* □

It follows immediately from Theorem 1.10 the contractivity of $\mathscr{W}_{p,q}$ for $p, q \in \mathscr{D}(\mathscr{I})$. In particular, if $p, q \in \mathscr{D}(\mathscr{I})$ are the endpoints of γ in Theorem 1.10, we can choose γ_0 *chronological*.

1.3.1.3 Localization (boundary to bulk)

Throughout this Subsubsection, it'll be assumed that $(\mathscr{M}, g)$ is an asymptotically simple AAdS spacetime satisfying the hypotheses of Theorem 1.3.

Knowing the localization of the physical procedures in wedges may not be enough for the complete reconstruction of the bulk quantum theory in the context of Chapter 4, only by employing boundary information and the Rehren bijection $\rho_{(\mathscr{M},g)}$ introduced by Definition 1.5. We need to be able to specify the localization of physical procedures in arbitrarily small open sets, or, which amounts to the same thing, its localization with respect to a base for the bulk topology. This can be achieved, in principle, by taking intersections of wedges, but it's by no means clear whether this results in a base or not. One thus needs to make this rough idea more precise.

It's known that, for strongly causal spacetimes, the topology generated by diamonds (Alexandrov topology) coincides with the manifold topology [BEE96]. Hence, in AdS, the above question has a positive answer, for *any diamond in AdS_d can be enveloped by wedges:* given

$$\mathscr{O}_{p,q} \doteq I^+(p, AdS_d) \cap I^-(q, AdS_d),\ p \ll_{AdS_d} q, \tag{1.33}$$

we can write

$$\mathscr{O}_{p,q} = \bigcap_{\substack{r\in\partial I^-(p,\overline{AdS_d})\cap\mathscr{I},\\ s\in\partial I^+(q,\overline{AdS_d}),\, r,s\in\mathscr{M}in(u)}} \mathscr{W}_{r,s}. \tag{1.34}$$

We shall see soon that the achronality of inextendible null geodesics in AdS_d is tantamount for a precise envelope, whereas in an asymptotically simple AAdS spacetime satisfying the hypotheses of Theorem 1.3, the question is more delicate, because of the following

Proposition 1.11 *Let $p \in \mathscr{M}$. Then, $\partial I^+(p,\overline{\mathscr{M}})$ intersects each time generator of $(\mathscr{I},\bar{g}^{(0)})$* precisely once.

Proof. Due to the achronality of $\partial I^+(p,\overline{\mathscr{M}})$, the latter intersects each time generator of $(\mathscr{I},\bar{g}^{(0)})$ *at most once*. Suppose that the assertion is false. Then, given a time generator T, we have the following possibilities:

(i) $T \subset I^+(p,\overline{\mathscr{M}})$ – Consider a complete null geodesic γ crossing p, and let q be the *past* endpoint of γ. Then there exists a value t of the affine parameter of T such that $T(t) \ll_{\mathscr{I}} q$. Hence, $T(t) \ll_{\overline{\mathscr{M}}} p$, which is absurd since $(\overline{\mathscr{M}},\bar{g})$ is chronological.

(ii) $T \cap J^+(p,\overline{\mathscr{M}}) = \varnothing$ – Consider a complete null geodesic γ crossing p, and let r the *future* endpoint of γ. Then there exists a value t of the affine parameter of T such that $T(t) \gg_{\mathscr{I}} r$. Hence, $T(t) \gg_{\overline{\mathscr{M}}} p$, contradicting the hypothesis.

□

Proposition 1.12 *Let $q, r \in \mathscr{M}$ such that $r \in \partial I^-(q,\mathscr{M})$, and γ a null generator of $\partial I^-(q,\mathscr{M})$ to which r belongs. Let $s_1(r), s_2(r), s_3(r) \in \mathscr{I}$ defined as follows:*

- *$s_1(r)$ is the future endpoint of γ;*
- *$s_2(r)$ is the point where $\partial I^+(q,\overline{\mathscr{M}})$ intersects the time generator of $T(s_1(r))$ to which $s_1(r)$ belongs;*
- *$s_3(r)$ is the point where $\partial I^+(r,\overline{\mathscr{M}})$ intersects $T(s_1(r))$.*

Then:

(i) $s_3(r) \leq_{\mathscr{I}} s_2(r) \leq_{\mathscr{I}} s_1(r)$.

(ii) $s_3(r) = s_2(r) = s_1(r)$ *if and only if the segment of γ linking r to $s_1(r)$ is achronal.*

PROOF. (i) Immediate, as (ii) $\Rightarrow$. It remains only to prove (ii) $\Leftarrow$. Namely, suppose that $s_3(r)$ coincides with $s_2(r)$. Then the null geodesic segment linking q to $s_2(r)$ must belong to γ, for otherwise there would exist a broken null geodesic segment linking r to $s_3(r)$, contradicting the definition of the latter (this, in particular, proves that $s_1(r) = s_3(r)$ even if we only assume $s_2(r) = s_3(r)$). If γ is not achronal, once again we have a contradiction with the definition of $s_3(r)$. $\square$

REMARK 1.8 *Results analogous to Propositions 1.11 and 1.12 are valid if we exchange future with past.*

Now, let $\mathscr{O}_{p,q} \subset \mathscr{M}$ be onwards a *relatively compact diamond possessing a contractible* CAUCHY *surface* – any sufficiently small diamond satisfies both conditions. Let us then consider the region

$$\mathscr{Q}_{p,q} = \bigcap_{\substack{r\in\partial I^-(p,\overline{\mathscr{M}})\cap\mathscr{I},\\ s\in\partial I^+(q,\overline{\mathscr{M}})\cap T(r)}} \mathscr{W}_{r,s}. \tag{1.35}$$

IT follows naturally from this definition that $\mathscr{Q}_{p,q} \supset \mathscr{O}_{p,q}$, is *causally complete*, as it's an intersection of causally complete regions, and

$$\mathscr{Q}_{p,q} \cap J^+(q,\overline{\mathscr{M}}) = \mathscr{Q}_{p,q} \cap J^-(p,\overline{\mathscr{M}}) = \varnothing. \tag{1.36}$$

In AdS_d, $\mathscr{Q}_{p,q} = \mathscr{O}_{p,q}$. For asymptotically simple AAdS spacetimes satisfying the as hypotheses of Theorem 1.3, however, it may happen that $\mathscr{Q}_{p,q} \supsetneqq \mathscr{O}_{p,q}$. Analogously, defining $\mathscr{E}_{p,q} \doteq \partial I^+(p,\mathscr{M}) \cap \partial I^-(q,\mathscr{M})$, let's start from

$$\widetilde{\mathscr{Q}}_{p,q} = \bigcap_{r\in\mathscr{E}_{p,q}} \mathscr{W}_{s_3'(r),s_3(r)}, \tag{1.37}$$

where $s_3'(r)$ corresponds to $s_3(r)$ if we exchange future with past in the statement of Proposition 1.12. Here, $\widetilde{\mathscr{Q}}_{p,q} \subset \mathscr{Q}_{p,q}$ is once more causally complete, *if nonvoid*. However, is the metric deep inside the bulk is sufficiently "distorted", making a sufficient number of null generators of, say, $\partial I^-(q,\mathscr{M})$ to acquire pairs of conjugate points between $\mathscr{E}_{p,q}$ and $\mathscr{I}$, for all we know (Proposition 1.12, page 24) $\widetilde{\mathscr{Q}}_{p,q}$ can very well be empty. This is suggested by the following remarks:

1. In a causally simple spacetime, any relatively compact diamond $\mathscr{O}_{p,q}$ is a globally hyperbolic region, for which any CAUCHY surface possess boundary $\mathscr{E}_{p,q}$;

2. Any causally complete region $\mathscr{U}$ possesses the following property: if $\mathscr{S} \subset \mathscr{U}$ closed, achronal set with respect to $\mathscr{U}$, then $D(\mathscr{S}) \subset \mathscr{U}$.

Both remarks together show that, if $r \in \mathscr{E}_{p,q}$ is such that a null generator of, say, $\partial I^-(q, \mathscr{M})$ crossing r acquires a pair of conjugate points between r and $s_1(r)$, then, by causal simplicity, there exists a neighbourhood of q causally disjoint from $s_3(r)$, and, thus, $I^-(s_3(r), \overline{\mathscr{M}})$ cannot contain a Cauchy surface for $\mathscr{O}_{p,q}$. Since, on the other hand, this doesn't exclude the possibility that $\widetilde{\mathscr{Q}}_{p,q}$ may contain points outside $\mathscr{O}_{p,q}$ either, it's by no means clear whether the collections of $\mathscr{Q}_{p,q}$'s and $\widetilde{\mathscr{Q}}_{p,q}$'s produce bases for the topology of $\mathscr{M}$ or not.

A way of circumventing these problems could be to restrict our considerations to sufficiently small diamonds, such that no null generator of $\partial I^+(q, \overline{\mathscr{M}})$ can pursue long enough a path beyond q which allows it to acquire a pair of conjugate points. There is, however, a situation in which no matter how small the extension, it will always cease to be achronal: it's when q *itself* is conjugate to $s_1(r)$. In this limit case, $s_1(r) = s_2(r)$ but $s_2(r) \neq s_3(r)$.

We'll show now that the key for the end of these problems is trying to build a region similar to $\widetilde{\mathscr{Q}}_{p,q}$, but employing, instead of the points $s_3(r)$, $s_3'(r)$ for $r \in \mathscr{E}_{p,q}$, the points for which the problem above, implied by Proposition 1.12, is, in a certain sense, "minimized". In order to undertake such a task and hence optimize our construction, we shall start from a different viewpoint, which will end up showing that the critical situation mentioned in the former paragraph is excluded by null geodesic completeness.

First, notice that, by means of an argument analogous to the one employed in [HE73] and [Wal84] to prove the existence of a topological (Lipschitz) manifold structure for achronal boundaries, one can show that $\mathscr{E}_{p,q}$ is locally the graph of a locally Lipschitz function with values in $\mathbb{R}$ with $d-2$ arguments, and thus an achronal, compact topological (Lipschitz) submanifold embedded into $\mathscr{M}$, with codimension two. Notice as well that one can parametrize in a $\mathscr{C}^\infty$ way the family of time generators of $(\mathscr{I}, \bar{g}^{(0)})$ by means of a Cauchy surface $\mathscr{S}$ of the latter, homeomorphic to S^{d-2} and hence also compact. Let t be the affine parameter common to the time generators of $(\mathscr{I}, b)$ mentioned above, and $\mathscr{F}_{p,q}$ some Cauchy surface for $\mathscr{O}_{p,q}$, whose boundary (edge) is of course $\mathscr{E}_{p,q}$. Define implicitly the functions

$$\tau^\pm : \overline{\mathscr{F}_{p,q}} \times \mathscr{S} \ni (r, \theta) \mapsto \tau^\pm(r, \theta) \in \mathbb{R}, \tag{1.38}$$

where

$$\partial I^\pm(R, \overline{\mathscr{M}}) \cap T(\theta) = \{T(\theta)(\tau^\pm(r, \theta))\}. \tag{1.39}$$

Proposition 1.11 shows that the definition of $\tau^\pm$ is not empty. Moreover:

Proposition 1.13 *τ^+ (resp. τ^-) is upper (resp. lower) semicontinuous in r at fixed θ.*

PROOF. We'll just prove it for τ^+ – the result for τ^- follows analogously. Let $\epsilon > 0$. r belongs to the chronological past of the point $T(\theta)(\tau^+(r,\theta)+\epsilon)$, and thus there exists an open neighbourhood U of r in $\overline{\mathscr{F}_{p,q}}$ (with respect to the relative topology) which stays in the chronological past of $T(\theta)(\tau^+(r,\theta)+\epsilon)$. Hence, for all $r' \in U$, we must have $\tau^+(r',\theta) < \tau^+(r,\theta)+\epsilon$. □

One may yet prove that $\tau^\pm$ are locally LIPSCHITZ in θ for fixed r, but this won't be used in what follows. The function $\tau^+(.,\theta)$ (resp. $\tau^-(.,\theta)$) will be called *future* (resp. *past*) FERMAT *potential* with respect to θ.[8] The name is reminiscent from the HUYGENS-FERMAT principle in geometrical optics (see, for instance, pages 249-250 of [Arn89] for a beautiful proof). Since $\mathscr{E}_{p,q} = \partial\mathscr{F}_{p,q}$ as well as $\overline{\mathscr{F}_{p,q}}$ are closed subsets of the compact set $\overline{\mathscr{O}_{p,q}}$, they are themselves compact. It then follows from a standard result in Analysis (see, for instance, pages 110-111 of [KF75]) that $\tau^+(.,\theta)$ (resp. $\tau^-(.,\theta)$) possesses a maximum (resp. minimum) in $\overline{\mathscr{F}_{p,q}}$ as well as in $\mathscr{E}_{p,q}$. The next Theorem shows that $\pm\tau^\pm(.,\theta)$ has, indeed, a typical property of potentials:

THEOREM 1.14 (MAXIMUM / MINIMUM PRINCIPLE FOR FERMAT POTENTIALS) *The maximum (resp. minimum) value of $\tau^+(.,\theta)$ (resp. $\tau^-(.,\theta)$) in $\overline{\mathscr{F}_{p,q}}$ is attained in $\mathscr{E}_{p,q}$.*

PROOF. As done in the proof of Proposition 1.13, we shall just prove the result for τ^+. Let r be a point of $\mathscr{E}_{p,q}$ where $\tau^+(.,\theta)$ attains its maximum in $\mathscr{E}_{p,q}$, and let r' be a point of $\mathscr{F}_{p,q}$ such that $\tau^+(r',\theta) \geq \tau^+(r',\theta)$. In this case, it's obvious that $\mathscr{E}_{p,q}$ belongs to the causal past of $T(\theta)(\tau^+(r',\theta))$. Pick a curve segment in $\overline{\mathscr{F}_{p,q}}$ starting in r', initially pointing outside $J^-(T(\theta)(\tau^+(r',\theta)),\overline{\mathscr{M}})$ and terminating at some point of $\mathscr{E}_{p,q}$. Then any such curve segment must cross $\partial I^-(T(\theta)(\tau^+(r',\theta)),\overline{\mathscr{M}})$ at least once more after r', and before (or when) hitting $\mathscr{E}_{p,q}$. This shows that $\partial I^-(T(\theta)(\tau^+(r',\theta)),\overline{\mathscr{M}}) \cap \overline{\mathscr{F}_{p,q}}$ contains an open subset X of $\mathscr{F}_{p,q}$ outside the causal past of $T(\theta)(\tau^+(r',\theta))$.

The rest of the proof is analogous to the proof of PENROSE's singularity theorem [HE73, Wal84]: namely, we'll show that the properties of ∂X imply the existence of an incomplete null geodesic in $(\mathscr{M},g)$. First, we shall show that the closed, acausal set $\partial X = \partial I^-(T(\theta)(\tau^+(r',\theta)),\overline{\mathscr{M}}) \cap \overline{\mathscr{F}_{p,q}}$ is *past trapped*, i.e., $\partial I^-(X,\mathscr{M})$ is *compact*. The past null geodesics "entering" ∂X constitute the past CAUCHY horizon of X, which is hence contained in $\overline{\mathscr{O}_{p,q}}$ and, thus, compact, for it's closed. The outgoing null geodesics are precisely the null generators of $\partial I^-(T(\theta)(\tau^+(r',\theta)),\overline{\mathscr{M}})$ which cross ∂X. Let us adopt a common affine parametrization for the null generators of $\partial I^-(T(\theta)(\tau^+(r',\theta)),\overline{\mathscr{M}})$ such that the zero of the affine parameter corresponds to ∂X. Then let t_0 the largest value of the affine parameter for which a past endpoint of $\partial I^-(T(\theta)(\tau^+(r',\theta)),\overline{\mathscr{M}})$ is attained. This value must be finite, for any inextendible null geodesic must

[8] Actually, $\tau^\pm$ also depend on the choice of foliation of $\mathscr{I}$ by CAUCHY surfaces, but such a dependence is irrelevant for the use we'll make of $\tau^\pm$ and will be ignored.

acquire a pair of conjugate point before hitting infinity, though the value of the affine parameter in a past endpoint of the null generator segment starting, say, in $r'' \in \partial X$ can be zero if r' happens to be a past endpoint himself. Anyway, the portion of $\partial I^-(T(\theta)(\tau^+(r',\theta)),\overline{\mathscr{M}})$ in the causal past of ∂X, being closed, has a closed inverse image in the compact $[0,t_0]\times\partial X$ by the parametrization chosen for the null generators, and is thus compact. Hence, the set $\partial I^-(\partial X,\mathscr{M}) = H^-(X)\cup\partial X\cup(\partial I^-(T(\theta)(\tau^+(r',\theta)),\overline{\mathscr{M}})\cap J^-(\partial X,\mathscr{M}))$ is a compact, achronal subset of $\mathscr{M}$, as stated.

However, any causally simple spacetime is stably causal [BEE96]. That is, one can foliate $\mathscr{M}$ in a $\mathscr{C}^\infty$ manner by "constant-time" spacelike, codimension-one hypersurfaces. Due to the structure of conformal infinity, these surfaces cannot be compact. Moreover, each timelike *orbit* of this foliation crosses an achronal set at most once. Following these orbits, one can continuously map $\partial I^-(\partial X,\mathscr{M})$ into a spacelike leaf. As the image of this map is compact, it must have a nonvoid boundary. It's, on the other hand, known that a set of the form $\partial I^-(Y,\mathscr{M}), Y\subset\mathscr{M}$ is a topological submanifold *without boundary* of $\mathscr{M}$. This shows that some null generator of $\partial I^-(T(\theta)(\tau^+(r',\theta)),\overline{\mathscr{M}})$ must hit a singularity before reaching its past endpoint. But this enters into conflict with the null geodesic completeness of $\widehat{\mathscr{M}}$, implied by asymptotic simplicity. Hence, no point of $\mathscr{F}_{p,q}$ can attain a maximum for $\tau^+(.,\theta)$ in $\overline{\mathscr{F}_{p,q}}$ – the former is always reached in $\mathscr{E}_{p,q}$. □

Proposition 1.13 and Theorem 1.14 together show that, for each θ, there is always a $r\in\mathscr{E}_{p,q}$ such that, given *any* Cauchy surface $\mathscr{F}_{p,q}$ for $\mathscr{O}_{p,q}$, the set $\overline{\mathscr{F}_{p,q}}$ always stays in the causal past of $T(\theta)(\tau^+(r,\theta))$. By Proposition 1.12 and the above remarks, this can only happen is the achronal null geodesic segment $\gamma(r,\theta)$ linking r to $T(\theta)(\tau^+(r,\theta))$ crosses q. Hence, this maximum point is *unique:* suppose otherwise. Then, there would exist another $r'\in\mathscr{E}_{p,q}$ such that there is an achronal null geodesic segment $\gamma(r',\theta)$ linking r' to $T(\theta)(\tau^+(r',\theta)) = T(\theta)(\tau^+(r,\theta))$ and crossing q. Now, consider the curve segment $\gamma'(r,\theta)$, which coincides with $\gamma(r,\theta)$ from r to q, and coincides with $\gamma(r',\theta)$ from q to $T(\theta)(\tau^+(r,\theta))$. This segment is necessarily broken, which conflicts with the achronality of $\gamma(r,\theta)$. Exchanging the roles of r and r', one can see that this argument also enters in conflict with the achronality of $\gamma(r',\theta)$.

Let us notice, however, that an arbitrary $r\in\mathscr{E}_{p,q}$ *need not* maximize $\tau^+(.,\theta)$ for some θ. The two situations where this indeed cannot occur are:

1. r is conjugate to q along a null generator of $\partial I^-(q,\mathscr{M})$ – any future extension of this generator beyond q won't be achronal;

2. q is conjugate to $s_2(r)$ along a null generator of $\partial I^+(q,\overline{\mathscr{M}})$, by the remarks made above.

The second instance, on the other hand, is excluded by our line of reasoning, for it makes impossible, by Proposition 1.12 and by Theorem 1.14, that $\tau^+(.,\theta)$ attains a maximum in $\mathscr{E}_{p,q}$. This cannot occur, since for each θ a maximum must exist by Proposition 1.13. The first instance can be circumvented if we pick $\mathscr{O}_{p,q}$ contained, say, in a convex normal neighbourhood, which can always be done, as here $(\mathscr{M}, g)$ is strongly causal. One can go beyond that and take $\mathscr{O}_{p,q}$ sufficiently small (albeit nonvoid) in a way that $r \in \mathscr{E}_{p,q}$ is a maximum of $\tau^+(.,\theta)$ for some θ, for the only obstacle to this would be the second instance above, excluded by the argument here presented. All these results possess a past counterpart, if we switch q with p and τ^+ with τ^-.

Summing up, we've shown that *any sufficiently small $\mathscr{O}_{p,q}$ can always be precisely enveloped by wedges.* In this case, any point that doesn't belong to $\overline{\mathscr{O}_{p,q}}$ belongs either to the chronological future of $\partial I^-(q,\mathscr{M})$ or to the chronological past of $\partial I^+(p,\mathscr{M})$, and, as such, will fail at belonging to any wedges enveloping $\mathscr{O}_{p,q}$. As the points in $\partial\mathscr{O}_{p,q}$ are automatically excluded from the intersection by construction, one concludes that $\mathscr{O}_{p,q} = \mathscr{Q}_{p,q}$ for $\mathscr{O}_{p,q}$ sufficiently small. Moreover, in this situation, each wedge in the definition (1.35) of $\mathscr{Q}_{p,q}$ is guaranteed to be contained in some POINCARÉ domain. Notice yet that the procedure we've proposed allows as well to envelope the *regular diamonds* (if sufficiently small) defined in [GLRV01].

Chapter 2

Dynamics and time evolution

Because I know that time is always time
And place is always and only place
And what is actual is actual only for one time
And only for one place

T. S. Eliot
"I. Because I do not hope to turn again"
(*Ash-Wednesday*, 1930)

One has made extensive use of the global structure of AAdS spacetimes in Chapter 1, but the Einstein equations have exerted a minor role in these developments. However, it has become clear along the previous Chapter that nontrivial geometries in the bulk of an AAdS spacetime provoke a distortion of the causal structure in the large, which, thus, must be noticeable from the viewpoint of a theory of local observables at the boundary. Hence, it's highly desirable to establish a more quantitative connection between these effects and the very gravitational dynamics, since Maldacena's conjecture and many of its consequences involve gravitational effects in an essential way.

2.1 Variational approach to classical gravity. Conceptual foundations

The Einstein equations (1.1) are obtained in the Lagrangian approach by means of the Einstein-Hilbert variational principle: for any compact set $K \Subset \mathscr{M}$ with piecewise $\mathscr{C}^\infty$ boundary (or, more generally, we assume that the set of non differentiable

points of ∂K have zero $(d-1)$-dimensional **Lebesgue** measure), the action functional

$$S_K[g] \doteq \frac{1}{16\pi G_d}\int_K \underbrace{(R(g)+2\Lambda)}_{\doteq \mathscr{L}_\Lambda(g);}\sqrt{|g|}dx \quad (2.1)$$

is stationary for arbitrary variations of g around a solution of the variational problem, supported in the interior of K. Namely, a finite variation of g is obtained from a $\mathscr{C}^\infty$ curve of metrics g_λ, $\lambda \in (-1,1)$, which differ from g in a compact set $K_1 \subset int(K)$ and such that $g_0 = g$. The corresponding infinitesimal variation (= tangent vector) around g is given by $\delta g = \frac{d}{d\lambda}|_{\lambda=0} g_\lambda$. The corresponding variation in (2.1) is given by

$$(\delta S_K)[g,\delta g] = \int_K \left[\nabla^a \underbrace{\left(\nabla^b \delta g_{ab} - g^{cd}\nabla_a \delta g_{cd}\right)}_{\doteq \theta_a(g,\delta g);} + \left(\mathrm{Ric}_{ab} - \frac{1}{2}Rg_{ab} + \Lambda g_{ab}\right)\delta g^{ab}\right]\sqrt{|g|}dx, \quad (2.2)$$

where the first term of the right hand side is the divergence of the Hodge-dual 1-form to the $(d-1)$-form $\theta^{a_1}\sqrt{|\det g|}_{[a_1\cdots a_d]}$, which vanishes in ∂K for the class of variations considered. The second term is the Einstein tensor $G(g)_{ab} \doteq \mathrm{Ric}(g)_{ab} - \frac{1}{2}R(g)g_{ab}$, plus the cosmological constant part Λ. Hence, we have the Einstein equations (1.1). Let us notice that the use of a compact domain of integration K reflects the fact that the variational principle that leads to 1.1 is *inherently local*, as any field theory.

We end this Section here emphasizing that, for any vector T^a, the expression $\left(\mathrm{Ric}(g)_{ab} - \frac{1}{2}R(g)g_{ab} + \Lambda g_{ab}\right)T^b$ doesn't have second order derivatives in direction of T^a. If T^a is timelike, thus determining (locally) a time flow, we see that the components $\left(\mathrm{Ric}(g)_{ab} - \frac{1}{2}R(g)g_{ab} + \Lambda g_{ab}\right)T^b = 0$ of the Einstein equations are actually *constraints*, expressing the symmetry of the latter under general coordinate transformations (*passive* viewpoint) or, equivalently in mathematical terms, under local diffeomorphisms (*active* viewpoint). The physical meaning of this fact, according to Einstein, is that gravity doesn't have an *intrinsic* notion of (dynamical) time evolution, for the very choice of time coordinate is a symmetry – such a choice must, hence, be made by means of a concrete physical procedure, in such a manner that the physical laws which rule such procedures in our gravitational background are independent of this choice. There are two ways to do this: one local – by the specification of a nontrivial energy-momentum tensor – and the other global, if the metric becomes close to some fixed background geometry at sufficiently long distances, for which such choice can be made naturally. The latter circumstance includes precisely the case of AAdS spacetimes.

[1] G_d is the d-dimensional Newton constant; henceforth, we choose units $G_d = (16\pi)^{-1}$, so as the numerical factor multiplying the right hand side of (2.1) becomes equal to one.

In the next Section, we'll make use of this fact to explore in detail the geometry of (locally) AAdS spacetimes near conformal infinity.

2.2 The Fefferman-Graham expansion

Let us consider the case of AdS-type spacetimes $(\mathscr{M}, g)$ as in Definition 1.4, whence we adopt the notation. Let's now scrutinize the form of the solutions of (1.1) for such spacetimes in a collar neighbourhood $\overline{\mathscr{U}}_\epsilon \cong \mathscr{I} \times [0, \epsilon) \ni (x, z)$, $\epsilon > 0$, of $\mathscr{I}$ in $\overline{\mathscr{M}}$. This will be made by means of the asymptotic expansion obtained by Fefferman and Graham [FG85], whose deduction we present below. In the subsequent argument, we closely follow [HIM05] and, in a smaller measure, [Gra00, GL91, HSS01].

An equivalent way of writing (1.1) is directly in terms of the Ricci tensor, resulting in

$$\mathrm{Ric}(g)_{ab} = \frac{2\Lambda}{d-2} g_{ab}. \tag{2.3}$$

Invoking now the formula (A.25) of the conformal transformation of $\mathrm{Ric}(g)$ into $\mathrm{Ric}(\bar{g})$, $\bar{g} = z^2 g$

$$\mathrm{Ric}(g)_{ab} = \mathrm{Ric}(\bar{g})_{ab} + \frac{d-2}{z}\bar{\nabla}_a\bar{\nabla}_b z + \bar{g}_{ab}\bar{g}^{cd}\left(\frac{1}{z}\bar{\nabla}_c\bar{\nabla}_d z - \frac{d-1}{z^2}\bar{\nabla}_c z\bar{\nabla}_d z\right),$$

we have the expression of (2.3) in terms of $\bar{g}$

$$\mathrm{Ric}(\bar{g})_{ab} + \frac{d-2}{z}\bar{\nabla}_a\bar{\nabla}_b z + \bar{g}_{ab}\bar{g}^{cd}\left(\frac{1}{z}\bar{\nabla}_c\bar{\nabla}_d z - \frac{d-1}{z^2}\bar{\nabla}_c z\bar{\nabla}_d z\right) = \frac{2\Lambda}{(d-2)z^2}\bar{g}_{ab}. \tag{2.4}$$

Notice, in particular, that if we multiply (2.4) by z^2 and take $z = 0$, it follows that dz necessarily satisfies

$$\bar{g}^{-1}(dz, dz)\restriction_{\mathscr{I}} = -\frac{2\Lambda}{(d-1)(d-2)} = \frac{1}{R^2}, \tag{2.5}$$

where R is the AdS radius of $(\mathscr{M}, g)$, defined in Chapter 1.

Let us now restrict (2.4) to the timelike hypersurfaces $\mathscr{I}'_\epsilon \doteq z^{-1}(\epsilon')$, $0 \leq \epsilon' < \epsilon$. First, we'll use the freedom we have in the choice of z to define a system of Gaussian normal coordinates in $\mathscr{U}_\epsilon$:

Lemma 2.1 *Given any AdS-type metric g satisfying the* Einstein *equations (2.3) in $\mathscr{U}_\epsilon$ for some $\epsilon > 0$, we can choose z and ϵ in a way that $^{(*)}$ $\bar{g}^{-1}(dz, dz) \doteq \zeta^2$ is constant in $\mathscr{U}_\epsilon$. In particular, $\zeta^2 = R^{-2}$.*

Proof. Let $z' = e^{\sigma} z$ and $\bar{g}' = e^{2\sigma}\bar{g}$; then, $dz' = e^{\sigma}(dz + z d\sigma)$, and hence

$$\bar{g}'^{-1}(dz', dz') = \bar{g}^{-1}(dz + z d\sigma, dz + z d\sigma) =$$
$$= \bar{g}^{-1}(dz, dz) + 2z\bar{g}^{-1}(dz, d\sigma) + z^2\bar{g}^{-1}(d\sigma, d\sigma).$$

Thus, the condition $^{(*)}$ for dz' is equivalent to the first-order nonlinear partial differential equation

$$2z\bar{g}^{-1}(dz, d\sigma) + z^2\bar{g}^{-1}(d\sigma, d\sigma) = \frac{\zeta^2 - \bar{g}^{-1}(dz, dz)}{z},$$

where the right hand side extends in a $\mathscr{C}^\infty$ way to $z = 0$ due to (2.5), which, on its turn, fixes $\zeta^2 = R^{-2}$. By the method of characteristics [Joh82] (see also Subsection 6.4 of [Hör90]), there exists $\epsilon > 0$ and a unique $\mathscr{C}^\infty$ solution of this equation in $\mathscr{U}_{2\epsilon}$ for arbitrary values of $\sigma\restriction_{\mathscr{I}}$. The result follows globally by employing a partition of unity subordinated to the covering $\{\mathscr{U}_{2\epsilon}, \mathscr{M} \setminus \overline{\mathscr{U}_\epsilon}\}$ of $\overline{\mathscr{M}}$. □

Applying the choice of z dictated by Lemma 2.1 to (2.4), we see that the last term of the left hand side cancels the right hand side, resulting in

$$\mathrm{Ric}(\bar{g})_{ab} + \frac{d-2}{z}\bar{\nabla}_a\bar{\nabla}_b z + \bar{g}_{ab}\bar{g}^{cd}\frac{1}{z}\bar{\nabla}_c\bar{\nabla}_d z = 0. \tag{2.6}$$

Notice yet that, by formulae (A.22–A.24) (pages 113–115), the multiplication of $\bar{g}$ by a constant $K = \kappa^2 > 0$ doesn't modify neither the Levi-Civita connection, nor the Riemann and Ricci tensors. Hence, we can make the substitution $g \mapsto -\frac{(d-1)(d-2)}{2\Lambda} g = R^2 g$ and, simultaneously, take $\Lambda = -\frac{(d-1)(d-2)}{2}$, which results in $\bar{g}^{-1}(dz, dz) = 1$ in $\mathscr{U}_\epsilon$. In this case, denoting by $Z_a \doteq \bar{\nabla}_a z$ the (unit) normal to the foliation of $\mathscr{U}_\epsilon$ induced by our choice of z, it follows that $K_{ab} \doteq -\bar{\nabla}_a Z_b$ is the extrinsic curvature of $\mathscr{I}_z$. We can, thus, write $\bar{g}$ as

$$\bar{g}(x, z)_{ab} = \bar{g}^{(0)}(x, z)_{ab} + Z_a Z_b,$$

where $\bar{g}^{(0)}(z, .)$ is the metric induced in each of these hypersurfaces ($\bar{g}^{(0)}(x, 0) \doteq \bar{g}^{(0)}(x)$). Noticing that

$$Z^a K_{ab} = -(\bar{\nabla}^a z)\bar{\nabla}_a\bar{\nabla}_b z = -(\bar{\nabla}^a z)\bar{\nabla}_b\bar{\nabla}_a z = -\frac{1}{2}\bar{\nabla}_b(\bar{\nabla}^a z\bar{\nabla}_a z) = 0,$$

we can project (2.6) to its part tangent to the $\mathscr{I}_z$'s:

$$\bar{g}^{(0)}(z)^c_a\bar{g}^{(0)}(z)^d_b\mathrm{Ric}(\bar{g})_{cd} - \frac{d-2}{z}K_{ab} - \frac{1}{z}\bar{g}^{(0)}(z)_{ab}\mathrm{Tr}K = 0. \tag{2.7}$$

where $\mathrm{Tr}K \doteq \bar{g}^{(0)}(z)^{cd}K_{cd}$ is the mean curvature. Finally, applying the Gauss equation (A.3) to $\mathrm{Ric}(\bar{g})$

$$\bar{g}^{(0)}(z)^c_a\bar{g}^{(0)}(z)^d_b\mathrm{Ric}(\bar{g})_{cd} = \mathrm{Riem}(\bar{g})_{acbd}Z^c Z^d + \mathrm{Ric}(\bar{g}^{(0)})_{ab} + K^e_a K_{eb} - (\mathrm{Tr}K)K_{ab}, \tag{2.8}$$

and making use of the identity (analogous to the Riccati equation (A.6))

$$\mathrm{Riem}(\bar{g})_{acbd}Z^cZ^d = Z^e\bar{\nabla}_eK_{ab} - K^e_aK_{eb}, \tag{2.9}$$

we arrive, after multiplication of both sides by z, at

$$z\underbrace{Z^e\bar{\nabla}_eK_{ab}}_{=-\frac{1}{2}\partial_z^2\bar{g}^{(0)}(z)} + z\mathrm{Ric}(\bar{g}^{(0)})_{ab} - z(\mathrm{Tr}K)K_{ab} - (d-2)K_{ab} - \bar{g}^{(0)}(z)_{ab}\mathrm{Tr}K = 0, \tag{2.10}$$

which is the equation of evolution we sought for.[2] Notice, however, that (2.10) possesses constraints, for the contraction of the Einstein equations (2.3) with any vector X^a doesn't have second-order derivatives in the direction of X^a. The contraction of (2.6) and (2.8) with $\bar{g}^{(0)ab}$, plus the Codazzi-Mainardi equation (A.4) together with (2.6), result respectively in the constraint equations (2.11) and (2.12) for the evolution of $(\bar{g}^{(0)}, K)$ dictated by (2.10):

$$R(\bar{g}^{(0)}) + K_{ab}K^{ab} - (\mathrm{Tr}K)^2 + 2\frac{d-2}{z}\mathrm{Tr}K = 0, \tag{2.11}$$

$$\bar{\nabla}^{(0)}_aK^a_b - \bar{\nabla}^{(0)}_b\mathrm{Tr}K = 0. \tag{2.12}$$

The conservation of the constraints (2.11–2.12) along the evolution is guaranteed by the contracted Bianchi identities $\nabla^aG_\Lambda(g)_{ab} = 0$, re-expressed in terms of $\bar{g}$.

(2.10) constitutes a nonlinear, second-order Fuchsian system (see, for instance, [Kic04] for a definition of a Fuchsian system in the nonlinear case), which can be formally solved by *initially* adopting the Taylor-series *ansatz*

$$\bar{g}^{(0)}(z) = \bar{g}^{(0)} + \sum_{j=1}^{\infty} z^j\bar{g}^{(j)}. \tag{2.13}$$

Notice that by taking $z = 0$ in (2.10), we have $K_{ab}\restriction_{\mathscr{I}} = 0$, thus fixing our initial data $(g^{(0)}, 0)$. The coefficients $g^{(j)}$ are obtained by applying ∂_z^{j-1} to both sides of (2.10), resulting in

$$\frac{d-1-j}{2}\partial_z^j\bar{g}^{(0)}(z)_{ab} + \frac{1}{2}\bar{g}^{(0)}(z)_{ab}\bar{g}^{(0)}(z)^{cd}\partial_z^j\bar{g}^{(0)}(z)_{cd} = \\ = (\text{just terms with } \partial_z^k\bar{g}^{(0)}(z),\ k < j). \tag{2.14}$$

[2] This equation is slightly different from the equation of evolution derived in [FG85] and studied in [Gra00, GL91, HSS01]. In fact, Fefferman and Graham depart from considerations different from ours in [FG85] to arrive at their analog of (2.10).

For future use, it's convenient to separate (2.12–2.14) into the trace part $\mathrm{Tr}K$ and the trace-free part $P_{ab} \doteq K_{ab} - \frac{1}{d-1}\bar{g}^{(0)}(z)_{ab}\mathrm{Tr}K$

$$(d-2-j)P_b^{(j)a} = \mathrm{Ric}(\bar{g}^{(0)})_b^{(j-1)a} - \frac{1}{d-1}R(\bar{g}^{(0)})^{(j-1)}\delta_b^a + - \sum_{m=0}^{j-1}(\mathrm{Tr}K)^{(m)}P_b^{(j-1-m)a}, \tag{2.15}$$

$$(2d-3-j)(\mathrm{Tr}K)^{(j)} = R(\bar{g}^{(0)})^{(j-1)} - \sum_{m=0}^{j-1}(\mathrm{Tr}K)^{(m)}(\mathrm{Tr}K)^{(j-1-m)}, \tag{2.16}$$

whence we can obtain by invoking the identity $\partial_z\bar{g}^{(0)}(z)_{ab} = -2\partial_z\bar{g}^{(0)}(z)_{bc}K_a^c$ the explicit form of (2.13)

$$j\bar{g}^{(j)} = -2\sum_{m=0}^{j-1}\left(\bar{g}_{bc}^{(m)}P_a^{(j-1-m)c} + \frac{1}{d-1}\bar{g}_{ab}^{(m)}K^{(j-1-m)}\right). \tag{2.17}$$

The "Hamiltonian" constraint (2.11) guarantees the invariance of the solution of (2.10), up to diffeomorphisms, under reparametrizations of the variable z, and doesn't exert any direct role in the determination of the coefficients of the expansion (2.13).

Hence, we see that $\bar{g}^{(j)}$, $j < d-1$, is uniquely determined by $\bar{g}^{(0)}$ and its tangential derivatives *at* $\mathscr{I}$ (in particular, if $\bar{g}^{(0)}$ is the ESU metric, these coefficients must match the ones of the pure AdS metric in a similar chart in a neighbourhood of $\mathscr{I}$). In particular, as (2.10) is invariant under the parity transformation $z \mapsto -z$, it follows that $\bar{g}^{(j)} = 0$ for j odd, $j < d-1$. For $j = d-1$, the recursion relations given by (2.14) are truncated, and, in this case, we have two possible scenarios:

(i) d even: Once more, parity invariance demands $\mathrm{Tr}\bar{g}^{(d-1)} = 0$, but the trace-free part of $\bar{g}^{(d-1)}$ is only subject to the conservation law $\bar{\nabla}^{(0)a}\bar{g}_{ab}^{(d-1)} = 0$ coming from (2.12).

(ii) d odd: (2.14) nontrivially fixes the trace of $\bar{g}^{(d-1)}$, and (2.12) results in $\bar{\nabla}^{(0)a}\bar{g}_{ab}^{(d-1)} = \bar{\nabla}_b^{(0)}\mathrm{Tr}\bar{g}^{(d-1)}$, thus prescribing the divergence of the trace-free part.

In the case of *d odd*, it's necessary to follow the usual strategy of solution of Fuchsian systems, by modifying the initially adopted *ansatz* so as to include terms in the form $z^k \log z$, $k \geq d-1$ (such terms don't exist for d even). Such an analysis was originally sketched by Fefferman and Graham [FG85] for $\bar{g}^{(0)}$ with arbitrary signature, having been analyzed in the case of Euclidean signature by several authors, among which we cite in particular the works of Graham and Lee [GL91] and Graham [Gra00]. Of great importance was the calculation of the coefficients $\bar{g}^{(j)}$, $j < d-1$, and the

coefficient $\bar{h}^{(d-1)}$ of the term proportional to $z^{d-1}\log z$, quite involved and realized in detail for certain values of d by de Haro, Skenderis and Solodukhin [HSS01]. Thus proceeding, the expansion of $\bar{g}$ around $z=0$, suitably corrected by the above considerations, becomes

$$\text{(i) } d \text{ even:}\quad \bar{g}(z,x) \sim_{z\searrow 0} dz^2+\bar{g}^{(0)}(x)+(z^2)\bar{g}^{(2)}(x)+\cdots+(z^2)^{\left(\frac{d-2}{2}\right)}\bar{g}^{(d-2)}(x)+ \\ +z^{d-1}\bar{g}^{(d-1)}+\cdots; \tag{2.18}$$

$$\text{(ii) } d \text{ odd:}\quad \bar{g}(z,x) \sim_{z\searrow 0} dz^2+\bar{g}^{(0)}(x)+(z^2)\bar{g}^{(2)}(x)+\cdots+(z^2)^{\left(\frac{d-3}{2}\right)}\bar{g}^{(d-3)}(x)+ \\ +(z^{d-1}\log z)\bar{h}^{(d-1)}+z^{d-1}\bar{g}^{(d-1)}+\cdots. \tag{2.19}$$

(we understand (2.18) and (2.19), for now, in a purely asymptotic way – the question of convergence will be discussed later). The coefficient $\bar{h}^{(d-1)}$, denominated Fefferman-Graham tensor, possesses remarkable properties: as $\bar{g}^{(j)}$ with $j<d-1$, it only depends on $\bar{g}^{(0)}$ and its derivatives tangential to $\mathscr{I}$; it's conformally invariant; and vanishes if and only if the conformal class of $g^{(0)}$ has a representative which solves the Einstein equations without matter and zero cosmological constant (as, for instance, ESU_{d-1}). In the latter case, there are no coefficients in the form $z^j\log z$ in (2.19), but this clearly ceases to be true for perturbations of $g^{(0)}$ with compact support and not coming from local conformal rescaling, which affects the calculation of energy-momentum tensors for field theories at the boundary obtained by the AdS/CFT correspondence, causing an anomaly in trace of this tensor to appear and, hence, breaking scale invariance. Such an anomaly, in this context, is called *holographic* Weyl *anomaly*.[HS98, HS00]

Here, instead of detailing these calculations (for which we shall have no use in the remaining of the present work, and which can be found in the above cited references), we shall explore in more detail the remaining components of $\mathrm{Riem}(\bar{g})$, so as to make more precise the part of $\bar{g}^{(d-1)}$ not determined by the recursion relations (2.14). First, we'll simplify (2.6) in terms of K_{ab}. For such, we introduce Schouten *tensor* associated to $\bar{g}$

$$S(\bar{g})_{ab} \doteq \frac{1}{d-2}\left(\mathrm{Ric}(\bar{g})_{ab}-\frac{1}{2(d-1)}R(\bar{g})\bar{g}_{ab}\right).$$

This tensor has a simpler conformal transformation rule and a simpler relation involving $\mathrm{Riem}(\bar{g})$ and $C(\bar{g})$ than the Ricci tensor; the latter becomes

$$\mathrm{Riem}(\bar{g})_{acbd}=C(\bar{g})_{acbd}+2\bar{g}_{a[b}S(\bar{g})_{c]d}-2\bar{g}_{c[b}S(\bar{g})_{d]a}, \tag{2.20}$$

whereas (2.6) becomes

$$S(\bar{g})_{ab}=\frac{1}{z}K_{ab}. \tag{2.21}$$

Gathering (2.9), (2.20) and (2.21) together, we have

$$C(\bar{g})_{acbd}Z^cZ^d = \left(\partial_z - \frac{1}{z}\right)K_{ab} - K^c_a K_{cb}. \tag{2.22}$$

Using th expansion of $\bar{g}^{(0)}(z)$ (2.13) and denoting by $K^{(j)}_{ab} \doteq -\frac{j+1}{2}\bar{g}^{(j+1)}$ the j-th coefficient of the corresponding expansion for K_{ab}, we obtain the coefficients of the expansion of $C(\bar{g})_{acbd}Z^cZ^d$ around $z = 0$:

$$(C(\bar{g})_{acbd}Z^cZ^d)^{(j)} = jK^{(j+1)}_{ab} + \sum_{m=0}^{j} K^{(j-m)c}_a K^{(m)}_{cb}. \tag{2.23}$$

(2.23) explicitly shows that the information lost by the truncation of the recursion relations (2.14) at $j = d-1$, referring to the trace-free part of $\bar{g}^{(d-1)}_{ab}$, is contained in $(C(\bar{g})_{acbd}Z^cZ^d)^{(d-3)}$. Indeed, if we are capable of determining $\bar{g}^{(d-1)}_{ab}$ by other means, we can proceed to higher orders and *completely* determine the coefficients of the expansion (2.14). Thus, we can say that the"correct" initial data for (2.10) are $(\bar{g}^{(0)}, \bar{g}^{(d-1)})$.

We shall now restrict ourselves to the class of AAdS metrics, i.e., $\bar{g}^{(0)}$ is the metric of ESU_{d-1}. In this case, the coefficients $\bar{g}^{(j)}$, $j < d-1$, must match, after an adequate choice of coordinates in $\mathscr{I}$, with those of the metric of the conformal completion of AdS_d (1.15). Hence, a general AAdS metric has in $\mathscr{U}_\epsilon$, after normalization of the cosmological constant, the form

$$ds^2 = \frac{1}{z^2}\left[-\left(1+\frac{z^2}{4}\right)^2 dt^2 + dz^2 + \left(1-\frac{z^2}{4}\right)^2 d\Omega^2_{d-2} + O(z^{d-1})\right]. \tag{2.24}$$

Thus we have from (2.20), (2.21) and (2.24) the following rough "peeling" property for the Weyl tensor: $C(\bar{g})^{(j)}_{abcd} = 0$ for $j < d-3$, and, due to the Codazzi-Mainardi equation (A.4) (page 107), $(\bar{g}^{(0)}(.)^e_a\bar{g}^{(0)}(.)^f_b\bar{g}^{(0)}(.)^h_c Z^d C(\bar{g})_{abcd})^{(j)} = 0$ for $j \leq d-3$. In AdS_d, $d \geq 6$, we know that $\bar{g}^{(d-1)}_{ab} = 0$ and, as AdS_d is a space form, $C(\bar{g})_{abcd} = 0$. Therefore, the sum in the right hand side of (2.23) also vanishes. As it's the same for all AAdS metrics and all $j \leq d-3$, it vanishes for all these cases. Hence,

$$K^{(d-2)}_{ab} = \frac{1}{d-3}(C(\bar{g})_{acbd}Z^cZ^d)^{(d-3)},\ \forall d \geq 6. \tag{2.25}$$

In the case $d = 4$, the sum in the right hand side of (2.23) reduces to $2K^{(1)c}_a K^{(0)}_{cb} = 0$, and for $d = 5$, $K^{(1)c}_a K^{(1)}_{cb} = \frac{1}{8}\bar{g}^{(0)}_{ab}$, by virtue of (2.15–2.17). Thus, we conclude that

$$\bar{g}^{(d-1)} = \begin{cases} -\frac{2}{d-1}E_{ab} & (d = 4 \text{ or } d \geq 6) \\ -\frac{1}{2}E_{ab} + \frac{1}{16}\bar{g}^{(0)}_{ab} & (d = 5) \end{cases}, \tag{2.26}$$

where $E_{ab} = \frac{1}{d-3}\lim_{z\searrow 0} z^{3-d}C(\bar{g})_{acbd}Z^cZ^d$ is the rescaled electric part of the Weyl tensor of $\bar{g}$ (the limit exists and is $\mathscr{C}^\infty$, for we've seen that the coefficients of order inferior to $d-3$ of the expansion of $C(\bar{g})_{acbd}Z^cZ^d$ vanish), which leaves a general AAdS metric satisfying (2.3) in the form

$$\begin{aligned}(2.27)\qquad \bar{g}^{(0)}(z) &= -\left(1+\frac{z^2}{4}\right)^2 dt^2 + dz^2 + \left(1-\frac{z^2}{4}\right)^2 d\Omega^2_{d-2} + \\ &\quad + \left\{ \begin{array}{ll} -\frac{2}{d-1}E_{ab}z^{d-1} & (d=4 \text{ or } d\geq 6) \\ \left(-\frac{1}{2}E_{ab} + \frac{1}{16}\bar{g}^{(0)}_{ab}\right)z^4 & (d=5) \end{array} \right\} + O(z^d).\end{aligned}$$

Remark 2.1 *The problem of convergence of the expansions (2.18–2.19) for $z<\epsilon$ sufficiently small was affirmatively solved, for $\bar{g}^{(0)}_{ab}$ and E_{ab}* real analytic*, by* Kichenassamy *[Kic04]. In the non analytic case, it's not possible, in the present state of the art, to proceed by means of analytic approximations, due to the lack of sufficiently strong estimates of the solution in terms of these boundary data. This, at a first glance, was to be expected, for the* Cauchy *problem for hyperbolic equations is know to be ill posed for initial data at timelike hypersurfaces [Had03]. However, such an intuition is possibly only partially correct, for the "true" initial data for our problem (i.e., $\bar{g}^{(0)}_{ab}$* and *E_{ab}) make the latter considerably different from the usual* Cauchy *problem for second-order hyperbolic equations (remember that $\partial_z\bar{g}^{(0)}(z)\restriction_{z=0} = 0$!) – indeed, weighted estimates involving the* Weyl *tensor exert a fundamental role in the proof global nonlinear stability of* Minkowski *spacetime [CK93]. Partial results in this direction can be found in [And06c, And05, And06b, And06a] and the references cited therein. Hence, in non analytic cases, we should understand the expansions (2.18–2.19) only as asymptotic expansions as $z\searrow 0$.*

Notice, finally, that, by virtue of the purely local character of the analysis made above, it remains valid without change for any Poincaré domain in an AAdS spacetime and, more in general, for any locally AAdS spacetime (the whole discussion preceding formula (2.24) is valid even for general locally AdS-type spacetimes). In the case of Poincaré domains, we must substitute the coordinates (1.8) for the coordinates (1.6), and the metric (1.9) multiplied by z^2 for the metric (1.15). An advantage of this chart in the present context is that $\bar{g}^{(j)}_{ab} = 0$ for $1\leq j\leq d-2$, whence it follows that $\bar{g}^{(d-1)}_{ab} = \frac{2}{d-1}E_{ab}$ for all $d\geq 4$, unlike (2.26). The asymptotic form of an AAdS metric in the neighbourhood of a Minkowski domain then becomes

$$(2.28)\qquad \bar{g}^{(0)}(z)_{ab} = \left(\eta_{ab} - \frac{2}{d-1}z^{d-1}E_{ab}\right) + O(z^d).$$

2.3 Gravitational "thermodynamics" in AAdS wedges

We'll show how to associate to each wedge $\mathscr{W}_{p,q}$ of a locally AAdS spacetime $(\mathscr{M}, g)$, $p, q \in \mathscr{D}(\mathscr{I})$, a flow of diffeomorphisms which extend to $\mathscr{D}_{p,q}$ in such a way that their action coincides in this region with the action of the one-parameter subgroup (1.26). In this case, it follows that these diffeomorphisms are *asymptotic isometries*, i.e., if T^a is the vector field tangent to the flow, then $\pounds_T g(x, z) = o(z)$ as $z \searrow 0$.

The construction we'll make is a particular case of a more general procedure, suitable to any relatively compact diamond $\mathscr{O}_{p,q} = I^+(p) \cap I^-(q)$ in a causally simple spacetime $(\mathscr{M}, g)$, closely related to the method of Geroch [Ger70] for the construction of global time functions in globally hyperbolic spacetimes.

2.3.1 Geometrical time evolution in diamonds

We'll show initially how to associate to each diamond $\mathscr{D}_{p,q}$ at conformal infinity of a locally AAdS spacetime $(\mathscr{M}, g)$, $p, q \in \mathscr{D}(\mathscr{I})$ a global time function associated to the action of the one-parameter subgroup (1.26).

Theorem 2.2 *Consider a d-dimensional AAdS spacetime $(\mathscr{M}, g)$, $r \in \mathscr{I}$ and $p_0, q_0 \in \mathscr{M}in(r)$ such that, if $\mathscr{M}in(r) \ni p \mapsto x^\mu(p)$ is the global Cartesian chart of* Minkowski *spacetime, we have $x^\mu(p_0) = (-1, \mathbf{0})$ and $x^\mu(q_0) = (1, \mathbf{0})$. Consider the one-parameter group $u^\lambda_{p_0,q_0}$ of conformal diffeomorphisms (1.26) of $(\mathscr{D}_{p_0,q_0}, \eta\restriction_{\mathscr{D}_{p_0,q_0}})$, given for p in the $x^0 - x^1$ plane by*

$$x^\pm(u^\lambda_{p_0,q_0}(p)) \doteq \frac{(1 + x^\pm) - e^{-\lambda}(1 - x^\pm)}{(1 + x^\pm) + e^{-\lambda}(1 - x^\pm)},\ \lambda \in \mathbb{R}$$

($x^\pm = x^0 \pm x^1$ are the light cone coordinates in the $x^0 - x^1$ plane) and extended to $p \in \mathscr{D}_{p_0,q_0}$ by spatial rotations. Define the following foliation $\lambda \mapsto \Sigma^\lambda_{p_0,q_0}$ of $\mathscr{D}_{p_0,q_0}$ by Cauchy *surfaces:*

- $\Sigma_{p_0,q_0} \doteq \Sigma^0_{p_0,q_0} = \mathscr{D}_{p_0,q_0} \cap \{x^0 = 0\}$;
- $\Sigma^\lambda_{p_0,q_0} = u^\lambda_{p_0,q_0}(\Sigma^0_{p_0,q_0})$, *i.e., the foliation is given by the diffeomorphism $F_{p_0,q_0} : \mathbb{R} \times \Sigma_{p_0,q_0} \ni (\lambda, p) \mapsto u^\lambda_{p_0,q_0}(p) \in \mathscr{D}_{p_0,q_0}$.*
- *λ is, hence, the global time function associated to F_{p_0,q_0}.*

Then,

$$\lambda = \frac{1}{d-1} \log \left[\frac{|\mathscr{D}_{p_0, u^\lambda_{p_0,q_0}(p)}|}{|\mathscr{D}_{u^\lambda_{p_0,q_0}(p), q_0}|} \right] = \log \left[\frac{d_\eta(p_0, u^\lambda_{p_0,q_0}(p))}{d_\eta(u^\lambda_{p_0,q_0}(p), q_0)} \right],\ \forall p \in \Sigma_{p_0,q_0}, \tag{2.29}$$

where d_η is the Lorentzian distance associated to η (see Appendix A). Recall that the LEBESGUE *measure is precisely the measure induced by the volume element $\sqrt{|\eta|}$ (we notice, for convenience, that $|\mathscr{D}_{p_0,q_0}| = 2\int_0^1 |B_r(0)|dr = 2\int_0^1 \frac{r^{d-2}}{d-2}\mathrm{Vol}S^{d-3}dr = \frac{2}{(d-1)(d-2)}\mathrm{Vol}S^{d-3}$).*

PROOF. Due to the rotational symmetry of $u^\lambda_{p_0,q_0}$, we can restrict our considerations to the $x^0 - x^1$ plane. In this case, let us write $x^\mu(p) = (0, x^1, 0, \ldots, 0)$, $x^1 \in [-1, 1]$, whence it follows that

$$x^\pm(u^\lambda_{p_0,q_0}(p)) = \frac{(1 \pm x^1) - e^{-\lambda}(1 \mp x^1)}{(1 \pm x^1) + e^{-\lambda}(1 \mp x^1)},$$

and, thus,

$$\begin{aligned} x^0(u^\lambda_{p_0,q_0}(p)) &= \frac{1}{2}(x^+(u^\lambda_{p_0,q_0}(p)) + x^-(u^\lambda_{p_0,q_0}(p))) = \\ &= \frac{(1-(x^1)^2)(1-e^{-2\lambda})}{(1-(x^1)^2)(1+e^{-2\lambda}) + 2e^{-\lambda}(1+(x^1)^2)} \end{aligned}$$

and

$$\begin{aligned} x^1(u^\lambda_{p_0,q_0}(p)) &= \frac{1}{2}(x^+(u^\lambda_{p_0,q_0}(p)) - x^-(u^\lambda_{p_0,q_0}(p))) = \\ &= \frac{4x^1 e^{-\lambda}}{(1-(x^1)^2)(1+e^{-2\lambda}) + 2e^{-\lambda}(1+(x^1)^2)}. \end{aligned}$$

Let $q = (t, \mathbf{0})$, $t \in (-1, 1)$. The diamond $\mathscr{D}_{p_0,q}$ is a translation of $\frac{1+t}{2}\mathscr{D}_{p_0,q_0}$, and the diamond $\mathscr{D}_{q,q_0}$, a translation of $\frac{1-t}{2}\mathscr{D}_{p_0,q_0}$ – hence, we have $|\mathscr{D}_{p_0,q}| = \left(\frac{1+t}{2}\right)^{d-1}|\mathscr{D}_{p_0,q_0}|$ and $|\mathscr{D}_{q,q_0}| = \left(\frac{1-t}{2}\right)^{d-1}|\mathscr{D}_{p_0,q_0}|$. More in general, if $q = (t, r, 0, \ldots, 0)$, there exists a LORENTZ boost in the $x^0 - x^1$ plane around p_0 which takes $\mathscr{D}_{p_0,q}$ to $\mathscr{D}_{p_0,q^+}$, where $q^+ = (((1+t)^2 - r^2)^{\frac{1}{2}} - 1, \mathbf{0})$, and a LORENTZ boost in the $x^0 - x^1$ plane around q_0 which takes $\mathscr{D}_{p_0,q}$ to $\mathscr{D}_{q^-,q_0}$, where $q^- = (1 - ((1-t)^2 - r^2)^{\frac{1}{2}}, \mathbf{0})$. As LORENTZ transformations preserve volume, we have

$$|\mathscr{D}_{p_0,q}| = \left(\frac{((1+t)^2 - r^2)^{\frac{1}{2}}}{2}\right)^{d-1} |\mathscr{D}_{p_0,q_0}|$$

and

$$|\mathscr{D}_{q,q_0}| = \left(\frac{((1-t)^2 - r^2)^{\frac{1}{2}}}{2}\right)^{d-1} |\mathscr{D}_{p_0,q_0}|.$$

Finally, taking $q = q(\lambda) = u^\lambda_{p_0,q_0}(p)$, it follows that

$$(1 + x^0(q(\lambda)))^2 - x^1(q(\lambda))^2 = \frac{4(1-(x^1)^2)}{(1-(x^1)^2)(1+e^{-2\lambda}) + 2e^{-\lambda}(1+(x^1)^2)}$$

and

$$(1-x^0(q(\lambda)))^2 - x^1(q(\lambda))^2 = \frac{4e^{-2\lambda}(1-(x^1)^2)}{(1-(x^1)^2)(1+e^{-2\lambda})+2e^{-\lambda}(1+(x^1)^2)},$$

and hence formula (2.29). The second identity follows from the fact that $|\mathscr{D}_{p,q}| = \frac{1}{2^{d-2}(d-1)(d-2)}\mathrm{Vol}S^{d-3}d_\eta(p,q)^{d-1}$. □

The importance of Theorem (2.2) will soon become clear.

It's important to have in mind that, in d-dimensional, strongly causal spacetimes $(\mathscr{M},g)$, the Lorentzian distance d_g coincides with the geodesic distance in convex normal neighbourhoods (see formula 2.31, page 44, and the discussion that follows for more details).

Let us suppose now that $(\mathscr{M},g)$ is causally simple, so as any nonvoid, relatively compact diamond $\mathscr{O}_{p,q}$ is globally hyperbolic, whose Cauchy surfaces $\mathscr{S}_{p,q}$ are acausal sets with edge $\partial I^+(p)\cap\partial I^-(q)$. Notice, however, that if $\overline{\mathscr{O}_{p,q}}$ is not contained in any geodesically convex neighbourhood, then $\mathscr{O}_{p,q}$ is *not* equal to $D(\mathscr{S}_{p,q})$ *in* $\mathscr{M}$, for in this case $\partial\mathscr{O}_{p,q} = (\partial I^+(p)\cap J^-(q))\cup(\partial I^-(q)\cap J^+(p))$ possesses geodesic segments with pairs of conjugate points, that is, there are points in $\partial\mathscr{O}_{p,q}$ which belong to null geodesics which cease to be achronal before reaching the edge of $\mathscr{S}_{p,q}$. Hence, we have that $intD(\mathscr{S}_{p,q}) \supsetneqq \mathscr{O}_{p,q}$ in this case (equality occurs only if the null geodesic segments which generate $\partial\mathscr{O}_{p,q}$ are achronal, as for example in the case that $\overline{\mathscr{O}_{p,q}}$ is contained in some geodesically convex neighbourhood).

Another obstacle to showing that (2.29) indeed generalises to a global time function for a relatively compact $\mathscr{O}_{p,q}$ whose level sets are Cauchy surfaces is that the *second* identity in (2.29) no longer holds if there is curvature present. The intuitive reason is that the "packing" number of small diamonds inside a larger one need not grow linearly with the volume of the latter (this argument can be made rigorous by employing semi-Riemannian volume comparison estimates [ES00]). Hence, Geroch's argument, which does yield a global time function with Cauchy level sets, is not directly applicable. However, simple causality happens to be just strong enough to guarantee that the *first* identity in (2.29) does define a function with the desired properties.

The property we'll use is the reverse triangular inequality (see Appendix A, page 113)

$$p\le r\le q \Rightarrow d_g(p,q)\ge d_g(p,r)+d_g(r,q).$$

It follows from this inequality that, if $\gamma:[0,1]\to\mathscr{M}$ is an inextendible causal curve in $\mathscr{O}_{p,q}$, then $\lambda\mapsto d_g(\gamma(\lambda),q)$ (resp. $\lambda\mapsto d_g(p,\gamma(\lambda))$) is a function bounded

by $d_g(p,q)$ ($< +\infty$ by virtue of the compactness of $\overline{\mathscr{O}_{p,q}}$ and the Definition A.1 of Lorentzian distance) and *strictly* decreasing (resp. increasing) in λ – recall that any *maximal* causal curve (i.e., whose arc length between any two of its points is equal to the Lorentzian distance) is necessarily a geodesic, up to reparametrization. Thus, the level surfaces of $d_g(.,q)$ and $d_g(p,.)$ are *acausal* for nonzero values. Moreover, thanks to causal simplicity, $d_g(p,.)$ (resp. $d_g(.,q)$) tends to zero along any past (resp. future) inextendible causal curve in $\mathscr{O}_{p,q}$. Employing the properties obtained above, we can repeat the proof of Proposition 3.1 of [AGH98b] within our context, arriving at the following

PROPOSITION 2.3 ([AGH98b]) *$d_g(p,.)$ and $d_g(.,q)$ are* semiconvex, *i.e., for any $r \in \mathscr{O}_{p,q}$ there exists a neighbourhood $\mathscr{U} \ni r$, a local chart $x : \mathscr{U} \to \mathbb{R}^d$ and $f \in \mathscr{C}^\infty(\mathscr{U})$ such that $(d_g(p,.)\restriction_{\mathscr{U}} + f) \circ x^{-1}$ and $(d_g(.,q)\restriction_{\mathscr{U}} + f) \circ x^{-1}$ are* convex *in $x(\mathscr{U})$.* □

Actually, the result we obtain is the following: there exist $\phi_{p,r}, \phi_{r,q} \in \mathscr{C}^\infty(\mathscr{U}$ such that $d_g(p,r) = \phi_{p,r}(r)$ and $d_g(r,q) = \phi_{r,q}(r)$, $d_g(p,.) \geq \phi_{p,r}$ and $d_g(.,q) \geq \phi_{r,q}$ in $\mathscr{U}$ and the Hessians $D^2\phi_{p,r}$ and $D^2\phi_{r,q}$ are such that $D^2\phi_{p,r}(r) - c_{p,r}\mathbb{1}$ and $D^2\phi_{r,q}(r) - c_{r,q}\mathbb{1}$ are positive semidefinite matrices for $c_{p,r}, c_{r,q} \in \mathbb{R}$, which not only implies semiconvexity in the sense of Proposition 2.3 [AGH98a], but also guarantees that the given definition is independent of coordinates. In these circumstances, we can invoke the classical result of ALEXANDROV, which tells us that a convex function is not only locally LIPSCHITZ, but is also twice differentiable almost everywhere with respect to LEBESGUE measure (see [EG92] for a proof of this fact). Such a result obviously extends to semiconvex functions. As the restriction of μ_g to normal neighbourhoods is absolutely continuous with respect to LEBESGUE measure, we thus obtain the following

PROPOSITION 2.4 *$d_g(p,.)$ and $d_g(.,q)$ are locally* LIPSCHITZ *and twice differentiable almost everywhere.* □

After this preparations, we can finally define our global time function in $\mathscr{O}_{p,q}$ associated to d_g:

$$\mathscr{O}_{p,q} \ni r \mapsto \lambda^g_{p,q}(r) \doteq \log\left[\frac{d_g(p,r)}{d_g(r,q)}\right] \in \mathbb{R}. \tag{2.30}$$

Our argument above shows that $\lambda^g_{p,q}$ is surjective and strictly increasing along any inextendible, future-directed causal curve $\gamma : (a,b) \to \mathscr{O}_{p,q}$ – as a consequence, $\lambda^g_{p,q}(\gamma(t)) : (a,b) \to \mathbb{R}$ is *surjective* and hence $(\lambda^g_{p,q})^{-1}(\lambda)$ is a CAUCHY surface for all λ, as desired.

We'll show now that the asymptotic behaviour of $\lambda^g_{p,q}(r)$ in the case that $\mathscr{O}_{p,q}$ is contained in a geodesically convex neighbourhood, as

(i) $d_g(p,r) \to 0$ keeping $d_g(r,q) \neq 0$, and

(ii) $d_g(r,q) \to 0$ keeping $d_g(p,r) \neq 0$.

The restriction on the size of $\mathscr{O}_{p,q}$ guarantees that $d_g(p,.)$, $d_g(.,q)$ and, hence, $\lambda^g_{p,q}$ are $\mathscr{C}^\infty$, but we emphasize that formulae that follow depend only on derivatives of order ≤ 2 of $d_g(p,.)$ and $d_g(.,q)$. We shall later discuss other consequences of this hypothesis.

Let $\mathscr{U}$ be a geodesically convex neighbourhood. Let us then define the Synge *world function*

$$\Gamma_g \mathscr{U} \times \mathscr{U} \ni (p,q) \mapsto \Gamma_g(p,q) \doteq -\frac{1}{2}\int_0^1 g(\dot\gamma_{p,q}(s), \dot\gamma_{p,q}(s))ds, \tag{2.31}$$

where $\gamma_{p,q} : [0,1] \to \mathscr{U}$ is the (unique) geodesic segment linking $p = \gamma_{p,q}(0)$ to $q = \gamma_{p,q}(1)$. Γ_g has the following properties (see [Fri75] for the proofs):

- $\Gamma_g \in \mathscr{C}^\infty\left(\bigcup_{p\in\mathscr{M}}\{p\} \times \mathscr{U}_p\right)$, where $\mathscr{U}_p$ is an open, geodesically convex neighbourhood of p;
- $\Gamma_g(p,q) = \Gamma_g(q,p)$;
- $\nabla^a\Gamma_g(p,.) = -\dot\gamma^a_{p,.}(.)$ and $\nabla^a\Gamma_g(.,q) = -\dot\gamma^a_{.,q}(.)$, where . denotes the variable on which ∇ acts. We immediately have the fundamental formula

  $$g^{-1}(d_p\Gamma_g(p,q), d_p\Gamma_g(p,q)) = g^{-1}(d_q\Gamma_g(p,q), d_q\Gamma_g(p,q)) = -2\Gamma_g(p,q), \tag{2.32}$$

  where d_p and d_q denote respectively the differential with respect to the first and second variables.
- $(\nabla_a\nabla_b\Gamma_g(p,.))(p) = -g_{ab}(p)$.

First, we must notice that d_g and $2(\Gamma_g)^{\frac{1}{2}}$ coincide in $\mathscr{U} \times J^+(\mathscr{U},\mathscr{U})$, for $(\mathscr{M},g)$ is causally simple by hypothesis and, thus, strongly causal (see Theorem 4.27 in [BEE96]). $(\Gamma_g)^{\frac{1}{2}}$ is continuous but non differentiable in $d_g^{-1}(0) \cap (\mathscr{U} \times J^+(\mathscr{U},\mathscr{U})$, by virtue of the presence of the square root. Hence, we employ the equivalent formula

$$\lambda^g_{p,q}(r) = \frac{1}{2}(\log(d_g(p,r)^2) - \log(d_g(r,q)^2)) = \frac{1}{2}(\log(\Gamma_g(p,r)) - \log(\Gamma_g(r,q))) \tag{2.33}$$

instead of (2.30) in what follows. We assume, from now on, p and q *fixed* and $\Gamma_g(p,r)$ and $\Gamma_g(r,q)$ as functions of r *alone*, that is, differentials and covariant derivatives act *only on* r. In this way, we can unclutter the notation quite a bit.

We'll first obtain the infinitesimal form of the diffeomorphism flow induced by the global time function $\lambda^g_{p,q}$:

$$\nabla_a \lambda^g_{p,q}(r) = \frac{1}{2}\left(\frac{\nabla_a \Gamma_g(p,r)}{\Gamma_g(p,r)} - \frac{\nabla_a \Gamma_g(r,q)}{\Gamma_g(r,q)}\right). \tag{2.34}$$

Let us now consider the vector field T^a in $\mathscr{O}_{p,q}$ uniquely determined by the following algebraic conditions:

- $T^a \nabla_a \lambda^g_{p,q}(r) = 1$ for all $r \in \mathscr{O}_{p,q}$;
- $g(T, X) = 0$ for all X^a tangent to $(\lambda^g_{p,q})^{-1}(t)$, $t \in \mathbb{R}$.

In other words, T^a is the vector field which generates the flow of diffeomorphisms induced by the foliation of $\mathscr{O}_{p,q}$ by the level hypersurfaces of $\lambda_{p,q}$. Hence, we have

$$g(T,T) = \frac{1}{g^{-1}(d\lambda^g_{p,q}, d\lambda^g_{p,q})}, \tag{2.35}$$

where

$$g^{-1}(d\lambda^g_{p,q}, d\lambda^g_{p,q}) = -\frac{1}{2}\left(\frac{1}{\Gamma_g(p,r)} + \frac{1}{\Gamma_g(r,q)} + \frac{g^{-1}(d\Gamma_g(p,r), d\Gamma_g(r,q))}{\Gamma_g(p,r)\Gamma_g(r,q)}\right). \tag{2.36}$$

Thus,

$$\begin{aligned} T^a &= \frac{1}{g^{-1}(d\lambda^g_{p,q}, d\lambda^g_{p,q})} \nabla^a \lambda^g_{p,q} = \\ &= \frac{\Gamma_g(p,r)\nabla^a \Gamma_g(r,q) - \Gamma_g(r,q)\nabla^a \Gamma_g(p,r)}{\Gamma_g(p,r) + \Gamma_g(r,q) + g^{-1}(d\Gamma_g(p,r), d\Gamma_g(r,q))}. \end{aligned} \tag{2.37}$$

It's clear from (2.37) that, asymptotically as $r \to p$ and $r \to q$, that is, respectively (—) $d_g(p,r) \searrow 0$, $d_g(r,q) \nearrow d_g(p,q)$ and (+) $d_g(r,q) \searrow 0$, $d_g \nearrow d_g(p,q)$, T^a approaches a conformal KILLING field, i.e.,

$$\begin{aligned} \nabla_a T_b + \nabla_b T_a &= f(.) g_{ab} + o(\min\{d_g(p,r), d_g(r,q)\}) = \\ &= \frac{2}{d}(\nabla_c T^c) g_{ab} + o(\min\{d_g(p,r), d_g(r,q)\}) \end{aligned} \tag{2.38}$$

as $r \to p$ or $r \to q$. More precisely, T^a (resp. $-T^a$) approaches an infinitesimal *dilation* around p (resp. q). In geodesic normal coordinates (x^μ) around p (resp. q), the components of T^a become

$$T^\mu = x^\mu - x^\mu(p) + O((x - x(p))^2) \text{ (resp. } x^\mu - x^\mu(q) + O((x - x(q))^2)). \tag{2.39}$$

2.3.2 Construction of asymptotic isometries

Once more specializing ourselves to (locally) AdS-type spacetimes $(\mathscr{M}, g)$, we see that, due to the fact that $(\mathscr{I}, \bar{g}^{(0)})$ is totally geodesic with respect to $(\overline{\mathscr{M}}, \bar{g})$ (and, hence, $\frac{1}{2}d^2_{\bar{g}}\restriction_{\mathscr{I}^2\cap\mathscr{U}^2} = \Gamma_{\bar{g}}\restriction_{\mathscr{I}^2\cap\mathscr{U}^2} = \Gamma_{\bar{g}^{(0)}}\restriction_{\mathscr{I}^2\cap\mathscr{U}^2}$, where $\mathscr{U}$ is a geodesically convex neighbourhood in $\overline{\mathscr{M}}$ – see Proposition 4.32 in [BEE96]), $\lambda^{\bar{g}}_{p,q}\restriction_{\mathscr{I}} \equiv \lambda^{\bar{g}^{(0)}}_{p,q}$ for any pair $p \ll_{\mathscr{I}} q \in \mathscr{I}$ such that $\overline{\mathscr{W}_{p,q}}$ is contained in a geodesically convex neighbourhood in $(\overline{\mathscr{M}}, \bar{g})$, for $\lambda^{\bar{g}}_{p,q}$ is constructed solely from $\Gamma_{\bar{g}}$. In particular, if $(\mathscr{M}, g)$ is (locally) AAdS, and $x(p) = (-1, \mathbf{0})$ and $x(q) = (1, \mathbf{0})$ in a Cartesian chart $x : (\mathscr{M}in(r), \bar{g}^{(0)}\restriction_{\mathscr{M}in(r)}) \to (\mathbb{R}^{1,d-2}, \eta)$ on a Minkowski domain $\mathscr{M}in(r)$ form some $r \in \mathscr{I}$, it follows that $\lambda^{\bar{g}}_{p,q}\restriction_{\mathscr{I}} \equiv \lambda$, where λ is given by formula (2.29), which is the conclusion of Theorem 2.2. That is, *T^a is not only tangent to $\mathscr{I}$ as $T^a\restriction_{\mathscr{I}}$ coincides with the vector field in $\mathscr{D}_{p,q}$ generated by the one-parameter subgroup $\lambda \mapsto u^{\lambda}_{p_0,q_0}$ of the conformal group of $(\mathscr{I}, \bar{g}^{(0)})$ given by (1.23).* Extending the assertion of Theorem 2.2 to $(p, q) \in \mathscr{D}(\mathscr{I})$ by the commutation relations of $SO_e(2, d-1)$ and recalling that (resp. the connected component to the identity of) the conformal group is precisely the set of diffeomorphisms which preserve the causal structure (resp., plus orientation and time orientation), we obtain:

Proposition 2.5 *Let $(\mathscr{M}, g)$ be a locally AAdS spacetime, $(p, q) \in \mathscr{D}(\mathscr{I})$ such that $\overline{\mathscr{W}_{p,q}}$ is contained in a geodesically convex neighbourhood in $(\overline{\mathscr{M}}, \bar{g})$. Then T^a is tangent to $\mathscr{I}$ and $\lambda^{\bar{g}}_{p,q}\restriction_{\mathscr{I}} = \lambda^{\bar{g}^{(0)}}_{p,q} = \lambda$, where $u^{\lambda}_{p,q}$ is the isotropy subgroup (1.26) of $\mathscr{D}_{p,q}$ of the conformal group of $(\mathscr{I}, \bar{g}^{(0)})$.* □

It follows from Proposition 2.5 that T^a generates a one-parameter group of *asymptotic isometries* of $(\mathscr{W}_{p,q}, g\restriction_{\mathscr{W}_{p,q}})$, i.e., $\pounds_T g = o(z)$ as $z \searrow 0$.

It's important to notice that, *in the case of AdS_d*, we see from the proof of Theorem 2.2 and by employing Poincaré charts that T^a is *precisely* the Killing field in $\mathscr{W}_{p,q}$ which generates the conformal isotropy subgroup of the latter, for any $(p, q) \in \mathscr{D}(\mathscr{I})$. Hence, our construction naturally generalizes the Killing fields of AdS_d.

Now what happens with "large" $\mathscr{W}_{p,q}$'s, i.e., such that $\overline{\mathscr{W}_{p,q}}$ is relatively compact in $\overline{\mathscr{M}}$, but *not* contained in some geodesically convex neighbourhood in $(\overline{\mathscr{M}}, \bar{g})$? We have problems of two sorts, both stemming from the same source (namely, the critical points of the exponential map), but with different consequences, more or less severe for us:

1. L_g ceases to be well defined, and d_g ceases to be $\mathscr{C}^\infty$ in $d_g(p, .)^{-1}((0, d_g(p, q)))$ and $d_g(., q)^{-1}((0, d_g(p, q)))$. However, as we know (Proposition 2.4) that d_g is twice differentiable $d_g(p, .)^{-1}((0, d_g(p, q)))$ and $d_g(., q)^{-1}((0, d_g(p, q)))$ up to some set $\mathscr{U}$ satisfying $\mu_g(\mathscr{U}) = \mu_{d_g}(\mathscr{U}) = 0$, all formulae we've obtained remain valid in the points where $d_g(p, .)^2$ and $d_g(., q)^2$ are twice differentiable. We cannot, however,

guarantee the validity of the limits of expressions involving derivatives of $d_g(.,q)^2$ as $r \to p$, or involving derivatives of $d_g(p,.)^2$ as $r \to q$, since we know nothing about the continuity of these outside geodesically convex neighbourhoods.

2. The second problem is specific of (locally) AAdS spacetimes. It may occur that (some) *maximal* (with respect a $d_{\bar{g}}$) timelike geodesic linking p to q traverses $\mathscr{M}$ instead of remaining at $\mathscr{I}$. This implies not only that T^a is no longer tangent to $\mathscr{I}$, but also causes T^a to acquire discontinuity points at $\mathscr{I}$. The hypothesis that such *focusing* of timelike geodesics doesn't occur is stronger and cannot be obtained as a consequence of all null geodesics in $\overline{\mathscr{M}}$ with endpoints in $\mathscr{I}$ and traversing $\mathscr{M}$ having a pair of conjugate points, employed for instance in Theorem 1.3. We can, of course, impose the hypothesis that maximal causal geodesics in $\overline{\mathscr{M}}$ connecting points p and q necessarily belong to $\mathscr{I}$, along lines similar to what we've done for null geodesics along Chapter 1. However, this hypothesis has the annoying characteristic of depending globally on the conformal factor z, unlike the weaker case of null geodesics, which depends only on the causal structure.

A way of circumventing these problems is to *regularize* $\lambda^g_{p,q}$ in a way that this regularization, here denoted $\bar{\lambda}^g_{p,q}$ for concreteness, satisfies $\bar{\lambda}^g_{p,q}\restriction_{\mathscr{D}_{p,q}} = \lambda$, where $u^\lambda_{p,q}$ is given by (1.26). Here the problem is how to regularize a global time function defined in a spacetime *with boundary* – we shall solve the problem in the following manner: let us define the *double* ${}^2\overline{\mathscr{M}}$ of a locally AdS-type spacetime $\overline{\mathscr{M}}$ as the disjoint union of two copies of $\overline{\mathscr{M}}$ with the points in $\mathscr{I}$ identified. ${}^2\overline{\mathscr{M}}$ can be endowed with a differentiable structure compatible with the one of each copy and such that the natural inclusion of each copy is a differentiable embedding [Hir76]. The metric in ${}^2\overline{\mathscr{M}}$ induced by $\bar{g}$ (also denoted by $\bar{g}$, as there's no confusion) is $\mathscr{C}^\infty$ in the interior of each copy, and $\mathscr{C}^{d-1,1}$ (resp. $\mathscr{C}^\infty$) in $\mathscr{I}$ for d odd (resp. d even [3]), due to the Fefferman-Graham expansion for locally AdS-type spacetimes constructed in the previous Section – the regularity is $\mathscr{C}^\infty$ for arbitrary d if $(\mathscr{M}, g)$ is locally AAdS or, more in general, if $\bar{g}^{(0)}$ possesses a representative $\bar{g}^{(0)'}$ of its conformal class which satisfies $\mathrm{Ric}(\bar{g}^{(0)'}) = 0$. This construction is a generalization of the conformal identification of AdS_d with half of ESU_d, seen in Chapter 1, page 9.

We can extend z a $z < 0$ in ${}^2\overline{\mathscr{M}}$, thus defining a system of Gaussian normal coordinates around $\mathscr{I}$. The double $({}^2\overline{\mathscr{M}}, \bar{g})$ has a group of *global* isometries, isomorphic to $\mathbb{Z}_2$, generated by the reflection ζ across $\mathscr{I}$ of a point in a copy of $\mathscr{M}$, having as image its counterpart in the other copy (locally close to $\mathscr{I}$, we have $z \circ \zeta = -z$).

[3] The extra regularity owes itself to (i) the absence of log terms in the Fefferman-Graham expansion and (ii) the parity symmetry in z in formula (2.10), page 35, which gives origin to the recursion relations.

We shall now define $\bar{\lambda}^{\bar{g}}_{p,q}$ *in a causally simple AAdS spacetime* $(\mathscr{M}, g)$ *which satisfies the focusing hypothesis of Theorem 1.3, page 16* (these requirements guarantee, as seen in Chapter 1, that $I^+(p, \overline{\mathscr{M}}) \cap I^-(q, \overline{\mathscr{M}}) \cap \mathscr{I} = \mathscr{D}_{p,q}$), in the following steps:

- First, we define $\lambda^{\bar{g}}_{p,q}$ *in* $({}^2\overline{\mathscr{M}}, \bar{g})$ (more precisely, in $\mathscr{O}_{p,q} = \mathscr{W}_{p,q} \cup \mathscr{D}_{p,q} \cup \zeta(\mathscr{W}_{p,q})$), with the same formula.

- Next, we regularize $\lambda^{\bar{g}}_{p,q}$ by employing the method of Bernal and Sánchez [BS03, BS05] (see [Sán05] for an introductory sketch of the method), obtaining a $\mathscr{C}^\infty$ global time function $\tilde{\lambda}^{\bar{g}}_{p,q}$ which foliates $\mathscr{O}_{p,q}$ by Cauchy surfaces. We assume yet that the obtained regularization satisfies $|\tilde{\lambda}^{\bar{g}}_{p,q} - \lambda^{\bar{g}}_{p,q}| < \delta$ for $\delta < 1$ sufficiently small [Sei77].

- Let us define $\hat{\lambda}^{\bar{g}}_{p,q} = \frac{1}{2}(\tilde{\lambda}^{\bar{g}}_{p,q} + \tilde{\lambda}^{\bar{g}}_{p,q} \circ \zeta)$. Since $\tilde{\lambda}^{\bar{g}}_{p,q}$ as well as $\tilde{\lambda}^{\bar{g}}_{p,q} \circ \zeta$ are global time functions in $\mathscr{O}_{p,q}$ whose level hypersurfaces are Cauchy surfaces in $\mathscr{O}_{p,q}$, and the collection of such functions is closed under convex linear combinations [Sei77], it follows that $\hat{\lambda}^{\bar{g}}_{p,q}$ also satisfies these properties, plus the key additional property (which soon will be crucial) $\partial_z \hat{\lambda}^{\bar{g}}_{p,q} \restriction_{\mathscr{I}} = 0$ and the estimate in the previous item. Henceforth, the usefulness of the double is over – we restrict $\hat{\lambda}^{\bar{g}}_{p,q}$ to $\overline{\mathscr{M}}$, and proceed only within this manifold.

- We extend $\lambda^{\bar{g}^{(0)}}_{p,q}$ from $\mathscr{D}_{p,q}$ to a collar neighbourhood $\mathscr{U}^{\delta'}_{p,q} \cong \mathscr{D}_{p,q} \times [0, 0 < \delta') \ni (x, w)$ of the latter in $\mathscr{W}_{p,q}$ by using the formula $\lambda^{\bar{g}^{(0)}}_{p,q}(x, w) = \lambda^{\bar{g}^{(0)}}_{p,q}(x)$, taking δ' small enough so that $\bar{\nabla}^a \lambda^{\bar{g}^{(0)}}$ remains timelike in $\mathscr{U}^{\delta'}_{p,q}$ and keeping the same notation for the extension (notice that such a restriction to δ' is independent of the restriction to δ in the first item).

- We choose a partition of unity $\{\phi, 1 - \phi\}$ of $\mathscr{W}_{p,q} \cup \mathscr{D}_{p,q}$ subordinated to the covering $\{\mathscr{D}_{p,q} \times [0, \frac{3\delta'}{4}), \mathscr{W}_{p,q} \setminus (\mathscr{D}_{p,q} \times [0, \frac{\delta'}{2}])\}$, satisfying $|\partial_w \phi| \leq K(\delta')^{-1}$ [Whi34] and *independent of* $x \in \mathscr{D}_{p,q}$ (as $\phi \equiv 0$ outside $\mathscr{U}^{\delta'}_{p,q}$, there are no problems with the second property). Let us define

$$\bar{\lambda}^{\bar{g}}_{p,q} \doteq \phi \lambda^{\bar{g}^{(0)}}_{p,q} + (1 - \phi)\hat{\lambda}^{\bar{g}}_{p,q}, \tag{2.40}$$

  which satisfies

$$d\bar{\lambda}^{\bar{g}}_{p,q} = (\lambda^{\bar{g}^{(0)}}_{p,q} - \hat{\lambda}^{\bar{g}}_{p,q})\partial_w \phi dw + \phi d\lambda^{\bar{g}^{(0)}}_{p,q} + (1 - \phi) d\hat{\lambda}^{\bar{g}}_{p,q}. \tag{2.41}$$

- We see immediately that we need to control the size of $|\lambda^{\bar{g}^{(0)}}_{p,q} - \hat{\lambda}^{\bar{g}}_{p,q}|$ in $\mathscr{U}^{\delta'}_{p,q}$ so that $|\bar{\lambda}^{\bar{g}^{(0)}}_{p,q} - \hat{\lambda}^{\bar{g}}_{p,q}|$ is as small as we want and, at the same time, $d\bar{\lambda}^{\bar{g}^{(0)}}_{p,q}$ is a future timelike covector field; the most severe obstacle to such requirements resides in the

spacelike covector proportional to $\partial_w \phi$ in (2.41), due to the WHITNEY estimate invoked in the former item, which is essentially optimal. We write, thus,

$$(2.42)\quad |\lambda^{\bar{g}^{(0)}}_{p,q}(x,w) - \hat{\lambda}^{\bar{g}}_{p,q}(x,w)| \leq \underbrace{|\lambda^{\bar{g}^{(0)}}_{p,q}(x) - \hat{\lambda}^{\bar{g}}_{p,q}(x,0)|}_{<\delta\,(a)} + \underbrace{|\hat{\lambda}^{\bar{g}}_{p,q}(x,w) - \hat{\lambda}^{\bar{g}}_{p,q}(x,0)|}_{\leq K'\delta'^2\,(b)},$$

where the estimate (a) is valid by construction, and the estimate (b) follows from the fact that $\hat{\lambda}^{\bar{g}}_{p,q}$ is $\mathscr{C}^\infty$ and $\partial_w \hat{\lambda}^{\bar{g}}_{p,q} \restriction_{\mathscr{D}_{p,q}} \equiv 0$.

- Taking $\delta = K'(\delta')^2$ and δ' sufficiently small, it follows that $\bar{\lambda}^{\bar{g}}_{p,q}$ is a $\mathscr{C}^\infty$ global time function which:
  1. satisfies $|\bar{\lambda}^{\bar{g}}_{p,q} - \lambda^{\bar{g}}_{p,q}| < \delta''$, where $\delta'' > 0$ is as small as we like,
  2. extends $\lambda^{\bar{g}^{(0)}}_{p,q}$ to $\mathscr{W}_{p,q} \cup \mathscr{D}_{p,q}$, and such that
  3. $\bar{\nabla}^a \bar{\lambda}^{\bar{g}}_{p,q}$ is a future directed timelike vector field, tangent to $\mathscr{I}$, and
  4. $(\bar{\lambda}^{\bar{g}}_{p,q})^{-1}(t)$ is a CAUCHY surface for $\mathscr{W}_{p,q} \cup \mathscr{D}_{p,q}$ for all $t \in \mathbb{R}$ (it follows directly from the first property [Die88] and the continuity of $\bar{\lambda}^{\bar{g}}_{p,q}$).
- By virtue of the results above, it follows that

$$(2.43)\qquad \bar{T}^a \doteq \frac{1}{\bar{g}^{-1}(d\bar{\lambda}^{\bar{g}}_{p,q}, d\bar{\lambda}^{\bar{g}}_{p,q})} \bar{\nabla}^a \bar{\lambda}^{\bar{g}}_{p,q}$$

is an asymptotic KILLING field in $(\mathscr{W}_{p,q}, g \restriction_{\mathscr{W}_{p,q}})$ satisfying $\bar{T}^a \restriction_{\mathscr{I}} = \frac{du^{\cdot}_{p,q}}{d\lambda}$, and, hence, the formulae (2.38) and (2.39), as desired.

Although $\bar{\lambda}^{\bar{g}}_{p,q}$ all the properties we need, unfortunately it no longer possesses in $(\mathscr{M}, g)$ the natural geometrical interpretation which $\lambda^{\bar{g}}_{p,q}$ does, though being as close to this as we desire. Nevertheless, we've shown that it's possible to explicitly build families of asymptotic isometries in (practically) any AAdS wedge. We shall make extensive use of this fact in Chapter 4.

2.3.3 "Return to equilibrium" in wedges

In the work of MARTINETTI and ROVELLI [MR03], it was proposed to associate an "UNRUH effect" to the group $u^{\lambda}_{p,q}$ associated to a diamond $\mathscr{D}_{p,q}$ in MINKOWSKI spacetime, whose orbits were imagined as the worldlines of "observers with finite lifetime". For quantum field theories with conformally invariant vacuum, it's truly possible to prove such an association [BGL93], but a proper discussion of this issue will have to await Chapter 4. Nevertheless, formulae (2.38) and (2.39) make natural the following question: *what if such a temperature were attained only in the limit* $\lambda \to \pm\infty$ *?* For

quantum field theories which enjoy asymptotic freedom, scale invariance is indeed realized only in the short-distance limit, and the corrections to "critical" behaviour are given, for instance, by the operator product expansion (OPE). Thus, we can think that the "thermal time hypothesis" of Martinetti and Rovelli, cited above, is realized only in the weaker form of a *return to equilibrium*, by means of an *asymptotic* scale invariance (we shall give a more precise sense to these considerations in Chapter 4).

We'll transplant this reasoning to AAdS wedges. From the viewpoint of the laws of black hole dynamics, if we imagine the boundary of a wedge (supposed geodesically convex, for simplicity) $\mathscr{W}_{p,q}$ as an "asymptotically stationary" *event horizon*, and $\mathscr{W}_{p,q}$ as a "domain of outer communications", we can ask ourselves if there exists some asymptotic analog for these laws. It's now our objective to show that this indeed occurs, at least for the zeroth law.

First, we shall give the precise definition of *surface gravity* within our context. Let T^a be as in (2.37). Taking, for instance, $r \to r_- \in \partial_-\mathscr{W}_{p,q}$, we have $\Gamma_{\bar{g}}(p, r_-) = 0$, $\Gamma_{\bar{g}}(r_-, q) \neq 0$ and $d\Gamma_{\bar{g}}(p, r_-) \neq 0$. Moreover,

$$\lim_{r\to r_-} \bar{g}(T,T) = -\lim_{r\to r_-} \frac{2\Gamma_{\bar{g}}(p,r)\Gamma_{\bar{g}}(r,q)}{\Gamma_{\bar{g}}(p,r)+\Gamma_{\bar{g}}(r,q)+\bar{g}^{-1}(d\Gamma_{\bar{g}}(p,r), d\Gamma_{\bar{g}}(r,q))} = 0 \tag{2.44}$$

by construction. Equally, if $r \to r_+ \in \partial_+\mathscr{W}_{p,q}$, we have $\Gamma_{\bar{g}}(r_+, q) = 0$, $\Gamma_{\bar{g}}(p, r_+) \neq 0$, $d\Gamma_{\bar{g}}(r_+, q) \neq 0$ and $\lim_{r\to r_+} \bar{g}(T,T) = 0$. In the case that $r \to r_0 \in \partial I^+(p, \overline{\mathscr{M}}) \cap \partial I^-(q, \overline{\mathscr{M}})$, we have $T^a \to 0$. Summing up, T^a is, at the same time, *tangent* and *normal* to the past horizon $\partial_-\mathscr{W}_{p,q}$ and to the future horizon $\partial_+\mathscr{W}_{p,q}$ of $\mathscr{W}_{p,q}$ (notice that the definition of horizon we've given in page 15 excludes precisely the points of $\partial\mathscr{W}_{p,q}$ where $T^a \to 0$). Hence, we have

$$\lim_{\to\partial_\pm\mathscr{W}_{p,q}} \bar{\nabla}_a\bar{g}(T,T) \doteq -2\bar{\kappa}_\pm(\lim_{\to\partial_\pm\mathscr{W}_{p,q}} T_a), \tag{2.45}$$

where $\bar{\kappa}_+$ (resp. $\bar{\kappa}_-$) is a $\mathscr{C}^\infty$ function on the submanifold $\partial_+\mathscr{W}_{p,q}$ (resp. $\partial_-\mathscr{W}_{p,q}$), given by

$$\bar{\kappa}_\pm = -\lim_{\to\partial_\pm\mathscr{W}_{p,q}} \frac{T^a\bar{\nabla}_a\bar{g}(T,T)}{2\bar{g}(T,T)} = \frac{g^{-1}(d\lambda^{\bar{g}}_{p,q}, d\bar{g}^{-1}(d\lambda^{\bar{g}}_{p,q}, d\lambda^{\bar{g}}_{p,q}))}{2\bar{g}^{-1}(d\lambda^{\bar{g}}_{p,q}, d\lambda^{\bar{g}}_{p,q})^2}. \tag{2.46}$$

Let us suppose for the moment that T^a is a conformal Killing field with respect to $\bar{g}$. In this case,

$$\begin{aligned} \bar{\nabla}_a\bar{g}(T,T) &= 2T^b\bar{\nabla}_aT_b = 2T^b(\bar{\nabla}_aT_b + \bar{\nabla}_bT_a) - 2T^b\bar{\nabla}_bT_a = \\ &= \frac{4}{d}(\bar{\nabla}_bT^b)T_a - 2T^b\bar{\nabla}_bT_a \overset{\to\partial_\pm\mathscr{O}_{p,q}}{\longrightarrow} -2\bar{\kappa}_\pm T_a, \end{aligned} \tag{2.47}$$

whence it follows that

$$\kappa_\pm \doteq \lim_{\to\partial_\pm\mathscr{W}_{p,q}} \frac{T^aT^b\bar{\nabla}_bT_a}{\bar{g}(T,T)} \doteq \lim_{\to\partial_\pm\mathscr{W}_{p,q}} \frac{T^aT^b\bar{\nabla}_aT_b}{\bar{g}(T,T)} + \bar{\kappa}_\pm = \lim_{\to\partial_\pm\mathscr{W}_{p,q}} \left(\frac{2}{d}\bar{\nabla}_aT^a\right) + \bar{\kappa}_\pm \tag{2.48}$$

defines precisely the failure of the parametrization of the orbits of T^a in $\partial_\pm\mathscr{D}_{p,q}$ to be an *affine* parametrization, since T^a is tangent to the null geodesics which generate $\partial_+\mathscr{W}_{p,q}$ and $\partial_-\mathscr{W}_{p,q}$. More precisely, if λ_- (resp. λ_+) is an affine parameter common to the future directed null geodesics which emanate from p (resp. reach q) such that $\lambda_- = 0$ at p (resp. $\lambda_+ = 0$ at q), we have

$$\lambda_\pm = \mp e^{\kappa_\pm\lambda^g_{p,q}}. \tag{2.49}$$

Another interpretation for $\kappa_\pm$, which follows from formulae (2.46)–(2.48), is the acceleration needed to maintain a test body at a fixed, but "infinitesimally small" distance, from the horizon, compensated by the redshift factor *"redshift"* $(-\bar{g}(T,T))^{\frac{1}{2}}$ of the orbit [Wal84, Wal94].

Definition 2.2 *κ_- (resp. κ_+) is said to be the* past *(resp.* future*) m*surface gravity *of $\mathscr{O}_{p,q}$.*

For arbitrary T^a, we use the first (two) identity(ies) in (2.48) to *define* $\kappa_\pm$, but the last identity in formula (2.48) is only *asymptotically* valid, as $r_- \to p$, $r_+ \to q$. With this caveat in mind, we can now formulate our "zeroth law of dynamics":

Theorem 2.6 (Zeroth Law of Dynamics of AAdS) *Let $(\mathscr{M}, g)$ be a (locally) AAdS spacetime, and $\mathscr{W}_{p,q}$ a wedge contained in a geodesically convex neighbourhood in $(\overline{\mathscr{M}}, g)$. Then* $\lim_{\to p}\kappa_- = -\lim_{\to q}\kappa_+ = 1$.

Proof. A direct calculation employing the properties of $\Gamma_{\bar{g}}$ gives us

$$T^a\bar{\nabla}_a\bar{g}(T,T) = \frac{2\Gamma_{\bar{g}}(p,r)\Gamma_{\bar{g}}(r,q)}{(\Gamma_{\bar{g}}(p,r)+\Gamma_{\bar{g}}(r,q)+\bar{g}^{-1}(d\Gamma_{\bar{g}}(p,r),d\Gamma_{\bar{g}}(r,q)))^2}\cdot$$
$$\cdot\left[(\Gamma_{\bar{g}}(r,q)-\Gamma_{\bar{g}}(p,r))\left(2+\frac{\bar{g}^{-1}(d\Gamma_{\bar{g}}(p,r),d\Gamma_{\bar{g}}(r,q))}{\Gamma_{\bar{g}}(p,r)+\Gamma_{\bar{g}}(r,q)+\bar{g}^{-1}(d\Gamma_{\bar{g}}(p,r),d\Gamma_{\bar{g}}(r,q))}\right)+\right.$$
$$\left.-\frac{\Gamma_{\bar{g}}(r,q)\bar{\nabla}^a\Gamma_{\bar{g}}(p,r)\bar{\nabla}^b\Gamma_{\bar{g}}(p,r)\bar{\nabla}_a\bar{\nabla}_b\Gamma_{\bar{g}}(r,q)-\Gamma_{\bar{g}}(p,r)\bar{\nabla}^a\Gamma_{\bar{g}}(r,q)\bar{\nabla}^b\Gamma_{\bar{g}}(r,q)\bar{\nabla}_a\bar{\nabla}_b\Gamma_{\bar{g}}(p,r)}{\Gamma_{\bar{g}}(p,r)+\Gamma_{\bar{g}}(r,q)+\bar{g}^{-1}(d\Gamma_{\bar{g}}(p,r),d\Gamma_{\bar{g}}(r,q))}\right],$$

and, thus,

$$-\frac{T^a\bar{\nabla}_a\bar{g}(T,T)}{2\bar{g}(T,T)} = -\frac{1}{2(\Gamma_{\bar{g}}(p,r)+\Gamma_{\bar{g}}(r,q)+\bar{g}^{-1}(d\Gamma_{\bar{g}}(p,r),d\Gamma_{\bar{g}}(r,q)))}\cdot$$
$$\cdot\left[(\Gamma_{\bar{g}}(r,q)-\Gamma_{\bar{g}}(p,r))\left(2+\frac{\bar{g}^{-1}(d\Gamma_{\bar{g}}(p,r),d\Gamma_{\bar{g}}(r,q))}{\Gamma_{\bar{g}}(p,r)+\Gamma_{\bar{g}}(r,q)+\bar{g}^{-1}(d\Gamma_{\bar{g}}(p,r),d\Gamma_{\bar{g}}(r,q))}\right)+\right.$$

$$-\frac{\Gamma_{\bar{g}}(r,q)\bar{\nabla}^a\Gamma_{\bar{g}}(p,r)\bar{\nabla}^b\Gamma_{\bar{g}}(p,r)\bar{\nabla}_a\bar{\nabla}_b\Gamma_{\bar{g}}(r,q)-\Gamma_{\bar{g}}(p,r)\bar{\nabla}^a\Gamma_{\bar{g}}(r,q)\bar{\nabla}^b\Gamma_{\bar{g}}(r,q)\bar{\nabla}_a\bar{\nabla}_b\Gamma_{\bar{g}}(p,r)}{\Gamma_{\bar{g}}(p,r)+\Gamma_{\bar{g}}(r,q)+\bar{g}^{-1}(d\Gamma_{\bar{g}}(p,r),d\Gamma_{\bar{g}}(r,q))}\Bigg].$$

$\bar{\kappa}_+$ (resp. $\bar{\kappa}_-$) is obtained by taking the limit $\Gamma_{\bar{g}}(r,q)\searrow 0$, $d\Gamma_{\bar{g}}(r,q)\neq 0$ (resp. $\Gamma_{\bar{g}}(p,r)\searrow 0$, $d\Gamma_{\bar{g}}(p,r)\neq 0$). Hence, we have

$$\bar{\kappa}_+=\frac{1}{2(\Gamma_{\bar{g}}(p,r)+\bar{g}^{-1}(d\Gamma_{\bar{g}}(p,r),d\Gamma_{\bar{g}}(r,q)))}\cdot$$

$$\cdot\Bigg[\Gamma_{\bar{g}}(p,r)\left(3-\frac{\Gamma_{\bar{g}}(p,r)}{\Gamma_{\bar{g}}(p,r)+\bar{g}^{-1}(d\Gamma_{\bar{g}}(p,r),d\Gamma_{\bar{g}}(r,q))}\right)+$$

$$-\frac{\Gamma_{\bar{g}}(p,r)\bar{\nabla}^a\Gamma_{\bar{g}}(r,q)\bar{\nabla}^b\Gamma_{\bar{g}}(r,q)\bar{\nabla}_a\bar{\nabla}_b\Gamma_{\bar{g}}(p,r)}{\Gamma_{\bar{g}}(p,r)+\bar{g}^{-1}(d\Gamma_{\bar{g}}(p,r),d\Gamma_{\bar{g}}(r,q))}\Bigg]$$

and

$$\bar{\kappa}_-=-\frac{1}{2(\Gamma_{\bar{g}}(r,q)+\bar{g}^{-1}(d\Gamma_{\bar{g}}(p,r),d\Gamma_{\bar{g}}(r,q)))}\cdot$$

$$\cdot\Bigg[\Gamma_{\bar{g}}(r,q)\left(3-\frac{\Gamma_{\bar{g}}(r,q)}{\Gamma_{\bar{g}}(r,q)+\bar{g}^{-1}(d\Gamma_{\bar{g}}(p,r),d\Gamma_{\bar{g}}(r,q))}\right)+$$

$$-\frac{\Gamma_{\bar{g}}(r,q)\bar{\nabla}^a\Gamma_{\bar{g}}(p,r)\bar{\nabla}^b\Gamma_{\bar{g}}(p,r)\bar{\nabla}_a\bar{\nabla}_b\Gamma_{\bar{g}}(r,q)}{\Gamma_{\bar{g}}(r,q)+\bar{g}^{-1}(d\Gamma_{\bar{g}}(p,r),d\Gamma_{\bar{g}}(r,q))}\Bigg].$$

The divergence of T^a, on its turn, leads to

$$\bar{\nabla}_aT^a=\frac{1}{\Gamma_{\bar{g}}(p,r)+\Gamma_{\bar{g}}(r,q)+\bar{g}^{-1}(d\Gamma_{\bar{g}}(p,r),d\Gamma_{\bar{g}}(r,q))}\cdot$$

$$\Bigg[\Gamma_{\bar{g}}(p,r)\bar{\nabla}_a\bar{\nabla}^a\Gamma_{\bar{g}}(r,q)-\Gamma_{\bar{g}}(r,q)\bar{\nabla}_a\bar{\nabla}^a\Gamma_{\bar{g}}(p,r)-\frac{1}{\Gamma_{\bar{g}}(p,r)+\Gamma_{\bar{g}}(r,q)+\bar{g}^{-1}(d\Gamma_{\bar{g}}(p,r),d\Gamma_{\bar{g}}(r,q))}\cdot$$

$$\cdot\Big((\Gamma_{\bar{g}}(p,r)-\Gamma_{\bar{g}}(r,q))\bar{g}^{-1}(d\Gamma_{\bar{g}}(p,r),d\Gamma_{\bar{g}}(r,q))+\Gamma_{\bar{g}}(r,q)\bar{\nabla}^a\Gamma_{\bar{g}}(p,r)\bar{\nabla}^b\Gamma_{\bar{g}}(p,r)\bar{\nabla}_a\bar{\nabla}_b\Gamma_{\bar{g}}(r,q)+$$

$$-\Gamma_{\bar{g}}(p,r)\bar{\nabla}^a\Gamma_{\bar{g}}(r,q)\bar{\nabla}^b\Gamma_{\bar{g}}(r,q)\bar{\nabla}_a\bar{\nabla}_b\Gamma_{\bar{g}}(p,r)\Big)\Bigg],$$

which in the future horizon $\partial_+\mathscr{W}_{p,q}$ becomes

$$\bar{\nabla}_aT^a\restriction_{\partial_+\mathscr{W}_{p,q}}=\frac{1}{\Gamma_{\bar{g}}(p,r)+\bar{g}^{-1}(d\Gamma_{\bar{g}}(p,r),d\Gamma_{\bar{g}}(r,q))}\cdot$$

$$\Bigg[\Gamma_{\bar{g}}(p,r)\bar{\nabla}_a\bar{\nabla}^a\Gamma_{\bar{g}}(r,q)-\frac{1}{\Gamma_{\bar{g}}(p,r)+\bar{g}^{-1}(d\Gamma_{\bar{g}}(p,r),d\Gamma_{\bar{g}}(r,q))}\cdot$$

$$\cdot\Big(\Gamma_{\bar{g}}(p,r)\bar{g}^{-1}(d\Gamma_{\bar{g}}(p,r),d\Gamma_{\bar{g}}(r,q))-\Gamma_{\bar{g}}(p,r)\bar{\nabla}^a\Gamma_{\bar{g}}(r,q)\bar{\nabla}^b\Gamma_{\bar{g}}(r,q)\bar{\nabla}_a\bar{\nabla}_b\Gamma_{\bar{g}}(p,r)\Big)\Bigg],$$

and in the past horizon $\partial_-\mathscr{W}_{p,q}$,

$$\bar{\nabla}_aT^a\restriction_{\partial_-\mathscr{W}_{p,q}}=-\frac{1}{\Gamma_{\bar{g}}(r,q)+\bar{g}^{-1}(d\Gamma_{\bar{g}}(p,r),d\Gamma_{\bar{g}}(r,q))}\cdot$$

$$\left[\Gamma_{\bar{g}}(r,q)\bar{\nabla}_a\bar{\nabla}^a\Gamma_{\bar{g}}(p,r) - \frac{1}{\Gamma_{\bar{g}}(r,q) + \bar{g}^{-1}(d\Gamma_{\bar{g}}(p,r), d\Gamma_{\bar{g}}(r,q))}\cdot\right.$$

$$\left.\cdot\left(\Gamma_{\bar{g}}(r,q)\bar{g}^{-1}(d\Gamma_{\bar{g}}(p,r), d\Gamma_{\bar{g}}(r,q)) - \Gamma_{\bar{g}}(r,q)\bar{\nabla}^a\Gamma_{\bar{g}}(p,r)\bar{\nabla}^b\Gamma_{\bar{g}}(p,r)\bar{\nabla}_a\bar{\nabla}_b\Gamma_{\bar{g}}(r,q)\right)\right].$$

We obtain immediately from the formulae above that

$$\lim_{\to q}\bar{\kappa}_+ = 1 = -\lim_{\to p}\bar{\kappa}_-,\ \lim_{\to q}\bar{\nabla}_a T^a = -d = -\lim_{\to p}\bar{\nabla}_a T^a.$$

The result hence follows from formula (2.48) and the discussion after Definition 2.2. □

REMARK 2.3 *The result and the proof of Theorem 2.6 hold for geodesically convex diamonds $\mathscr{O}_{p,q}$ in arbitrary causally simple spacetimes with no change.*

There are two caveats about Theorem 2.6 as far as the interpretation of the wedge $\mathscr{W}_{p,q}$ as the exterior region of an event horizon is concerned.

- All results of this Subsection involve the metric $\bar{g}$ of the conformal closure, and not the physical metric g. We have, though, that

$$\frac{T^a\bar{\nabla}_a\bar{g}(T,T)}{\bar{g}(T,T)} = \frac{T^a\nabla_a g(T,T)}{g(T,T)} + 2z\langle dz, T\rangle \tag{2.50}$$

and

$$\begin{aligned}\bar{\nabla}_a T^a &= \bar{g}^{ab}\bar{\nabla}_a T_b = \frac{1}{z^2}g^{ab}\bar{\nabla}_a T_b = \\ &= \frac{1}{z^2}\nabla_a T^a - \frac{1}{z^2}g^{ab}C^c_{ab}T_c = \frac{1}{z^2}\nabla_a T^a + \frac{d-2}{2z^3}\langle dz, T\rangle \Rightarrow \\ &\Rightarrow \nabla_a T^a = z^2\bar{\nabla}_a T^a - \frac{d-2}{2z}\langle dz, T\rangle,\end{aligned} \tag{2.51}$$

where C^c_{ab} is the tensor (A.27) which relates the connections $\bar{\nabla}$ and ∇ (see page 118). As T^a is tangent to $\mathscr{I}$ vanishes at p and q (see formula (2.39), page 45), it follows that $\langle dz, T\rangle = o(z)$ as we get close to p or q. Taking into account that T^a is an asymptotic KILLING field, we obtain that $\kappa_\pm$ also defines the asymptotic past/future surface gravity with respect to g as we get close to p, q. As the asymptotic values of $\kappa_\pm$ are the same for all asymptotic KILLING fields $\frac{du^\lambda_{p,q}}{d\lambda}$ in $\mathscr{W}_{p,q}$ as $z \searrow 0$, it follows that the conclusion of Theorem 2.6 also holds for non geodesically convex wedges and the asymptotic KILLING fields constructed in Subsection 2.3.2.

- The sign of κ_+ we've obtained is minus what it's expected from an event horizon, for the origin of our affine parameter for the (future oriented) generators of $\partial_+ \mathscr{W}_{p,q}$ corresponds to q and not to $\partial I^+(p, \overline{\mathscr{M}}) \cap \partial I^-(q, \overline{\mathscr{M}})$, which would correspond to a "bifurcation surface" of the horizon (notice the change of sign in (2.49)). A correction of this problem leads to the desired change of the sign of κ_+.

Let us notice, finally, that there is a hypothesis about the normalization of T^a implicit in the definition of $\kappa_\pm$. More precisely, our definition is conditioned to the following fact: if $\Gamma_{\bar{g}^{(0)}}(p, r) = \Gamma_{\bar{g}^{(0)}}(r, q)$, i.e., r is the middle point of the maximal timelike geodesic in $\mathscr{M}in(r) \ni p, q$ linking p to q, clearly implying that ${}^{(0)}\bar{\nabla}^a \Gamma_{\bar{g}^{(0)}}(p, r) = -{}^{(0)}\bar{\nabla}^a \Gamma_{\bar{g}^{(0)}}(r, q)$, then $\bar{g}^{(0)}(T, T)(r) = -\frac{d_{\bar{g}^{(0)}}(p,q)^2}{16}$. If we rescale T^a by a factor $R \in \mathbb{R}$, $\kappa_\pm$ is rescaled by the same factor, by definition. This has, for instance, the consequence that, if we take $p_0 = (-1, \mathbf{0})$, $q_0 = (1, \mathbf{0})$ and rescale the T^a associated to the diamond $\mathscr{D}_{Rp_0, Rq_0}$ by a factor $\frac{2}{R}$, so as to maintain $\bar{g}^{(0)}(T, T)(r) = \eta(T, T)(0) = -1$, it follows that $\kappa_\pm \overset{R \to +\infty}{\longrightarrow} 0$, in a way consistent with the fact that $\frac{2}{R} T^a \overset{R \to +\infty}{\longrightarrow} (\partial_0)^a$ (physically, the "Unruh temperature" associated to time translations in Minkowski spacetime is zero).

We can understand Proposition 1.8 as a formulation of the second law of black hole dynamics for AAdS wedges, as the "shrinking" of AAdS wedges expressed there provokes a decrease of the area of the transversal sections of the past and future horizons, the larger the deeper we penetrate the bulk, that is, in the opposite orientation to the one of the time evolution (notice that, unlike Hawking's Area Theorem, we cannot express this result directly in terms of the physical spacetime $(\mathscr{M}, g)$, for the transversal sections of horizons of AAdS wedges are noncompact and hence have always infinite area). The case in which there is no increase of "entropy", which characterizes *reversible* processes, becomes:

Theorem 2.7 (Characterization of reversible processes in AAdS wedges) *Let $(\mathscr{M}, g)$ be am asymptotically simple AAdS spacetime satisfying the hypotheses of Proposition 1.8, and $(p, \bar{q}) \in \mathscr{D}(\mathscr{I})$, $r \in \mathscr{D}_{p,\bar{q}}$. Suppose that, for r sufficiently close to q, we have $\mathscr{W}'_{r,\bar{q}} \cap \mathscr{W}'_{q,\bar{r}} = \varnothing$. Then $\partial_+ \mathscr{W}_{s,\bar{q}}$ is a null $\mathscr{C}^\infty$ hypersurface which is totally geodesic with respect to g for all $s \in \mathscr{I}$ such that $(s, \bar{q}) \in \mathscr{D}(\mathscr{I})$. In particular, $\mathscr{W}'_{s,\bar{q}} \cap \mathscr{W}'_{q,\bar{s}} = \varnothing$ for all $s \in \mathscr{I}$ such that $(s, \bar{q}) \in \mathscr{D}(\mathscr{I})$.*

Proof. Follows immediately from Proposition 1.8 and Remark 1.7. □

Notice that, by virtue of Theorem 2.6, it makes no sense to speak of a "third law" for the dynamics of AAdS wedges and diamonds.

— A lo que yo imagino — dijo don Quijote —, no hay historia humana en el mundo que no tenga sus altibajos, especialmente las que tratan de caballerías, las cuales nunca pueden estar llenas de prósperos sucesos.[ix]

Miguel de Cervantes Saavedra
El ingenioso hidalgo don Quijote de la Mancha, Segundo Libro, Cap. III

Part II

Holographic avatars of Local Quantum Physics

Chapter 3

Locally Covariant Quantum Theories

(Hamlet.) — *O God! I could be bounded in a nutshell, and count myself a king of infinite space, were it not that I have bad dreams.*

William Shakespeare
Hamlet, Prince of Denmark, Act II, Scene II

Having convinced ourselves about the need for rethinking the implementation of asymptotic isometries of spacetime already in the context of classical gravity, we shall now introduce a proper formalism for the discussion of this question at the level of local quantum theories. Such a formalism was recently proposed by Brunetti, Fredenhagen and Verch [BFV03], and one can base it on the observation, related to theăremark at the beginning of Chapter 2 (see page 32), that the specification of physical procedures and its relative ordering are inherently *local* – these are insensitive to the structure of the Universe as a whole, into which the localization region of these procedures is embedded. This, as we'll see, generalizes the notions of isotony and covariance of physical procedures in a single blow. We can even raise this observation to the category of *principle* — say, of *local covariance*.

On the other hand, we are immediately forced, in the light of these considerations, to consider *all regions of all possible spacetimes on an equal footing.* More precisely, we must specify the possible physical procedures of a theory for each of these regions, in a coherent manner. Mathematically, we must specify a *functor* from the *category* of "admissible" spacetime regions to the category of possible physical procedures (we advise the reader not familiar with the notions of category and functor to consult Appendix C, page 145). It's now our objective to make more precise the structure we've just sketched, and investigate some of its properties.

3.1 Basic definitions

First, we shall consider a *locally covariant quantum theory* on globally hyperbolic spacetimes, for this class is "dynamically closed" with respect to equations of motion which lead to causal propagation of Cauchy data. In this case, the causal commutation relations between observables can be uniquely determined (the necessary modifications of this concept in non globally hyperbolic cases will be exposed in Chapter 4, page 67).

Definition 3.1 *Let the categories*

$$\mathscr{G}lh_d = \begin{cases} \textit{objects:} & Obj\mathscr{G}lh_d = \textit{d-dimensional, globally hyperbolic spacetimes } (\mathscr{M}, g); \\ \textit{morphisms:} & Hom_{\mathscr{G}lh_d}((\mathscr{M}, g), (\mathscr{M}', g')) = \{\psi : (\mathscr{M}, g) \to (\mathscr{M}', g') \\ & \textit{isometric embeddings with open, causally convex image}\}; \end{cases}$$

and

$$\mathscr{A}lg = \begin{cases} \textit{objects:} & Obj\mathscr{A}lg = \textit{unital } C^*\textit{-algebras } \mathfrak{A}; \\ \textit{morphisms:} & Hom_{\mathscr{A}lg}(\mathfrak{A}, \mathfrak{A}') = \{\alpha : \mathfrak{A} \to \mathfrak{A}' \textit{ unital } ^*\textit{-monomorphisms}\}. \end{cases}$$

A locally covariant quantum theory *is a covariant functor* $\mathfrak{A}$ *from* $\mathscr{G}lh_d$ *to* $\mathscr{A}lg$.

We say that $\mathfrak{A}$ is:

- *locally causal* if, for any pair $(\mathscr{O}_i, g_i), (\mathscr{M}, g) \in Obj\mathscr{G}lh_d$, $\psi_i \in \mathrm{Hom}_{\mathscr{G}lh_d}((\mathscr{O}_i, g_i), (\mathscr{M}, g))$, $i = 1, 2$, such that $\psi_1(\mathscr{O}_1)$ and $\psi_2(\mathscr{O}_2)$ are causally disjoint in $(\mathscr{M}, g)$, then
$$[(\mathfrak{A}\psi_1)(\mathfrak{A}(\mathscr{O}_1, g_1)), (\mathfrak{A}\psi_2)(\mathfrak{A}(\mathscr{O}_2, g_2))] = 0$$
in $\mathfrak{A}(\mathscr{M}, g)$.

- *primitively causal* if, given $(\mathscr{O}, g), (\mathscr{M}, g) \in Obj\mathscr{G}lh_d$, $\psi \in \mathrm{Hom}_{\mathscr{G}lh_d}((\mathscr{O}, g), (\mathscr{M}, g))$, such that $\psi(\mathscr{O})$ contains a Cauchy surface for $(\mathscr{M}, g)$, then $(\mathfrak{A}\psi)(\mathfrak{A}(\mathscr{O}, g)) = \mathfrak{A}(\mathscr{M}, g)$, i.e., $\mathfrak{A}\psi$ is a *C^*-isomorphism*.

- *additive* if, given $(\mathscr{O}_i, g_i) \in Obj\mathscr{G}lh_d$, $\psi_i \in \mathrm{Hom}_{\mathscr{G}lh_d}((\mathscr{O}_i, g_i), (\mathscr{M}, g))$, $i \in I$, such that the collection $\{\psi_i(\mathscr{O}_i)\}$ is an open covering of $(\mathscr{M}, g)$, then
$$\overline{\bigvee_{i \in I} (\mathfrak{A}\psi_i)(\mathfrak{A}(\mathscr{O}_i, g_i))} = \mathfrak{A}(\mathscr{M}, g)$$
, where $\vee_{i \in I}$ denotes the *-algebra generated by the *-algebras indexed by $i \in I$ and the above closure is in the C*-norm of $\mathfrak{A}(\mathscr{M}, g)$.

Given $(\mathscr{M}_i, g_i) \in Obj\mathscr{G}lh$, $i = 1, 2$, such that $\mathscr{M}_1 \subset \mathscr{M}_2$ and $g_1 = g_2{\restriction}_{\mathscr{M}_1}$, we denote, for future convenience, by $i_{\mathscr{M}_1,\mathscr{M}_2}$ the natural inclusion of $\mathscr{M}_1$ into $\mathscr{M}_2$.

The thoughtful reader will notice that we could generalize Definition 3.1 so as to incorporate any external classical bundle-valued field configuration over the objects of $\mathscr{G}lh$, and not only the metric. A particular case would be the dimensional parameters (masses, coupling constants, etc.) of the quantum field theory in question. This possibility will become more evident in Section 3.4 (page 62, where we define locally covariant quantum fields.

3.2 Relation with the usual axioms

We'll now show how one can recover the axiomatic scheme of Haag and Kastler [HK64, Haa96] for theories of local observables on a *fixed* spacetime starting from a locally covariant quantum theory $\mathfrak{A}$.

Consider a *fixed* spacetime $(\mathscr{M}, g) \in Obj\mathscr{G}lh_d$, with (possibly trivial) isometry group $G(\mathscr{M}, g)$. Let

$$\mathscr{K}(\mathscr{M}, g) \doteq \{\mathscr{O} \subset \mathscr{M} \text{ open, causally convex}\}. \tag{3.1}$$

Notice that, by the definition (3.1), $(\mathscr{O}, g{\restriction}_{\mathscr{O}})$ is automatically globally hyperbolic for all $\mathscr{O} \in \mathscr{K}(\mathscr{M}, g)$, by virtue of the global hyperbolicity of $(\mathscr{M}, g)$. Defining

$$\mathfrak{A}_{\mathscr{M}}(\mathscr{O}) \doteq \mathfrak{A}i_{\mathscr{O},\mathscr{M}}(\mathfrak{A}(\mathscr{O}, g{\restriction}_{\mathscr{O}})), \tag{3.2}$$

it follows that

1. The correspondence $\mathscr{K}(\mathscr{M}, g) \ni \mathscr{O} \mapsto \mathfrak{A}_{\mathscr{M}}(\mathscr{O})$ is *isotonous*, i.e., given $\mathscr{O}_1, \mathscr{O}_2 \in \mathscr{K}(\mathscr{M}, g)$ such that $\mathscr{O}_1 \subset \mathscr{O}_2$, we have $\mathfrak{A}_{\mathscr{M}}(\mathscr{O}_1) \subset \mathfrak{A}_{\mathscr{M}}(\mathscr{O}_2)$.

   To show this, notice that

$$\begin{aligned} \mathfrak{A}i_{\mathscr{O}_1,\mathscr{O}_2}(\mathfrak{A}(\mathscr{O}_1, g{\restriction}_{\mathscr{O}_1})) \subset \mathfrak{A}(\mathscr{O}_2, g{\restriction}_{\mathscr{O}_2}) \quad &\Rightarrow \\ \Rightarrow \mathfrak{A}i_{\mathscr{O}_2,\mathscr{M}}(\mathfrak{A}i_{\mathscr{O}_1,\mathscr{O}_2}(\mathfrak{A}(\mathscr{O}_1, g{\restriction}_{\mathscr{O}_1}))) \quad &= \quad \mathfrak{A}i_{\mathscr{O}_1,\mathscr{M}}(\mathfrak{A}(\mathscr{O}_1, g{\restriction}_{\mathscr{O}_1})) \subset \\ &\subset \quad \mathfrak{A}i_{\mathscr{O}_2,\mathscr{M}}(\mathfrak{A}(\mathscr{O}_2, g{\restriction}_{\mathscr{O}_2})), \end{aligned}$$

   where the equality follows from the covariance of $\mathfrak{A}$.

   In other words, $\mathscr{K}(\mathscr{M}, g) \ni \mathscr{O} \mapsto \mathfrak{A}_{\mathscr{M}}(\mathscr{O})$ is a *precosheaf* of C*-algebras over the causally convex open sets of $\mathscr{M}$, ordered by inclusion (see page 148 in Appendix C for the categorical definition of the concept). If $\mathscr{K}(\mathscr{M}, g)$ is *directed* (i.e., given $\mathscr{O}_1, \mathscr{O}_2 \in \mathscr{K}(\mathscr{M}, g)$, there exists $\mathscr{O}_3 \in \mathscr{K}(\mathscr{M}, g)$ such that $\mathscr{O}_1, \mathscr{O}_2 \subset \mathscr{O}_3$), then $\mathscr{K}(\mathscr{M}, g) \ni \mathscr{O} \mapsto \mathfrak{A}_{\mathscr{M}}(\mathscr{O})$ is a *net*.

2. There exists a representation of $G(\mathscr{M},g) \ni u$ by *-automorphisms $\alpha_u \doteq \mathfrak{A}u$ of $\mathfrak{A}(\mathscr{M},g)$ coherent with the precosheaf (net) structure above, that is, $\alpha_{u_1} \circ \alpha_{u_2} = \alpha_{u_1 u_2}$ and $\alpha_u(\mathfrak{A}_{\mathscr{M}}(\mathscr{O})) = \mathfrak{A}_{\mathscr{M}}(u(\mathscr{O}))$, $\forall u, u_1, u_2 \in G(\mathscr{M},g)$, $\mathscr{O} \in \mathscr{K}(\mathscr{M},g)$.

   The first property follows immediately from the covariance of $\mathfrak{A}$. To show the second property, notice that $u \restriction_{\mathscr{O}}$ is a isometric embedding of $(\mathscr{O}, g \restriction_{\mathscr{O}})$ into $(u(\mathscr{O}), g\restriction_{u(\mathscr{O})})$, and that $i_{\mathscr{O},\mathscr{M}} = i_{u(\mathscr{O}),\mathscr{M}} \circ u$, for all $u \in G(\mathscr{M},g)$, $\mathscr{O} \in \mathscr{K}(\mathscr{M},g)$. The result then follows, once more, from the covariance of $\mathfrak{A}$, as in item 1.

3. If $\mathfrak{A}$ is *locally causal*, it follows that the precosheaf (net) $\mathscr{O} \mapsto \mathfrak{A}_{\mathscr{M}}(\mathscr{O})$ is locally causal in the usual sense of Haag and Kastler, i.e., given $\mathscr{O}_1, \mathscr{O}_2 \in \mathscr{K}(\mathscr{M},g)$ such that $\mathscr{O}_1 \perp_{\mathscr{M}} \mathscr{O}_2$, then $[\mathfrak{A}_{\mathscr{M}}(\mathscr{O}_1), \mathfrak{A}_{\mathscr{M}}(\mathscr{O}_2)] = \{0\}$ (follows immediately from Definition 3.1).

4. Analogously to item 3, if $\mathfrak{A}$ is *primitively causal* (resp. *additive*), the precosheaf (net) $\mathscr{O} \mapsto \mathfrak{A}_{\mathscr{M}}(\mathscr{O})$ is primitively causal (resp. additive) in the usual sense of Haag and Kastler (up to weak closure of the local algebras, which we take for granted – see the end of this Chapter for more on this issue).

Definition 3.2 *Let $\mathfrak{A}$ a locally covariant quantum theory, and $(\mathscr{M},g) \in Obj\mathscr{G}lh$. The precosheaf $\mathscr{K}(\mathscr{M},g) \ni \mathscr{O} \mapsto \mathfrak{A}_{\mathscr{M}}(\mathscr{O})$ (net, if $\mathscr{K}(\mathscr{M},g)$ is directed) is said to be a* realization of $\mathfrak{A}$ in $(\mathscr{M},g)$.

There is a detail on the realization of $\mathfrak{A}$ in $(\mathscr{M},g)$ which requires a certain care with respect to the concept of a Haag-Kastler precosheaf (net) of local observables. It concerns the definition of *quasilocal algebra:* the latter is given by the inductive limit (see Appendix C, formula C.1, page 150 for the categorical definition of the concept)

$$\mathfrak{A}_{\mathscr{M}} \doteq \overline{\bigcup_{\mathscr{O} \in \mathscr{K}(\mathscr{M},g)} \mathfrak{A}(\mathscr{O})^{\|\cdot\|}}, \tag{3.3}$$

where $\|.\|$ is the norm of $\mathfrak{A}(\mathscr{M},g)$. Although clearly $\mathfrak{A}_{\mathscr{M}} \subset \mathfrak{A}(\mathscr{M},g)$, *it's not necessarily true that* $\mathfrak{A}_{\mathscr{M}} = \mathfrak{A}(\mathscr{M},g)$! Nevertheless, is we ignore $\mathfrak{A}(\mathscr{M},g)$ and understand $\mathfrak{A}(\mathscr{O})$ as local C*-subalgebras of $\mathfrak{A}_{\mathscr{M}}$, then we see that the realization of $\mathfrak{A}$ in $(\mathscr{M},g)$ indeed defines a Haag-Kastler precosheaf. A generalization of the concept of quasilocal algebra that allows one to define locally covariant quantum theories in non globally hyperbolic spacetimes and incorporate the specification of boundary conditions will be presented in Chapter 4, page 67.

We'll now investigate some dynamical properties of a *primitively causal* $\mathfrak{A}$. Consider a foliation of $(\mathscr{M},g)$ by Cauchy surfaces $\mathbb{R} \ni t \mapsto \Sigma_t = \tau^{-1}(t)$, given by the global time function τ – more precisely, such foliation is given by the diffeomorphism $F^\tau : \mathbb{R} \times \Sigma \ni (t,x) \mapsto F^\tau(t,x) \in \Sigma_t \subset \mathscr{M}$, where we make the convention $\Sigma \doteq \Sigma_0$ (i.e., $F^\tau(t,\Sigma) = \Sigma_t$

and $F^\tau(\mathbb{R}, x)$ is the orbit of $x \in \Sigma_0$ induced by the foliation). Attribute to each Σ_t an open, causally convex neigbourhood $\mathscr{N}_t \supset \Sigma_t$,[1] and define the two-parameter family of *-isomorphisms [BF07, Ver96]

$$\mathbb{R}^2 \ni (t_1, t_2) \mapsto \alpha^\tau_{t_1,t_2} \doteq (\mathfrak{A}i_{\mathscr{N}_{t_2},\mathscr{M}})^{-1} \circ \mathfrak{A}i_{\mathscr{N}_{t_1},\mathscr{M}} : \mathfrak{A}(\mathscr{N}_{t_1}, g\restriction_{\mathscr{N}_{t_1}}) \to \mathfrak{A}(\mathscr{N}_{t_2}, g\restriction_{\mathscr{N}_{t_2}}), \tag{3.4}$$

denominated *propagator* of $\mathfrak{A}$ in $(\mathscr{M}, g)$ associated to τ. The propagator $\alpha^\tau_{\cdot,\cdot}$ satisfies

$$\alpha^\tau_{t_2,t_3} \circ \alpha^\tau_{t_1,t_2} = \alpha^\tau_{t_1,t_3},\ \alpha^\tau_{t,t} = \mathrm{id}_{\mathfrak{A}(\mathscr{M},g)},\ \forall t_1, t_2, t_3, t \in \mathbb{R}, \tag{3.5}$$

i.e., $\alpha^\tau_{\cdot,\cdot}$ implements the *pair groupoid* $\mathscr{P}\mathbb{R}$ associated to $\mathbb{R}$.[2] It's important to notice that, although the propagator preserves, by definition, the localization of the physical procedures throughout time (i.e., of the orbits of the foliation), it doesn't implement in a *geometrical* way the flux of diffeomorphisms $F(. + t, .) \circ F^{-1}$, $t \in \mathbb{R}$: consider, for instance, an open set $\mathscr{S}$ of Σ, and define $\mathscr{S}_t \doteq F^\tau(t, \mathscr{S})$. It's *false* that $\alpha^\tau_{t_1,t_2}(\mathfrak{A}_{\mathscr{N}_{t_1}}(D(\mathscr{S}_{t_1}) \cap \mathscr{N}_{t_1}, g\restriction_{D(\mathscr{S}_{t_1})\cap\mathscr{N}_{t_1}})) = \mathfrak{A}_{\mathscr{N}_{t_2}}(D(\mathscr{S}_{t_2}) \cap \mathscr{N}_{t_2}, g\restriction_{D(\mathscr{S}_{t_2})\cap\mathscr{N}_{t_2}})$ ($D(.)$ denotes the Cauchy development; notice that the intersection of two causally convex open sets is causally convex), showing that $\alpha^\tau_{\cdot,\cdot}$ is, indeed, a *dynamical* and not a kinematic object.

We also emphasize that the size of $\mathscr{N}_t$ is essentially irrelevant. Indeed, we can take the set $\mathfrak{N}_t$ of all open, causally convex neighbourhoods of Σ_t and define the projective limits (see Appendix C, formula C.1, page 150 for the categorical definition of the concept)

$$\begin{aligned} \mathfrak{A}_{\Sigma_t} &\doteq \bigcup_{\mathscr{N}_t \in \mathfrak{N}_t} \mathfrak{A}(\mathscr{N}_t, g\restriction_{\mathscr{N}_t})/\sim, \\ \mathfrak{A}_{\Sigma_t}(\mathscr{S}_t) &\doteq \bigcup_{\mathscr{N}_t \in \mathfrak{N}_t} \mathfrak{A}_{\mathscr{N}_t}(D(\mathscr{S}_t) \cap \mathscr{N}_t, g\restriction_{D(\mathscr{S}_t\cap\mathscr{N}_t)})/\sim',\ \mathscr{S} \subset \Sigma \text{ open}, \end{aligned} \tag{3.6}$$

where the equivalence relation $\sim$ identifies the elements of the orbits of the family of *-isomorphisms $\mathfrak{A}i_{\mathscr{N}_t^1,\mathscr{N}_t^2}$ and the equivalence relation $\sim'$ identifies the elements of

[1] Notice that we haven't first defined $\mathscr{N}_{t_0}$ for a fixed t_0 and later on defined $\mathscr{N}_t$ as the image of $\mathscr{N}_{t_0}$ under the flux from t_0 to t induced by the foliation. As this family of diffeomorphisms doesn't necessarily preserve the causal structure of $\mathscr{N}_{t_0}$ – for such, it's necessary and sufficient that the diffeomorphisms here are *conformal transformations* [BEE96] –, the image of $\mathscr{N}_{t_0}$ may not be causally convex. Moreover, it's sometimes convenient to treat two *disjoint* Cauchy surfaces without necessarily having the flux of a foliation connecting both; a situation when this occurs will be illustrated later on.

[2] Given a set S, the *pair groupoid* $\mathscr{P}S$ associated to S is defined in the following way (for the definition of groupoid, see Appendix C, page 147): $Obj\mathscr{P}S \doteq S$, $Arr\mathscr{P}S = S^2 \ni (s_1, s_2)$, where $D((s_1, s_2)) = s_1$, $CD((s_1, s_2)) = s_2$, $\mathbb{1}_s = (s, s)$ and the composition law is given by $(s_2, s_3)(s_1, s_2) = (s_1, s_3)$.

the orbits of the family of *-isomorphisms $\mathfrak{A}i_{D(\mathscr{S}_t)\cap\mathscr{N}_t^1, D(\mathscr{S}_t)\cap\mathscr{N}_t^2}$, $\mathscr{N}_t^i \in \mathfrak{N}_t$, $i = 1, 2$, $\mathscr{N}_t^1 \subset \mathscr{N}_t^2$. This procedure defines the quasilocal algebra and the local subalgebras of *germs* of physical procedures at instant t with respect to the foliation induced by τ. We also denote by $\alpha^{\tau}_{t_1,t_2}$ the propagator linking $\mathfrak{A}_{\Sigma_{t_1}}$ to $\mathfrak{A}_{\Sigma_{t_2}}$ induced by (3.6). Notice that the projective limit is *not* the "restriction of $\mathfrak{A}(\mathscr{M}, g)$ to Σ_t" (such an identification is possible only for free quantum field theories, for which the restriction to spacelike hypersurfaces is a mathematically well defined operation).

3.3 Quantum states in curved spacetimes

Now, the notion of *states* requires more care in this formalism, for, unlike physical procedures, states *are inherently nonlocal* – they are sensitive to the global geometry of spacetime and can as well respond in a nonlocal manner to local perturbations of the metric.

Definition 3.3 *Let the category*

$$\mathscr{S}ts = \left\{ \begin{array}{rl} \textit{objects:} & \textit{Obj}\mathscr{S}ts = \textit{sets } \mathfrak{S} \textit{ of states over unital} \\ & \textit{C*-algebras;} \\ \textit{morphisms:} & \textit{Hom}_{\mathscr{S}ts}(\mathfrak{S}, \mathfrak{S}') = \{\sigma : \mathfrak{S} \to \mathfrak{S}' \textit{ linear positive maps;}\}. \end{array} \right.$$

Given a locally covariant quantum theory $\mathfrak{A}$*, a* (locally contravariant) state space *is a contravariant functor* $\mathfrak{S}$ *from* $\mathscr{G}lh_d$ *to* $\mathscr{S}ts$*, such that* $\mathfrak{S}\psi = (\mathfrak{A}\psi)^*$ *is the* dual (pull-back) *of* $\mathfrak{A}\psi$ *for all* $\psi \in Arr\mathscr{G}lh$*, i.e., given* $\omega \in \mathfrak{S}(\mathscr{M}', g')$ *and* $\psi \in \textit{Hom}_{\mathscr{G}lh}((\mathscr{M}, g), (\mathscr{M}', g'))$*, we have* $\mathfrak{S}\psi \in \textit{Hom}_{\mathscr{S}ts}(\mathfrak{S}(\mathscr{M}', g'), \mathfrak{S}(\mathscr{M}, g))$ *is defined by* $\mathfrak{S}\psi(\omega) = (\mathfrak{A}\psi)^*(\omega) \doteq \omega \circ \mathfrak{A}\psi$*.*

3.4 Locally covariant quantum fields

In the developments presented so far, the topological structure of C*-algebras attributed to the objects of $\mathscr{A}lg$ have had only a secondary role (definition of quasilocal algebras, etc.). Indeed, we could have defined the objects of $\mathscr{A}lg$ as unital *-algebras, and attributed a topological structure suited to convenience. Before proceeding further, let us define the category

$$\mathscr{T}alg = \left\{ \begin{array}{rl} \text{objects:} & \textit{Obj}\mathscr{T}alg = \text{unital topological *-algebras } \mathfrak{F}; \\ \text{morphisms:} & \text{Hom}_{\mathscr{T}alg}(\mathfrak{F}, \mathfrak{F}') = \{\alpha : \mathfrak{F} \to \mathfrak{F}' \text{ unital *-monomorphisms}\}. \end{array} \right.$$

and the pushforward functor of test functions $\mathfrak{D} : \mathscr{G}lh_d \to \mathscr{T}vs$ given by

$$\begin{aligned} \mathfrak{D}(\mathscr{M}, g) &= \mathscr{C}_c^{\infty}(\mathscr{M}), \\ \mathfrak{D}\psi(f)(p) &= \psi_*(f)(p) \doteq \left\{ \begin{array}{ll} f(q) & (p = \psi(q)) \\ 0 & \text{otherwise} \end{array} \right. , \end{aligned} \tag{3.7}$$

where $\mathscr{T}vs$ is the category of topological vector spaces (see Appendix C). We can also, sometimes, understand $\mathfrak{D}$ as a functor from $\mathscr{G}lh$ to the category $\mathscr{T}op$ of topological spaces.

DEFINITION 3.4 *A* locally covariant quantum field theory *is a covariant functor* $\mathfrak{F} : \mathscr{G}lh \to \mathscr{T}alg$. *A* locally covariant scalar quantum field *is a natural transformation* $\Phi : \mathfrak{D} \to \mathfrak{F}$, *where* $\mathfrak{D}$ *and* $\mathfrak{F}$ *are here understood as functors from* $\mathscr{G}lh$ *to* $\mathscr{T}op$. *We say that* Φ *is* linear *if* $\Phi_{(\mathscr{M},g)} \in Arr\mathscr{T}vs$ *for all* $(\mathscr{M}, g) \in Obj\mathscr{G}lh$.

REMARK 3.5 *Notice that the hypothesis of* $\Phi_{(\mathscr{M},g)}$ *being linear in the sense of Definition 3.4 has* nothing *to do with the nature of its self interaction!*

The notions of *local causality, primitive causality, additivity, realization* of $\mathfrak{F}$ in $(\mathscr{M}, g) \in Obj\mathscr{G}lh$ and of *state space* are naturally obtained as extensions of the corresponding concepts for locally covariant quantum theories (the only needed modification is the substitution of "*-algebra" for "C*-algebra" in formula (3.7) of Definition 3.3, page 62).

A typical example (albeit apparently trivial) of a locally covariant quantum field theory is the attribution, to each $(\mathscr{M}, g) \in Obj\mathscr{G}lh$, of the BORCHERS-UHLMANN *algebra* $\mathscr{F}(\mathscr{M})$ (see Appendix B.3, Definition B.11, page 143) – it's immediate to see that $\mathfrak{F}$ satisfies Definition 3.4. This locally covariant quantum field theory serves uniquely the purpose of defining the kinematics of an Hermitian scalar field, and it's onwards called the BORCHERS-UHLMANN *functor*. The realization of this $\mathfrak{F}$ in $(\mathscr{M}, g) \in Obj\mathscr{G}lh$ has as its quasilocal algebra $\mathfrak{F}_{\mathscr{M}} = \mathfrak{F}(\mathscr{M})$ (it suffices to apply partitions of unity), and possible examples of locally covariant scalar quantum fields associated to $\mathfrak{F}$ are:

- $\Phi_{(\mathscr{M},g)} = \mathrm{id}_{\mathscr{C}_c^\infty(\mathscr{M})}$, i.e, Φ identifies $f \in \mathscr{C}_c^\infty(\mathscr{M})$ with the corresponding element of $\mathfrak{F}(\mathscr{M})$;
- $\Phi_{(\mathscr{M},g)}(f) = [f]$, where $[f]$ is the equivalence class of f under the quotient of $\mathfrak{F}(\mathscr{M})$ modulo a *-ideal $\mathfrak{I}(\mathscr{M}, g)$. In order to $(\mathscr{M}, g) \mapsto \mathfrak{F}(\mathscr{M})/\mathfrak{I}(\mathscr{M}, g)$ to define a locally covariant quantum field theory, it's necessary that the correspondence $(\mathscr{M}, g) \mapsto \mathfrak{I}(\mathscr{M}, g)$ defines a covariant functor from $\mathscr{G}lh$ to $\mathscr{T}alg$. In this case, the quotient map defines a natural transformation, by virtue of the its universal property [Jac85] (see also Appendix C, page 143). The specification of *-ideals as $\mathfrak{I}(\mathscr{M}, g)$ constitutes a way to impose a dynamics to the BORCHERS-UHLMANN algebra. This example, as well as the previous one, result in linear fields.
- Here we elaborate the previous example, basing on the considerations at the end of Section B.3. Consider a state space $\mathfrak{S}$, where $\mathfrak{S}(\mathscr{M}, g)$ consists of a set of

hierarchies ω of k-point distributions $\omega_k \in \mathscr{D}'(\mathscr{M}^k)$ and $\mathfrak{S}\psi$ is the pullback ψ^* of distributions under $\psi \in Arr\mathscr{G}lh$, such that

$$\mathfrak{I}(\mathscr{M},g) \subset \bigcap_{\omega\in\mathfrak{S}(\mathscr{M},g)} \mathrm{Ann}\omega.$$

In this case, the locally covariant quantum field Φ of the former example naturally induces a set of states $\Phi^*_{(\mathscr{M},g)}\mathfrak{S}(\mathscr{M},g)$ over each $\widetilde{\mathfrak{F}}(\mathscr{M},g) \doteq \mathfrak{F}(\mathscr{M})/\mathfrak{I}(\mathscr{M},g)$

$$(\Phi^*_{(\mathscr{M},g)}\omega_k)([f_1],\ldots,[f_k]) = \omega_k(f_1,\ldots,f_k),$$

in a way that $\Phi^*_{\cdot}\mathfrak{S}(.)$ is a locally contravariant state space.

The possibility of proceeding as in the formalism of Wightman and defining quantum fields starting from states by means of the Wightman-GNS representation can, of course, can, of course, be conveyed for the realization of the Borchers-Uhlmann functor in some $(\mathscr{M},g) \in \mathscr{G}lh$, although it should be considered with care as far as the Principle of Local Covariance is concerned, for *individual* states are not locally contravariant. We may, nevertheless, consider the following scenario – by the way, already assessed by Haag and Kastler in [HK64], and which served them as a motivation for the introduction of the algebraic approach to QFT – under the light of the examples addressed above.

Let $\mathfrak{S}$ be a state space associated to $\mathfrak{F}$, such that

(3.8)
$$\mathrm{Ann}\omega_{(\mathscr{M},g)} = \mathrm{Ann}\omega'_{(\mathscr{M},g)} \doteq \mathfrak{I}(\mathscr{M},g),\ \forall\omega_{(\mathscr{M},g)},\omega'_{(\mathscr{M},g)} \in \mathfrak{S}(\mathscr{M},g),\ (\mathscr{M},g) \in Obj\mathscr{G}lh.$$

Employing the contravariance of $\mathfrak{S}$, we can show that, in this case, the correspondence $(\mathscr{M},g) \mapsto \mathfrak{I}(\mathscr{M},g)$ defines a locally covariant quantum field theory. Hence, as we've seen, $\Phi_{(\mathscr{M},g)}(f) = [f] = f/\mathfrak{I}(\mathscr{M},g)$ defines then a locally covariant scalar quantum field, thus realizing our *desideratum* about incorporating the dynamical information of a locally covariant quantum field theory into the specification of a state space, showing at the same time that the quantum field obtained is independent of a choice of a representative from the set of states, by the First *-Isomorphism Theorem.

From the viewpoint of a locally covariant quantum field theory $\mathfrak{A}$, (3.8) has an even deeper interpretation: the GNS representations of $\mathfrak{A}(\mathscr{M},g)$ associated to each $\omega_{(\mathscr{M},g)} \in \mathfrak{S}(\mathscr{M},g)$ are not only *quasiequivalent* (i.e., the sets of normal states over $\pi_{\omega_{(\mathscr{M},g)}}(\mathfrak{A}(\mathscr{M},g))''$ and $\pi_{\omega'_{(\mathscr{M},g)}}(\mathfrak{A}(\mathscr{M},g))''$ coincide, or, equivalently, these two von Neumann algebras are *-isomorphic – see Theorem 2.4.26 of [BR87]), but also, as a consequence, the GNS representations of the quasilocal algebra $\mathfrak{A}_{\mathscr{M}}$ of the realization of $\mathfrak{A}$ in any $(\mathscr{M},g)$ are *locally quasiequivalent*, that is, the sets of normal states over

$\pi_{\omega_{(\mathscr{M},g)}}(\mathfrak{A}(\mathscr{O}))''$ and $\pi_{\omega'_{(\mathscr{M},g)}}(\mathfrak{A}(\mathscr{O}))''$ coincide. Hence, local quasiequivalence becomes a highly desirable requirement for physically relevant state spaces and essentially allows us to pass from local C*-algebras to local VON NEUMANN algebras in the same manner as in the HAAG-KASTLER formalism. An even more detailed discussion about the role of local quasiequivalence in the context of the Principle of Local Covariance can be found in [BFV03].

Chapter 4

Implementing the Holographic Principle in AAdS spacetimes

In due modi si raggiunge Despina: per nave o per cammello. La città si presenta differente a chi viene da terra e a chi dal mare.

(...)

Ogni città riceve la sua forma dal deserto a cui si oppone; e così il cammelliere e il marinaio vedono Despina, città di confine tra due deserti.[x]

Italo Calvino
"Le città e il desiderio 3" (*Le città invisibili*)[xi]

4.1 Local covariance with boundary conditions

In Chapter 3, we've built a general framework for the description of locally covariant quantum theories in globally hyperbolic spacetimes. Within such a class, commutation relations among local procedures (given by the elements of the local algebras) can be established without ambiguities, for globally hyperbolic spacetimes constitute "closed" systems with respect to *causal* dynamical laws, i.e., which propagate local physical effects with speed inferior or equal of that of the light. However, we know that we can always cover *any* spacetime with open, globally hyperbolic and causally convex neighbourhoods – in the case of strongly causal spacetimes, such neighbourhoods even constitute a basis for the spacetime topology. And not even for this should we expect that, in these "open" systems, we may assume local causality *a priori*.

In the case of the realization of $\mathfrak{A}$ in a globally hyperbolic spacetime, local causality is coherent with the inclusions of the algebras, for by means of an adequate choice of the family of globally hyperbolic neighbourhoods, it's always possible to find, given any two causally disjoint such neighbourhoods (one of them, eventually, being sufficiently small), a third one which contains both.[1] It's precisely the violation of this property that causes difficulties in the case of non globally hyperbolic spacetimes.

To pinpoint such difficulties proper in the extension of the Principle of Local Covariance to spacetimes such as (A)AdS, we shall now exhibit a more general way to construct the realization of a locally covariant quantum theory $\mathfrak{A}$, valid even for non globally hyperbolic spacetimes $(\mathscr{M}, g)$. This path was proposed by Sommer [Som06] and makes more transparent the necessity and the role of boundary conditions on the imposition of local and primitive causality – in order to make clearer the context of the latter property, we'll restrict ourselves to stably causal spacetimes, for these, as seen in Appendix A, can be foliated by "constant time" hypersurfaces, for which it's meaningful to formulate a (mixed boundary-)initial-value problem.

Let, then, $(\mathscr{M}, g)$ be a stably causal spacetime, and $\mathscr{K}(\mathscr{M}, g)$ the set of globally hyperbolic, causally convex open regions $(\mathscr{O}, g\restriction_{\mathscr{O}})$, $\mathscr{O} \subset \mathscr{M}$ (we sometimes omit $g\restriction_{\mathscr{O}}$ whenever such procedure doesn't cause confusion). Let us consider initially, given a locally covariant quantum theory $\mathfrak{A}$,

$$\mathfrak{A}_{\mathscr{M}} = \bigvee \{(A, \mathscr{O}) : A \in \mathfrak{A}(\mathscr{O}, g\restriction_{\mathscr{O}}),\ \mathscr{O} \in \mathscr{K}(\mathscr{M}, g)\}, \tag{4.1}$$

the *free* *-algebra generated by the elements indicated above. Let now the *-ideal $\mathfrak{I}_{\mathscr{M}} \doteq \mathfrak{A}_{\mathscr{M}} \mathfrak{E}_{\mathscr{M}} \mathfrak{A}_{\mathscr{M}}$, where $\mathfrak{E}_{\mathscr{M}}$ is the *-algebra generated by the relations

$$\begin{gathered} (A, \mathscr{O}) + (B, \mathscr{O}) - (A + B, \mathscr{O}), \\ (A, \mathscr{O})(B, \mathscr{O}) - (AB, \mathscr{O}), \\ \lambda(A, \mathscr{O}) - (\lambda A, \mathscr{O}),\ \forall \lambda \in \mathbb{C} \text{ and} \\ (A, \mathscr{O}_1) - (\mathfrak{A}i_{\mathscr{O}_1, \mathscr{O}_2}(A), \mathscr{O}_2),\ \forall \mathscr{O},\ \mathscr{O}_1,\ \mathscr{O}_2 \in \mathscr{K}(\mathscr{M}, g),\ \mathscr{O}_1 \subset \mathscr{O}_2. \end{gathered} \tag{4.2}$$

We know that each element $(A, \mathscr{O})$ possesses the (C*-)norm $\|(A, \mathscr{O})\| = \|A\|$, consistent with the relations (4.2) (recall that $\mathfrak{A}i_{\mathscr{O}_1, \mathscr{O}_2}$ is a **-monomorphism* for $\mathscr{O}_1, \mathscr{O}_2 \in \mathscr{K}(\mathscr{M}, g)$, and, hence, $\|(\mathfrak{A}i_{\mathscr{O}_1, \mathscr{O}_2}(A), \mathscr{O}_2)\| = \|(A, \mathscr{O}_1)\| = \|A\|$!). Applying now the universal construction of Blackadar [Bla85, Bla06],[2] to the generators in (4.1) and to the relations (4.2), we define the collection $\Pi(\mathfrak{A}_{\mathscr{M}}, \mathfrak{I}_{\mathscr{M}})$ of *admissible *-representations* of the pair $(\mathfrak{A}_{\mathscr{M}}, \mathfrak{I}_{\mathscr{M}})$, consisting in the *-representations π of $\mathfrak{A}_{\mathscr{M}}$ such that $\pi(\mathfrak{I}_{\mathscr{M}}) = \{0\}$.

[1] See [GLRV01]. This fact if of the utmost importance in the algebraic theory of superselection sectors.

[2] I thank Prof. Severino Toscano do Rêgo Melo for bringing the reference [Bla85] to my attention.

It thus follows that the *universal* C*-seminorm $\|A\|_u = \sup_{\pi\in\Pi(\mathfrak{A}_{\mathscr{M}},\mathfrak{I}_{\mathscr{M}})}\|\pi(A)\|$ is *finite* for all A. Denoting the *-ideal $\bar{\mathfrak{I}}_{\mathscr{M}} \doteq \{A : \|A\|_u = 0\}$, then we obtain the *quasilocal* C*-algebra $\mathfrak{A}(\mathscr{M}, g)$ as the universal C*-algebra given by the generators $\mathfrak{A}_{\mathscr{M}}$ and relations $\mathfrak{I}_{\mathscr{M}}$:[3]

$$\mathfrak{A}(\mathscr{M}, g) \doteq \overline{(\mathfrak{A}_{\mathscr{M}}/\bar{\mathfrak{I}}_{\mathscr{M}})}^{\|.\|_u}. \tag{4.3}$$

Let's now see how the *-morphisms induced by isometric embeddings $\psi : (\mathscr{M}_1, g_1) \to (\mathscr{M}_2, g_2)$ with causally convex image behave. Obviously, $\psi(\mathscr{K}(\mathscr{M}_1, g_1)) \subset \mathscr{K}(\mathscr{M}_2, g_2)$, and $\psi\restriction_{\mathscr{O}}$, $\mathscr{O} \in \mathscr{K}(\mathscr{M}_1, g_1)$, defines and isometric embedding of $\mathscr{O}$ into $\psi(\mathscr{O})$, naturally defining a *-monomorphism from $\mathfrak{A}_{\mathscr{M}_1}$ to $\mathfrak{A}_{\mathscr{M}_2}$. Let us define the following *-morphism:

$$\begin{aligned} \mathfrak{A}\psi : \mathfrak{A}(\mathscr{M}_1, g_1) &\to \mathfrak{A}(\mathscr{M}_2, g_2) \\ A + \bar{\mathfrak{I}}_{\mathscr{M}_1} &\mapsto \mathfrak{A}\psi(A) + \bar{\mathfrak{I}}_{\mathscr{M}_2}. \end{aligned} \tag{4.4}$$

Notice that, due to the considerations of the previous paragraph, if $[A]_{\mathscr{M}_i}$ is the equivalence class of $A \in \mathfrak{A}_{\mathscr{M}_i}$ (under the quotient) modulo $\bar{\mathfrak{I}}_{\mathscr{M}_i}$ (i.e., $[A]_{\mathscr{M}_i} = [A+J]_{\mathscr{M}_i}$ if $J \in \bar{\mathfrak{I}}_{\mathscr{M}_i}$), $i = 1, 2$, it follows that $\mathfrak{A}\psi([A + J]_{\mathscr{M}_1}) = [\mathfrak{A}\psi \restriction_{\mathscr{O}}(A)]_{\mathscr{M}_2}$, for all $A \in \mathfrak{A}(\mathscr{O}, g\restriction_{\mathscr{O}})$, showing that the definition (4.4) isn't empty and indeed defines a nontrivial *-morphism. $\mathfrak{A}\psi$ is unital, but *not necessarily injective!* Hence, we must weaken the properties of the morphisms of the category of (C)*-algebras so as to $\mathfrak{A}$ to define a covariant functor from the (full super)category of the stably causal spacetimes to the latter – the covariance of $\mathfrak{A}$ is guaranteed by the First *-Isomorphism Theorem (Theorem B.2, page 123). We'll now exhibit a situation which illustrates with perfection how this fact complicates the imposition of local causality.

Consider $\mathscr{O}_1$, $\mathscr{O}_2 \in \mathscr{K}(\mathscr{M}_1, g_1)$ causally disjoint, such that there is *no* $\mathscr{O}_3 \in \mathscr{K}(\mathscr{M}_1, g_1)$ such that $\mathscr{O}_1$, $\mathscr{O}_2 \subset \mathscr{O}_3$, but there exists an $\mathscr{O}_4 \in \mathscr{K}(\mathscr{M}_2, g_2)$ such that $\psi(\mathscr{O}_1)$, $\psi(\mathscr{O}_2) \subset \mathscr{O}_4$. Let $\mathfrak{A}$ be locally causal (i.e., with respect to globally hyperbolic regions). Then we have

$$[(\mathfrak{A}(i_{\psi(\mathscr{O}_1),\mathscr{O}_4} \circ \psi)(A), \mathscr{O}_4), (\mathfrak{A}(i_{\psi(\mathscr{O}_2),\mathscr{O}_4} \circ \psi)(B), \mathscr{O}_4)] = 0.$$

However, there is nothing under our hypotheses preventing

$$[(A, \mathscr{O}_1), (B, \mathscr{O}_2)] = C \neq 0,$$

unless $A = 0$ or $B = 0$. Hence, for $A, B \neq 0$, it follows that $C \notin \mathfrak{I}_{\mathscr{M}_1}$ and $[C]_{\mathscr{M}_1} \in \mathrm{Ker}\mathfrak{A}\psi$, potentially spoiling the injectivity of $\mathfrak{A}\psi$ (it may occur, though, that $C \in \bar{\mathfrak{I}}_{\mathscr{M}_1}$,

[3]Notice that we *haven't* taken the quotient modulo the *-ideal $\mathfrak{I}_{\mathscr{M}}$. Were there a way to guarantee the existence of an *admissible* *-representation π such that $\mathrm{Ker}\pi = \mathfrak{I}_{\mathscr{M}}$, we could equivalently take such a quotient. However, this is in general impossible, for $\mathfrak{I}_{\mathscr{M}}$ is not necessarily closed in the C*-norm defined for the elements of $\mathfrak{A}_{\mathscr{M}}$.

but there is no way of guaranteeing this *a priori*). Nevertheless, the *-morphisms $\mathfrak{A}(\psi \circ i_{\mathscr{O},\mathscr{M}_1})$ are always injective, for all $\mathscr{O} \in \mathscr{K}(\mathscr{M}_1, g_1)$.

The construction above allows one to write the following

Definition 4.1 *Let the categories*

$$\mathscr{S}tc_d = \begin{cases} \textit{objects:} & Obj\mathscr{G}lh_d = \textit{d-dimensional, stably causal spacetimes } (\mathscr{M}, g); \\ \textit{morphisms:} & Hom_{\mathscr{S}tc_d}((\mathscr{M}, g), (\mathscr{M}', g')) = \{\psi : (\mathscr{M}, g) \to (\mathscr{M}', g') \\ & \textit{isometric embeddings with open, causally convex image}\}; \end{cases}$$

($\mathscr{G}lh$ is, hence, a full subcategory of $\mathscr{S}tc$) and

$$\mathscr{A}lgb = \begin{cases} \textit{objects:} & Obj\mathscr{A}lg = \textit{unital } C^*\textit{-algebras } \mathfrak{A}; \\ \textit{morphisms:} & Hom_{\mathscr{A}lg}(\mathfrak{A}, \mathfrak{A}') = \{\alpha : \mathfrak{A} \to \mathfrak{A}' \textit{ unital *-morphisms} \\ & \textit{(not necessarily injective)}\}. \end{cases}$$

An extended locally covariant quantum theory *is a covariant functor $\mathfrak{A}$ from $\mathscr{S}tc_d$ to $\mathscr{A}lgb$, such that $\mathfrak{A}\restriction_{\mathscr{G}lh}$ is a locally covariant quantum theory in the sense of Definition 3.1, page 58. Given $(\mathscr{M}, g) \in Obj\mathscr{S}tc_d$, the* realization *of $\mathfrak{A}$ in $(\mathscr{M}, g)$ is given by the precosheaf $\mathscr{K}(\mathscr{M}, g) \ni \mathscr{O} \mapsto \mathfrak{A}(\mathscr{O}) \doteq \overline{(\{(A, \mathscr{O}) : A \in \mathfrak{A}(\mathscr{O}, g\restriction_{\mathscr{O}})\}/\bar{\mathfrak{I}}_{\mathscr{M}})^{\|\cdot\|}}$, following* Blackadar*'s construction given above. We say yet that $\mathfrak{A}$ is* regular *is each local algebra $\mathfrak{A}(\mathscr{M}, g)$ coincides with the quasilocal algebra of the realization of$\mathfrak{A}$ in $(\mathscr{M}, g)$, given by (4.3).*[4]

It's immediate to see that the realization of $\mathfrak{A}$ in $(\mathscr{M}, g) \in Obj\mathscr{G}lh_d$ in the sense of Definition 4.1 coincides with Definition 3.2 up to an *-isomorphism. The loss of injectivity of the *-morphisms induced by $\mathfrak{A}$ translates itself to the freedom of choice of boundary conditions. Before making this idea more precise, we'll show what becomes of the remaining properties of $\mathfrak{A}$.

- If $(\mathscr{M}, g)$ possesses a nontrivial group of isometries $G \ni \psi$, it becomes clear that ψ induces a bijection of $\mathscr{K}(\mathscr{M}, g) \ni \mathscr{O}$ onto itself. Thus, we see that the *-morphism $\mathfrak{A}\psi$ is explicitly invertible, with inverse given by $\mathfrak{A}(\psi^{-1})$, for all $\psi \in G$. Thus, the Principle of Local Covariance allows the implementation of the action of isometry groups even in the non globally hyperbolic case.

[4]Here, we indulge in a slight abuse of notation, for we'll only deal with the regular case in what follows.

- *Local causality* holds in the following sense: if the restriction of $\mathfrak{A}$ to the category of globally hyperbolic spacetimes is locally causal, and given, as in the example above, $\mathscr{O}_1$, $\mathscr{O}_2 \in \mathscr{K}(\mathscr{M}_1, g_1)$ causally disjoint with respect to $\mathscr{O}_3 \in \mathscr{K}(\mathscr{M}_1, g_1)$ (i.e., $\mathscr{O}_3 \supset \mathscr{O}_1, \mathscr{O}_2$ and $\mathscr{O}_1 \perp_{\mathscr{O}_3} \mathscr{O}_2$), it follows by local covariance that

$$[\mathfrak{A}i_{\mathscr{O}_1,\mathscr{M}}(A_1), \mathfrak{A}i_{\mathscr{O}_2,\mathscr{M}}(A_2)] = 0,$$

for all $A_i \in \mathfrak{A}(\mathscr{O}_i, g\restriction_{\mathscr{O}_i})$, $i = 1, 2$.

- *Additivity* remains valid with no changes, as long as the covering consists in globally hyperbolic regions.

- *Primitive causality* has no immediate extension to non globally hyperbolic spacetimes. In the case of *stably causal* spacetimes, a tentative definition could be the following: let τ be a global time function in $(\mathscr{M}, g)$, and $\mathscr{N} \subset \mathscr{M}$ an open, causally convex neighbourhood of, say, $\Sigma = \tau^{-1}(0)$. Then we say that $\mathfrak{A}$ is *weakly primitively causal* if it's primitively causal with respect to globally hyperbolic regions and $\mathfrak{A}i_{\mathscr{N},\mathscr{M}}$ is a **-epimorphism*. Such a definition is convenient, for the ambiguity in the choice of boundary conditions becomes expressed in a purely algebraic manner, as follows: consider any *-ideal $\mathfrak{J}$ of $\mathfrak{A}(\mathscr{N}, g\restriction_{\mathscr{N}})$ containing $\mathrm{Ker}\mathfrak{A}\psi$. If $\mathfrak{A}$ is weakly primitively causal, then the map $A + \mathfrak{J} \mapsto \mathfrak{A}\psi(A) + \mathfrak{A}i_{\mathscr{N},\mathscr{M}}(\mathfrak{J})$ is a *-isomorphism from $\mathfrak{A}(\mathscr{N}, g\restriction_{\mathscr{N}})/\mathfrak{J}$ onto $\mathfrak{A}(\mathscr{M}, g)/\mathfrak{A}i_{\mathscr{N},\mathscr{M}}(\mathfrak{J})$, by the First *-Isomorphism Theorem.

The specification of the ideal $\mathfrak{J}$ in the context of *weak primitive causality* corresponds precisely to the choice of boundary conditions. That a "minimal" choice is always possible, is guaranteed by the definition of weak primitive causality (it suffices to take $\mathfrak{J} = \mathrm{Ker}\mathfrak{A}i_{\mathscr{N},\mathscr{M}}$). A "maximal" choice would be the case in which $\mathfrak{J}$ is a *maximal* ideal of $\mathfrak{A}(\mathscr{N}, g\restriction_{\mathscr{N}})$ (the existence of maximal *-ideals follows from the Zorn lemma; see Appendix C for its statement in a categorical context) – then the quotient *-algebra $\mathfrak{A}(\mathscr{N}, g\restriction_{\mathscr{N}})/\mathfrak{J}$ is *simple*, i.e., it has no nontrivial closed *-ideals, and any *-epimorphism with domain in $\mathfrak{A}(\mathscr{N}, g\restriction_{\mathscr{N}})/\mathfrak{J}$ is automatically a *-isomorphism.

It's important to choose *boundary *-ideals* $\mathfrak{J}$ which are preserved by $\{\mathfrak{A}\psi : \psi \in G(\mathscr{M}, g)\}$, for only then is $G(\mathscr{M}, g)$ implemented in $\mathfrak{A}(\mathscr{M}, g)/\mathfrak{A}i_{\mathscr{N},\mathscr{M}}(\mathfrak{J})$ by a group *-automorphisms. Local covariance is preserved by the quotient, for $\mathfrak{A}i_{\mathscr{O},\mathscr{N}}(\mathfrak{A}(\mathscr{O}, g\restriction_{\mathscr{O}}))/(\mathfrak{J} \cap \mathfrak{A}i_{\mathscr{O},\mathscr{N}}(\mathfrak{A}(\mathscr{O}, g\restriction_{\mathscr{O}}))) \cong (\mathfrak{A}i_{\mathscr{O},\mathscr{N}}(\mathfrak{A}(\mathscr{O}, g\restriction_{\mathscr{O}})) + \mathfrak{J})/\mathfrak{J}$, by the Second *-Isomorphism Theorem (Theorem B.3, page 124).

Remark 4.2 *As the constructions above make reference uniquely to *-algebraic properties, they naturally extend to* Borchers-Uhlmann *algebras and, as a consequence,*

to locally covariant quantum fields. Indeed, in the case of free fields, the algebraic imposition of boundary conditions supersedes the corresponding functional analytic concept in the one-particle Hilbert *space (i.e., self-adjoint extensions of the Hamiltonian generator of unitary time evolution) – see [Som06] for the particular case of the* Weyl *algebra of the* Klein-Gordon *field in* $\{x \in \mathbb{R}^{1,d-1} : x^{d-1} > 0\}$ *with* Dirichlet *or* Neumann *boundary conditions at* $x^{d-1} = 0$*. Both *-ideals in this example are maximal.*

4.2 Locally covariant quantum theories in AdS

We shall now consider the realization of a locally covariant quantum theory $\mathfrak{A}$ in AdS_d in the spirit of Section 4.1, following the work of Buchholz, Florig and Summers [BFS00]. (The connected component to identity of) the group of isometries of AdS_d induces, as seen in Section 4.1, a group of *-automorphisms $\{\alpha_g : g \in SO_e(2, d-1)\}$ in $\mathfrak{A}(AdS_d)$ which preserves the local structure of $\mathfrak{A}$.

Of particular interest are the (quasi)local algebras associated to wedges $\mathscr{W}_{p,q}$, $p \ll q \in \mathscr{M}in(r)$, $r \in \mathscr{I}$. As seen in Subsection 1.2.2, each $\mathscr{W}_{p,q}$ possesses a one-parameter subgroup of isometries $\{u^\lambda_{p,q}, \lambda \in \mathbb{R}\}$ of AdS_d which preserves the former, given by formula (1.26). We emphasize here a property of $u^\lambda_{p,q}$ obtained in the more general context of Chapter 2 (see page 48) and which will be of great importance in the present Chapter: the asymptotic behaviour of $g^\lambda_{p,q}$ as $\lambda \to \pm\infty$. It follows immediately from formulae (1.26) (page 11), (2.40) (page 48) and the adjoint action of $SO_e(2, d-1)$ that, in Poincaré coordinates,

$$u^\lambda_{p,q}(x,z) \sim \begin{cases} (e^{-\lambda}(x - x(q)), e^{-\lambda}z) & (\lambda \to +\infty) \\ (e^{\lambda}(x - x(p)), e^{\lambda}z) & (\lambda \to -\infty) \end{cases}, \forall (x,z) \in \mathscr{W}_{p,q}, \tag{4.5}$$

i.e., $u^\lambda_{p,q}$ acts asymptotically as scaling transformations around p (resp. q) as $\lambda \to -\infty$ (resp. $\lambda \to +\infty$). Gone through these geometrical preliminaries, we'll specify what we assume from our physical models:

1. The model is given by the realization of a regular, extended locally covariant quantum theory $\mathfrak{A}$ in the collection $\mathscr{W}(AdS_d)$ of the wedges of AdS_d, which, on its turn, provide by local covariance a realization of $\mathfrak{A}$ in AdS_d, according to the construction of Section 4.1.

2. We suppose that the group $\mathfrak{A}u^\lambda_{p,q}$ of *-automorphisms of $\mathfrak{A}(\mathscr{W}_{p,q})$ is strongly continuous for all $(p,q) \in \mathscr{D}(\mathscr{I})$.

3. $\mathfrak{A}$ is locally causal with respect to globally hyperbolic regions.[5]

[5] This condition isn't assumed in [BFS00], which employs a minimal adaptation of the formalism of Haag and Kastler to AdS spacetimes.

We opt for realizing $\mathfrak{A}$ not only in AdS_d, but also in each wedge individually, for this allows us not only to introduce physically reasonable boundary conditions but also to extend a good deal of our following considerations to locally AAdS spacetimes. The condition on the "elementary" states will be discussed in Subsection 4.2.1 below.

We conclude this Subsection by outlining, for completeness, the original form of Rehren duality: [Reh00].

Definition 4.3 *The* Rehren-dual *quantum theory to the realization of* $\mathfrak{A}$ *in* AdS_d *is the* conformally covariant *precosheaf* $\mathscr{D}_{p,q} \mapsto \mathfrak{A}(\mathscr{D}_{p,q}) \doteq \mathfrak{A}(\rho_{AdS_d}^{-1}(\mathscr{D}_{p,q})) = \mathfrak{A}(\mathscr{W}_{p,q})$ *of C*-algebras, indexed by the collection of diamonds in* $\mathscr{I}$*. Here,* ρ_{AdS_d} *is the* Rehren *bijection (1.32) (page 15).*

4.2.1 Boundary values as a dynamical condition

It's natural to understand each $\mathfrak{A}(\mathscr{W}_{p,q})$ as a C*-dynamical system, whose group of *-automorphisms is given by $\mathfrak{A}u^{\lambda}_{p,q}$. The "elementary" states ω in $\mathfrak{A}(AdS_d)$ should represent a situation of thermodynamic equilibrium, since the dynamics is *-automorphic – physically, the quantum dynamical system $(\mathfrak{A}(\mathscr{W}_{p,q}), \mathfrak{A}u^{\cdot}_{p,q})$ is *closed*.

We demand that, for any uniformly accelerated observer associated to $\mathfrak{A}u^{\lambda}_{p,q}$, a state ω over $\mathfrak{A}(\mathscr{W}_{p,q})$ be *passive*, i.e.,

$$-i\omega(U^*\delta_{p,q}U) \geq 0 \tag{4.6}$$

for all $U \in \mathfrak{U}_1(\mathfrak{A}(\mathscr{W}_{p,q})) \cap D(\delta_{p,q})$, where $\delta_{p,q}$ is the derivation which generates $\mathfrak{A}u^{\lambda}_{p,q}$, with domain $D(\delta_{p,q}) \subset \mathfrak{A}(\mathscr{W}_{p,q})$, and $\mathfrak{U}_1(\mathfrak{A}(\mathscr{W}_{p,q}))$ is the connected component to identity of the group of unitary elements of $\mathfrak{A}(\mathscr{W}_{p,q})$. The requirement of passivity means that it's not possible to extract energy from ω by means of a cyclic process – by this, one understands time-dependent perturbations internal to the system which are shut down outside a finite time interval. Equivalently, ω satisfies the second law of thermodynamics in the formulation of Kelvin (in particular, ω is *invariant* under $\mathfrak{A}u^{\lambda}_{p,q}$, guaranteeing the unitary implementability of $\mathfrak{A}u^{\lambda}_{p,q}$ in the GNS Hilbert space $\mathscr{H}_{\omega}$), and the first member of (4.6) corresponds to the *(relative) entropy* of the state $\omega(U^*.U)$ with respect to the state ω (see Appendix B, page 136). We also require that ω satisfies *weak ergodicity (in mean):*

$$\lim_{T\to+\infty} \frac{1}{T} \int_0^T \omega(A\mathfrak{A}u^{\lambda}_{p,q}(B))d\lambda = \omega(A)\omega(B),\ \forall A, B, \tag{4.7}$$

which means that ω is an *extremal* invariant state, that is, any order parameter (i.e., an element of the commutant of the GNS representation of $\mathfrak{A}$ associated to ω which is invariant under the action of $\mathfrak{A}u^{\lambda}_{p,q}$) is a c-number. In this case, it follows that ω

is a KMS state at some inverse temperature $\beta \in [0, +\infty]$, i.e., for all $f \in \mathscr{S}(\mathbb{R})$ with $\hat{f} \in \mathscr{C}_c^\infty(\mathbb{R})$, we have

$$\int_{-\infty}^{+\infty} \omega(A\mathfrak{A}u_{p,q}^\lambda(B))f(\lambda)d\lambda = \int_{-\infty}^{+\infty} \omega(\mathfrak{A}u_{p,q}^\lambda(B)A)f(\lambda + i\beta)d\lambda, \forall A, B. \tag{4.8}$$

The case $\beta = +\infty$, corresponding to the case where ω is a ground state, is excluded by the following argument: considering, for instance, the wedge $\mathscr{W} = \{X : \eta(X, X) = -1, X^1 > |X^0|, X^d > 0\}$ in the fundamental domain (1.5) of AdS_d (page 5), we have that the generator of the isotropy subgroup of $\mathscr{W}$ is the generator M_{01} of $SO_e(2, d-1)$. As $e^{-i\pi M_{12}} M_{01} e^{-i\pi M_{12}} = -M_{01}$, it follows from the invariance of the spectrum under unitary transformations [RS80] that the Hamiltonian in the GNS Hilbert space $\mathscr{H}_\omega$ implementing M_{01} cannot be a positive operator – the same holds, thus, for *all* wedges. Buchholz, Florig and Summers [BFS00] have shown that all ω satisfying (4.6)–(4.7) presents the Unruh *effect*, i.e., they are KMS states at $\beta = \frac{1}{kT} = \frac{2\pi}{\kappa} = 2\pi$, where $\kappa = 1$ is the surface gravity at $\partial_\pm \mathscr{W}_{p,q}$ (Theorem 2.6, page 51). Moreover, we obtain that the realization of $\mathfrak{A}$ in AdS_d is ω-*weakly locally causal* with respect to $\mathscr{W}(AdS_d)$: given $A \in \mathfrak{A}(\mathscr{W}_{p,q})$ and $B \in \mathfrak{A}(\mathscr{W}_{p',q'})$, $\mathscr{W}_{p',q'} \subset \mathscr{W}'_{p,q}$, we have $\omega(AB) = \omega(BA)$, even if we don't demand local causality from $\mathfrak{A}$ in any way.

We'll now establish a sense in which conditions (4.6)–(4.7) correspond to a choice of boundary conditions at conformal infinity. If a state ω over $\mathfrak{A}(\mathscr{W}_{p,q})$ is passive (4.6) and weakly ergodic in mean (4.7) with respect to the group of *-automorphisms $\mathfrak{A}u_{p,q}^\lambda$, then the cyclic vector Ω in the GNS Hilbert space $\mathscr{H}_\omega$ associated to ω is separating with respect to $\pi_\omega(\mathfrak{A}(\mathscr{W}_{p,q}))''$, due to the KMS condition, and $\mathfrak{A}u_{p,q}^\lambda$ extends to a group of *-automorphisms of $\pi_\omega(\mathfrak{A}(\mathscr{W}_{p,q}))''$ which coincides with the Tomita-Takesaki modular group $\sigma_\omega^{(2\pi)^{-1}\lambda}$ (see Appendix B, in particular Lemma B.18, page 140, and Theorem B.19, page 141). Since we've assume passivity and weak ergodicity in mean from an "elementary" state ω over $\mathfrak{A}_{AdS_d}$ with respect to all C*-dynamical systems $(\mathfrak{A}(\mathscr{W}_{p,q}), \mathfrak{A}u_{p,q}^\lambda)$, $p \ll q \in \mathscr{I}$, it follows that such an ω satisfies:

- The Reeh-Schlieder *property* with respect to the collection of wedges $\mathscr{W}(AdS_d)$, i.e., ω is *cyclic* and *separating* with respect to the von Neumann subalgebras $\pi_\omega(\mathfrak{A}(\mathscr{W}_{p,q}))''$,[6] and
- The action of $\mathfrak{A}u_{p,q}^\lambda$ on $\pi_\omega(\mathfrak{A}(\mathscr{W}_{p,q}))''$ is given precisely by the Tomita-Takesaki modular group $\sigma^{(2\pi)^{-1}\lambda}_{\omega\restriction_{\pi_\omega(\mathfrak{A}(\mathscr{W}_{p,q}))''}}$.

$\Delta_{p,q}$ is just the modular operator which implements $\mathfrak{A}u_{p,q}^\lambda$, i.e., $\pi_\omega(\mathfrak{A}u_{p,q}^{2\pi\lambda}(A)) = \Delta_{p,q}^{-i\lambda}\pi_\omega(A)\Delta_{p,q}^{i\lambda}$ for all $A \in \mathfrak{A}(\mathscr{W}_{p,q})$. Let us consider now the inclusion of von Neumann

[6] This property is known by specialists in Conformal Field Theory as the "state-operator correspondence".

algebras $\pi_\omega(\mathfrak{A}(\mathscr{W}_{r,q}))'' \subset \pi_\omega(\mathfrak{A}(\mathscr{W}_{p,q}))''$, where $p \ll_{\mathscr{I}} r \ll_{\mathscr{I}} q$ and, hence, $\mathscr{W}_{r,q} \subset \mathscr{W}_{p,q}$ and $\partial_+\mathscr{W}_{r,q} \subset \partial_+\mathscr{W}_{p,q}$. The inclusion of the respective underlying C*-algebras satisfies the following properties, due to the regularity and the local covariance of $\mathfrak{A}$:

1. $\cup_{\lambda\in\mathbb{R}}\mathfrak{A}u^\lambda_{p,q}(\mathfrak{A}(\mathscr{W}_{r,q}))$ is C*-dense in $\mathfrak{A}(\mathscr{W}_{p,q})$;
2. $\mathfrak{A}u^\lambda_{p,q}(\mathfrak{A}(\mathscr{W}_{r,q})) \subset \mathfrak{A}(\mathscr{W}_{r,q})$ for all $\lambda \geq 0$.

It follows from 1.) and 2.) that the inclusion $\pi_\omega(\mathfrak{A}(\mathscr{W}_{r,q}))'' \subset \pi_\omega(\mathfrak{A}(\mathscr{W}_{p,q}))''$ satisfies the conditions proposed in the work of BORCHERS and YNGVASON [BY99]. The main results of this work, rephrased within our present context, are:

THEOREM 4.1 ([BY99], THEOREM 2.1) *Define* $\mathfrak{M} \doteq \pi_\omega(\mathfrak{A}(\mathscr{W}_{p,q}))''$, $\mathfrak{N} \doteq \pi_\omega(\mathfrak{A}(\mathscr{W}_{r,q}))''$, $\Delta^{i\lambda}_{p,q} \doteq T(-2\pi\lambda)$ *and* $\mathfrak{N}(\lambda) = \mathrm{Ad}T(\lambda)\mathfrak{N} \doteq T(\lambda)\mathfrak{N}T(-\lambda)$. *In this case,*

(i) $\mathrm{Ad}\Delta^{iu}_{r,q}\mathfrak{N}(\lambda) = \mathfrak{N}(\varphi(u,\lambda))$, *where*

$$\varphi(u,\lambda) = \log(1 + e^{-2\pi u}(e^\lambda - 1))$$

for all $(u,\lambda) \in \mathbb{R}^2$ *for which the right hand side is defined. In particular,* $\mathrm{Ad}\Delta^{iu}_{r,q}\mathfrak{M} \subset \mathfrak{M}$ *for all* $u \geq 0$, *and* $\mathfrak{N} = \cap_{u\geq 0}\mathrm{Ad}\Delta^{iu}_{r,q}\mathfrak{M}$.

(ii) The operator $G \doteq \log\Delta_{p,q} - \log\Delta_{r,q}$ *is nonnegative and essentially self-adjoint in a core common to* $\log\Delta_{p,q}$ *and* $\log\Delta_{r,q}$.[7] *The one-parameter unitary group* $\Gamma(\tau) \doteq e^{(2\pi)^{-1}i\tau\bar{G}}$ *satisfies* $T(\lambda)\Gamma(\tau)T(\lambda) = \Gamma(e^\lambda\tau)$ *and* $\mathrm{Ad}\Gamma(\tau)\mathfrak{N}(\lambda) = \mathfrak{N}(\psi(\tau,\lambda))$, *where*

$$\psi(\tau,\lambda) = \lambda + \log(1 + \tau e^{-\lambda})$$

for all $(\tau,\lambda) \in \mathbb{R}^2$ *for which the right hand side is defined. In particular,* $\mathrm{Ad}\Gamma(\tau)\mathfrak{M} \subset \mathfrak{M}$ *and* $\mathrm{Ad}\Gamma(\tau)\mathfrak{N} \subset \mathfrak{N}$ *for all* $\tau \geq 0$, *and* $\mathfrak{N} = \mathrm{Ad}\Gamma(1)\mathfrak{M}$. □

Summing up, Theorem 4.1 not only gives a precise shape to the commutation relations between $\Delta^{i\lambda}_{p,q}$ and $\Delta^{iu}_{r,q}$, but also provides a geometrical realization for the action of $u \mapsto \mathrm{Ad}\Delta^{iu}_{r,q}$ from that of $\lambda \mapsto \mathrm{Ad}\Delta^{i\lambda}_{p,q}$ without our having to assume the existence of the former. The next two Theorems show that the actions of these unitary groups become essentially indistinguishable if we are geometrically sufficiently far from $\partial_-\mathscr{W}_{r,q}$:

THEOREM 4.2 ([BY99], THEOREM 2.2) *Employing the notation of Theorem 4.1, if* $A \in \mathfrak{N}(\lambda)$ *and* $B \in \mathfrak{N}'$, *we have*

$$|\langle B\Omega, \Delta^{iu}_{r,q}A\Omega\rangle - \langle B\Omega, T(-2\pi u)A\Omega\rangle| \leq 2\max\{\|A\Omega\|\|B\Omega\|, \|A^*\Omega\|\|B^*\Omega\|\}.$$

[7] A common core here is a linear subset D dense in $\mathscr{H}_\omega$ such that $D \subset D(\log\Delta_{p,q}) \cap D(\log\Delta_{r,q})$ and such that $\overline{\log\Delta_{p,q}\restriction_D} = \log\Delta_{p,q}$ and $\overline{\log\Delta_{r,q}\restriction_D} = \log\Delta_{r,q}$, where $D(\log\Delta_{p,q}), D(\log\Delta_{r,q}) \subset \mathscr{H}_\omega$ are respectively the domains of the operators $\log\Delta_{p,q}$ and $\log\Delta_{r,q}$.

$$. \min\left\{\frac{|2e^{2\pi u}-1|}{e^{\lambda}-1}, 1\right\}$$

for all $\lambda > 0$, $u \in \mathbb{R}$. □

Employing the notation of Theorem 4.1, we have:

(i)

$$\lim_{\lambda\to+\infty} \|\Delta_{r,q}^{iu}(\mathrm{Ad}T(\lambda)A)\Psi - T(-2\pi u)(\mathrm{Ad}T(\lambda)A)\Psi\| = 0$$

for all $A \in \mathfrak{M}$ *and* $\Psi \in \mathscr{H}_\omega$, *with uniform convergence in* $u \in (-\infty, u_0]$ *for all* $u_0 < +\infty$;

(ii)

$$\lim_{\lambda\to+\infty} \|\Delta_{r,q}^{iu}\mathrm{Ad}T(\lambda)A - T(-2\pi u)\mathrm{Ad}T(\lambda)A\| = 0$$

for all A *in a strongly dense subalgebra of* $\mathfrak{M}$, *with uniform convergence in* $u \in (-\infty, u_0]$ *for all* $u_0 < +\infty$. □

Summing up, denoting $\mathscr{D}_{p,q} = I^-(q, \mathscr{M}in(p) \cong \mathbb{R}^{1,d-2})$, it follows that such groups are essentially indistinguishable from dilations by a factor $e^{-2\pi\lambda}$ around q for $\lambda \to +\infty$. Taking into account the action of the dilation subgroup of $\mathscr{R}^{1,d-2}$ in AdS_d, we obtain that the modular groups associated to ω and the wedges of AdS_d asymptotically implement scaling transformations around points of $\mathscr{I}$, in the precise sense given by Theorems 4.2 and 4.3 – all the above considerations remain valid if we invert the time orientation, i.e., exchanging the roles of p and q.

More in general, inspired in the formalism proposed in [BV95], we write the following

Definition 4.4 *Let* $p \ll q \in \mathscr{M}in(r)$ *for some* $r \in \mathscr{I}$, *where it's assumed that* $\mathscr{M}in(r)$ *is endowed with the vector space operations of* $\mathbb{R}^{1,d-2}$. *The* future scaling algebra $\mathfrak{A}^+(\mathscr{W}_{p,q})$ *(resp.* past scaling algebra $\mathfrak{A}^-(\mathscr{W}_{p,q})$*) in* $\mathscr{W}_{p,q}$ *consists in the uniformly norm-limited functions* $(0,1] \ni \theta \mapsto \underline{A}^{\pm}(\theta) \doteq A^{\pm}_\theta \in \mathfrak{A}(\mathscr{W}_{p,q})$ *satisfying* $A^+_1 \doteq A^+ \in \mathfrak{A}(\mathscr{W}_{r,s}) \Rightarrow A^+_\theta \in \mathfrak{A}(\mathscr{W}_{\theta r+(1-\theta)q,\theta s+(1-\theta)q})$ *(resp.* $A^-_1 \doteq A^- \in \mathfrak{A}(\mathscr{W}_{r,s}) \Rightarrow A^-_\theta \in \mathfrak{A}(\mathscr{W}_{\theta r+(1-\theta)p,\theta s+(1-\theta)p})$*), for all* $p \leq_{\mathscr{I}} r \ll_{\mathscr{I}} s \leq_{\mathscr{I}} q$, $\theta \in (0,1]$. *The algebraic operations are defined pointwise, and the* C^**-norm is given by* $\|\underline{A}^{\pm}\| \doteq \sup_{0<\theta\leq 1} \|A^{\pm}_\theta\|$.

Remark 4.5 *We can say that the concept of scaling algebras defines an abstract version of the* renormalization group *in Local Quantum Physics. A different, cruder definition of scaling algebras associated to wedges in* AdS_d *was proposed by the author in [Rib04].*

Notice that the definition above preserves isotony and local causality of the local subalgebras of $\mathfrak{A}(\mathscr{W}_{p,q})$.

Given a state ω in $\mathfrak{A}(\mathscr{W}_{p,q})$, we can perform its *lift* $\omega^{\pm}_{\theta}$ to $\mathfrak{A}^{\pm}(\mathscr{W}_{p,q})$ at scale $\theta \in (0,1]$ by defining $\omega^{+}_{\theta}(\underline{A}^{+}) \doteq \omega(A^{+}_{\theta})$ and $\omega^{-}_{\theta}(\underline{A}^{-}) \doteq \omega(A^{-}_{\theta})$. Obviously, ω^{+}_{θ} defines a state in $\mathfrak{A}^{\pm}(\mathscr{W}_{p,q})$.

Let's consider $s = q$ in Definition 4.4, i.e., we have the inclusion considered in Theorems 4.1 to 4.3. In this case, we have that $\theta \mapsto A^{+}_{\theta} \doteq \mathfrak{A}u^{f^{-1}_{r,+}(\theta)}_{p,q}(A)$, where $A \in \mathfrak{A}(\mathscr{W}_{r,q})$ and

$$f_{r,+}(\lambda) = \frac{d_{\bar{g}^{(0)}}(u^{\lambda}_{p,q}(r), q)}{d_{\bar{g}^{(0)}}(r,q)},\ f_{r,-}(\lambda) = \frac{d_{\bar{g}^{(0)}}(p, u^{-\lambda}_{p,q}(r))}{d_{\bar{g}^{(0)}}(p,r)}$$

are bijective, $\mathscr{C}^{\infty}$ and strictly decreasing functions from $[0,+\infty) \ni \lambda$ to $(0,1] \ni \theta$, defines an element of $\mathfrak{A}^{+}(\mathscr{W}_{r,q})$ and, thus, of $\mathfrak{A}^{+}(\mathscr{W}_{p,q})$. Conversely, we can define a family of *-automorphisms $\delta^{\mu,+}_{p,q}$ of $\mathfrak{A}^{+}(\mathscr{W}_{p,q})$ implementing scaling transformations around q by a factor $\mu \in (0,1]$ through formula $(\delta^{\mu,+}_{p,q}\underline{A}^{+})(\theta) \doteq A^{+}_{\mu\theta}$. Taking into account the commutation relation

$$\theta u^{\lambda}_{p,q}(r) + (1-\theta)q = u^{\theta\lambda}_{\theta p+(1-\theta)q,q}(\theta r + (1-\theta)q), \tag{4.9}$$

we can induce an action of $\mathfrak{A}u^{\lambda}_{p,q}$ on $\mathfrak{A}^{+}(\mathscr{W}_{p,q})$ through formula

$$\mathfrak{A}^{+}u^{\lambda}_{p,q}(\underline{A}^{+})(\theta) \doteq \mathfrak{A}u^{\theta\lambda}_{\theta p+(1-\theta)q,q}(A^{+}_{\theta}). \tag{4.10}$$

Analogous considerations hold for past scaling algebras by changing notation accordingly, if we invert the time orientation, exchange the roles of r and s and substitute $f_{s,-}$ for $f_{r,+}$. We assume that the action (4.10) and its past counterpart are strongly continuous respectively in $\mathfrak{A}^{+}(\mathscr{W}_{p,q})$ and $\mathfrak{A}^{-}(\mathscr{W}_{p,q})$.

The *raison d'être* of the redundant description of the realization of $\mathfrak{A}$ in AdS_d provided by the past and future scaling algebras comes from the fact that we can use it to define *scaling limits:* let us notice that the net of lifted states $\{\omega^{+}_{\theta} : \theta \in (0,1]\}$ induced by ω has always pointwise convergent subnets (i.e., in the *-weak topology of the dual of $\mathfrak{A}^{+}(\mathscr{W}_{p,q})$) to states $\omega^{+}_{0,\iota}$ over $\mathfrak{A}^{+}(\mathscr{W}_{p,q})$ ($I \ni \iota$ is some index set), by the Banach-Alaoglu theorem [RS80]. The GNS representation $\pi^{+}_{0,\iota}$ associated to each $\omega^{+}_{0,\iota}$ of the elements of $\mathfrak{A}^{+}(\mathscr{W}_{p,q})$ thus have the interpretation of (weak) scaling limits of the local procedures. However, there remains the question of uniqueness of the representations $\pi^{+}_{0,\iota}$ up to *net isomorphisms*, i.e., up to the existence or not of *-isomorphisms $\phi_{\iota,\iota'}$ from $\pi^{+}_{0,\iota}(\mathfrak{A}^{+}(\mathscr{W}_{p,q}))$ to $\pi^{+}_{0,\iota'}(\mathfrak{A}^{+}(\mathscr{W}_{p,q}))$ that preserve the local structure – $\phi_{\iota,\iota'}(\pi^{+}_{0,\iota}(\mathfrak{A}^{+}(\mathscr{W}_{r,s}))) = \pi^{+}_{0,\iota'}(\mathfrak{A}^{+}(\mathscr{W}_{r,s}))$ for all $p \leq_{\mathscr{I}} r \ll_{\mathscr{I}} s \leq_{\mathscr{I}} q$, $\iota, \iota' \in I$. Such a question is partially clarified by the following

Proposition 4.4 *Suppose that ω is a state in $\mathfrak{A}_{AdS_d}$, passive and weakly ergodic in mean with respect to the C*-dynamical systems $(\mathfrak{A}(\mathscr{W}_{p,q}), \mathfrak{A}u^{\lambda}_{p,q})$, for all pairs $(p,q) \in \mathscr{D}(\mathscr{I})$. Then,*

(i) *The limit state* $\omega^+_{0,\iota}$ *is* β*-KMS with* $\beta = 2\pi$ *with respect to the* C^**-dynamical system* $(\pi^+_{0,\iota}(\mathfrak{A}^+(\mathscr{W}_{p,q})), \mathfrak{A}^+_{0,\iota} u^\lambda_{p,q})$*, where* $\mathfrak{A}^+_{0,\iota} u^\lambda_{p,q}(\pi^+_{0,\iota}(A)) \doteq \pi^+_{0,\iota}(\mathfrak{A}^+ u^\lambda_{p,q}(A))$.

(ii) *The scaling limits* $\pi^+_{0,\iota}(\mathfrak{A}^+(\mathscr{W}_{r,s}))$ *and* $\pi^+_{0,\iota}(\mathfrak{A}^+(u^\lambda_{p,q}(\mathscr{W}_{r,s})))$ *are unitarily equivalent, for all* $p \leq_{\mathscr{I}} r \ll_{\mathscr{I}} s \leq_{\mathscr{I}} q$.

(iii) *If* ω *is* primary, *i.e.* $\pi_\omega(\mathfrak{A}_{AdS_d})''$ *is a factor (see Appendix B, page 130), it then satisfies the following property of* holographic local definiteness: *given* $r \in \mathscr{I}$*, we have that*

$$\bigcap_{\substack{(p,q)\in\mathscr{D}(\mathscr{I}),\\ \mathscr{W}_{p,q}\ni r}} \pi_\omega(\mathfrak{A}(\mathscr{W}_{p,q}))'' = \mathbb{C}\mathbb{1}.$$

It follows from this property that the sets of scaling limits $\{(\omega')^+_{0,\iota'} : \iota' \in I'\}$ *of all states* ω' *in the local folium of* ω *with respect to the collection of wedges (i.e.,* $\omega'|_{\mathfrak{A}(\mathscr{W}_{p,q})}$ *defines a normal state over* $\pi_\omega(\mathfrak{A}(\mathscr{W}_{p,q}))''$ *for all* $(p,q) \in \mathscr{D}(\mathscr{I})$*) coincide with* $\{\omega^+_{0,\iota} : \iota \in I\}$*. As a consequence of this and of (i), each* von Neumann *algebra* $\pi_\omega(\mathfrak{A}(\mathscr{W}_{p,q}))''$*, if infinite (which is generally the case in Quantum Field Theory), is type III (see Appendix B, Definition B.5, page 133).*

(iv) *Suppose that the linear map from* $\pi_\omega(\mathfrak{A}(\mathscr{W}_{p,q}))$ *to* $\mathscr{H}_\omega$ *given by*

$$\pi_\omega(A) \mapsto E_{p,q}(K)\pi_\omega(A)\Omega,$$

where $E_{p,q}$ *is the* Borel *measure with values in the projections of* $\mathscr{H}_\omega$ *which defines the spectral resolution of the operator* $\log\Delta_{p,q}$ *and* K *is a compact set in* $\mathbb{R} = \sigma(\log\Delta_{p,q})$*, is a* compact *map, i.e., it takes bounded subsets of the domain to relatively compact subsets of the image, for all* $(p,q) \in \mathscr{D}(\mathscr{I})$.[8] *Then the maps* $\phi_\iota : \pi^+_{0,\iota}(\underline{A}^+) \mapsto w - \lim_\kappa \pi_\omega(\mathfrak{A} u^{-f^{-1}_{p,+}(\theta_\kappa)}_{p,q}(A^+_{\theta_\kappa}))$*, where* $\{\omega^+_{\theta_\kappa}\}_\kappa$ *is a subnet* **-weakly converging to* $\omega^+_{0,\iota}$*, define net isomorphisms to the GNS representation of the original net associated to* ω*. In particular, by virtue of (i), the scaling limits are all unitarily equivalent, and the modular group of any scaling limit acts asymptotically as the modular group associated to* ω^+_1 *as* $\lambda \to +\infty$.

Proof (sketch).

[8] As compact linear maps are characterized for being able to be approximated in norm by finite-rank linear maps with arbitrary precision, such a condition tells us that the spectral density of local excitations of the reference state doesn't grow too fast with the energy. Such a condition on the phase space behaviour of quantum field theories was proposed by Haag and Swieca for the (Hamiltonian) generator of time translations in Minkowski spacetime. IT would be interesting to know is the compactness condition we've given can be deduced, in the same way that the Haag-Swieca compactness condition, from other structural conditions which have been explored in the literature of Algebraic QFT, such as modular nuclearity [Haa96].

(i) It follows from Theorem 5.3.30 in [BR97] that the *-weak limit of a net of β-KMS states is β-KMS.

(ii) It follows immediately from the continuity hypothesis on $\mathfrak{A}^+ u^\lambda_{p,q}$ the existence of a *-isomorphism, which is unitarily implemented due to the existence of cyclic and separating vector for both algebras (see Theorem 2.5.32 in [BR87] for a proof of the latter fact).

(iii) The first assertion is obtained by employing Rehren duality: By virtue of (i) and (ii), the conformal group of $\mathscr{M}in(r) \cong \mathbb{R}^{1,d-2}$ for all $r \in \mathscr{I}$ is unitarily implemented in $\mathscr{H}_\omega$ by the Tomita-Takesaki modular groups associated to $\mathfrak{A} \circ \rho^{-1}_{AdS_d}(\mathscr{D}_{p,q}) = \mathfrak{A}(\mathscr{W}_{p,q})$, and ω is a conformally invariant state of the Rehren-dual theory $\mathfrak{A} \circ \rho^{-1}_{AdS_d}$. By virtue of the resulting commutation relations between the different modular groups, it follows from Proposition 2.3 in [BS93] that the generators P^μ of translations in $\mathscr{M}in(r)$ satisfy the *spectral condition*, i.e., $\sigma(P) \subset J^+(0, \mathbb{R}^{1,d-2})$ – summing up, ω is a vacuum. As ω is primary, it then follows from Theorem 4.6 in [Ara99] that ω is pure and every translationally invariant element of $\mathscr{H}_\omega$ is proportional to Ω. Let now $Z \in \bigcap_{(p,q)\in\mathscr{D}(\mathscr{I}),\ \mathscr{W}_{p,q}\ni r} \pi_\omega(\mathfrak{A}(\mathscr{W}_{p,q}))'' \doteq \mathfrak{N}(r)$. In this case, $Z^* \in \mathfrak{N}(r)$ and, if $\mathbb{R}^{1,d-2} \ni x \alpha_x$ is the group of *-automorphisms which implements translations in $\mathscr{M}in(s) \ni 0 = x(r)$, where x is a Cartesian chart in $\mathscr{M}in(s)$, then $[Z^*, \alpha_x(Z)] = 0$ for all spacelike x and, by weak continuity of α_x, also for $x = 0$ and x lightlike. Consider the bounded function $\mathbb{R} \ni t \mapsto \langle \Omega, Z^* \alpha_{te}(Z)\Omega\rangle$, where e is a lightlike vector in $\mathbb{R}^{1,d-2}$. The Fourier transform of this function has, due to the spectral condition, support in $\bar{\mathbb{R}}_+$. Due to the lightlike commutativity established above, such a function coincides with the function $\mathbb{R} \ni t \mapsto \langle \Omega, \alpha_{te}(Z) Z^* \Omega\rangle$, whose Fourier transform has support in $\bar{\mathbb{R}}_-$. It thus follows that the Fourier transform of the former is supported in $\{0\}$ and, since it's bounded, it's necessarily constant. Therefore, if $x \mapsto U(x)$ is the unitary implementation of translations in $\mathscr{H}_\omega$, it follows that $U(te)Z\Omega = Z\Omega$ for all $t \in \mathbb{R}$ and e lightlike, hence $U(x)Z\Omega = Z\Omega$ for all x. By the uniqueness of Ω, we thus have $Z\Omega = \langle \Omega, Z\Omega\rangle\Omega$. As Ω is separating for $\pi_\omega(\mathfrak{A}(\mathscr{W}_{p,q}))''$ for all $(p,q) \in \mathscr{D}(\mathscr{I})$, we have $Z = \langle \Omega, Z\Omega\rangle \mathbb{1}$, as claimed. The second assertion follows from the fact that, by virtue of holographic local definiteness, $\lim_{\theta \searrow 0} \|(\omega - \omega')\restriction_{\mathfrak{A}(\mathscr{W}_{\theta p + (1-\theta)q, q})}\| = 0$ (this can be proven in a manner analogous to the second part of the proof of Lemma 4.1 in [BV95], whose first part is essentially the proof of the first assertion of (iii) above). The third assertion follows from Driessler's criterion (Theorem B.17, page 137), which, in a certain way, was specially devised for our situation.

(iv) One proves it in a way analogous to Proposition 5.1 in [BV95]. The last assertion then follows from Theorems 4.2 and 4.3.

□

Remark 4.6 *The analog of Proposition 4.4 for past scaling algebras holds, if we invert the time orientation and substitute $f_{q,-}$ for $f_{p,+}$ in item (iv). Moreover, there is a unitary equivalence between past and future scaling limits by virtue of item (ii) of this Proposition.*

Remark 4.7 *The unitary implementability of conformal transformations in the GNS* Hilbert *space of a conformally invariant vacuum by the* Tomita-Takesaki *modular groups was previously demonstrated by* Brunetti, Guido *and* Longo *[BGL93]. Thus, our line of reasoning can be seen as an "holographic" version of this result. Moreover, it follows from Theorem 2.3 (i) in [BGL93] that the *-representation π_ω satisfies* Haag duality *in $\mathscr{I}$ and essential* Haag *duality in $\mathscr{M}in(r)$ for all $r \in \mathscr{I}$ (for the definition and an explanation of the usefulness and meaning of (essential)* Haag *duality within our context, see Subsection 4.4.1, page 86). We can make use of the fact that essential* Haag *duality holds for the scaling limits and refine the conclusion of item (iii), proving that $\pi_\omega(\mathfrak{A}(\mathscr{W}_{p,q}))''$ is a type-III_1 factor if we employ a different criterion for the structure of this algebra [BV95]. We haven't proven this stronger result because we don't need this more refined information in what follows, and* Driessler*'s criterion, besides being simpler, has quite a natural interpretation within our context.*

The equivalence in item (iv) of Proposition 4.4 shows that, indeed, (4.6) and (4.7) act effectively as a condition on scaling limits around points in $\mathscr{I}$, thus revealing the relation between this set of hypotheses and the formulation of Bertola et al. [BBMS00], based in k-point Wightman functions in AdS_d. The boundary conditions employed in this work, similar to the ones proposed by Witten [Wit98] for the AdS/CFT correspondence, are given precisely by a scaling limit of the k-point functions around $\mathscr{I}$. The important item (iii) shows that the scaling limit is, indeed, a condition on the local folium of ω and, hence, on the *-ideals $\mathrm{Ann}\omega\restriction_{\mathfrak{A}(\mathscr{W}_{p,q})}$. Such a conditions is locally covariant by virtue of (ii) and the First *-Isomorphism Theorem (Theorem B.2, page 123), thus defining a locally covariant boundary condition in the sense of Section 4.1.

There is, though, a very important point to be stressed: in the work of Bertola et al., the scaling limit *defines* the dual quantum theory at the boundary, or, equivalently, the boundary quantum theory corresponds in this case to the *universality class* of the bulk quantum theory under the "renormalization group flow" induced by scaling transformations around points in $\mathscr{I}$), which doesn't happen to Rehren duality, which consists in *transplanting* the set that indexes the precosheaf which realizes the locally covariant quantum theory $\mathfrak{A}$ in AdS_d by means of the Rehren bijection. A structural property which illustrates rather well this difference is the Borchers's *timelike tube property:* if $\mathfrak{A}$ is *additive*, and there exists a collection of regions $\mathscr{O}_i \subset \mathscr{M}$ covering an open, arbitrarily small neighbourhood of a timelike curve segment with endpoints

$r \ll s \in \mathscr{M}$ such that $I^+(r) \cap I^-(s)$ is simply connected, then

$$\overline{\bigvee_i \mathfrak{A}(\mathscr{O}_i)} = \mathfrak{A}(I^+(r) \cap I^-(s)).$$

That is, the *germ* of local algebras around a timelike curve segment and, in particular, of a timelike submanifold is *-isomorphic to the local algebra associated to its causal completion (a proof of this fact for quantum fields in real analytic spacetimes can be found in [Str00]). Such a result shows that REHREN duality (Definition 4.3, page 73) really concerns the germs of algebras around $\mathscr{I}$ (whose definition demands, indeed, the imposition of some sort of boundary condition as the one elaborated above [9]), thus being richer than a mere scaling limit. Such a difference also manifests itself in the properties of the local algebras of the dual quantum theory at the boundary in both approaches, as discussed by DĄBROWSKI and REHREN [DR03, Reh05].

Item (iii) of Proposition 4.4 and Theorems 4.2 and 4.3 may, on the other hand, be understood as properties of *return to equilibrium*, if we imagine the isometries $u^\lambda_{p,q}$ in AdS_d as CAUCHY evolutions with boundary conditions at $\mathscr{I}$, in the light of the results of Chapter 2. Hence, we have a strict (we could say "holographic") relation between *scaling limits* around points in $\mathscr{I}$ and *thermalization* of the "elementary" states ω through the dynamics generated by $\mathfrak{A}u^\lambda_{p,q}$. In the case of AdS_d, this return is trivial, for we've seen that ω is a KMS state with respect to this dynamics and, at the same time, ω is a state over the REHREN-dual theory which is invariant under scaling transformations. In the next Section, we'll make use of this more general viewpoint and consider "elementary" states which, by virtue of nontrivial gravitational effects, return, accordingly, "nontrivially" to equilibrium and, equivalently, present only asymptotic scaling invariance around $\mathscr{I}$.

4.3 Algebraic holography in AAdS spacetimes

We've finally arrived at the moment of defining the quantum theories of interest to us in AAdS spacetimes $(\mathscr{M}, g)$ satisfying the hypotheses of Theorem 1.3, which guarantees that $I^+(p, \overline{\mathscr{M}}) \cap I^-(q, \overline{\mathscr{M}}) \cap \mathscr{I} = I^+(p, \mathscr{I}) \cap I^-(q, \mathscr{I}) \doteq \mathscr{D}_{p,q}$ for all $(p, q) \in \mathscr{D}(\mathscr{I})$. The most delicate part concerns the implementation of the asymptotic isometries associated to wedges constructed in Chapter 2 (pages 46–50). However, before that, let us write:

[9] Actually, we've committed a slight abuse in our definition of BORCHERS's timelike tube property. The correct formulation (which we won't use) involves the local VON NEUMANN algebras and weak topology closures, associated to a GNS representation generated by a state ω. Hence, the definition also involves properties of ω, or, more precisely, of its local folium.

Definition 4.8 *The quantum theory* Rehren-dual *to the realization of* $\mathfrak{A}$ *in* $(\mathscr{M}, g)$ *is the precosheaf* $\mathscr{D}_{p,q} \mapsto \mathfrak{A}(\mathscr{D}_{p,q}) \doteq \mathfrak{A}(\rho^{-1}_{(\mathscr{M},g)}(\mathscr{D}_{p,q})) = \mathfrak{A}(\mathscr{W}_{p,q})$ *of* C^**-algebras, indexed by the collection of diamonds in* $\mathscr{I}$*. Here,* $\rho_{(\mathscr{M},g)}$ *is the* Rehren *bijection (1.32), introduced in page 15.*

Let us now consider a wedge $\mathscr{W}_{p,q}$ with geodesically convex closure. We'll define, for $\lambda \in \mathbb{R}$, $\epsilon > 0$ fixed, the regions

$$\mathscr{W}^{+}_{p,q}(\lambda) \doteq (\lambda^{\bar{g}}_{p,q})^{-1}((\lambda - \epsilon, +\infty)) \cap \mathscr{M},\ \mathscr{W}^{-}_{p,q}(\lambda) \doteq (\lambda^{\bar{g}}_{p,q})^{-1}((-\infty, \epsilon - \lambda)) \cap \mathscr{M}, \tag{4.11}$$

and, correspondingly in $\mathscr{I}$,

$$\mathscr{D}^{+}_{p,q}(\lambda) \doteq (\lambda^{\bar{g}}_{p,q})^{-1}((\lambda - \epsilon, +\infty)) \cap \mathscr{I},\ \mathscr{D}^{-}_{p,q}(\lambda) \doteq (\lambda^{\bar{g}}_{p,q})^{-1}((-\infty, \epsilon - \lambda)) \cap \mathscr{I}. \tag{4.12}$$

Let $\mathfrak{A}$ be a regular, extended locally covariant quantum theory, satisfying the following hypotheses:

(a) The realization of $\mathfrak{A}$ in $(\mathscr{M}, g)$ is locally causal with respect to the collection of wedges $\mathscr{W}(\mathscr{M}, g)$;

(b) $\mathfrak{A}$ is weakly primitively causal with respect to the wedges of $(\mathscr{M}, g)$ (thus guaranteeing the possibility of imposing a time evolution subject to boundary conditions);

(c) There exists a state space $\mathfrak{S}$ such that $\mathfrak{S}(\mathscr{M}, g)$ possesses a primary representative ω satisfying the property of holographic local definiteness (defined in item (iii) of Proposition 4.4, page 77) and such that its local folium with respect to the causally convex subregions of wedges is contained in $\mathfrak{S}(\mathscr{M}, g)$.

In what follows, we've uncluttered the notation by omitting the symbols of restriction of the metric g to subregions and those of quotients modulo nontrivial *-ideals which annihilate the propagators (*boundary *-ideals*), having always in mind the *-Isomorphism Theorems. It follows from (b) that $\mathfrak{A}(\mathscr{W}^{\pm}_{p,q}(\lambda), g)$ is *-isomorphic to $\mathfrak{A}(\mathscr{W}_{p,q})$ for all γ, $(p, q) \in \mathscr{D}(\mathscr{I})$, modulo boundary *-ideals. Moreover, we can define the *germ* $\mathfrak{A}(\partial_{\pm}\mathscr{W}_{p,q})$ of $\mathfrak{A}(\mathscr{W}_{p,q})$ at the past and future horizons $\partial_{\pm}\mathscr{W}_{p,q}$ by the projective limits (modulo boundary *-ideals)

$$\mathfrak{A}(\partial_{\pm}\mathscr{W}_{p,q}) \doteq \varprojlim_{\lambda \nearrow +\infty, \mathfrak{A}} \mathfrak{A}(\mathscr{W}^{\pm}_{p,q}(\lambda), g), \tag{4.13}$$

recalling that the equivalence relation which defines the limit is given by the identification (modulo boundary *-ideals) $A \in \mathfrak{A}(\mathscr{W}^{\pm}_{p,q}(\lambda), g) \sim (\mathfrak{A}i_{\mathscr{W}^{\pm}_{p,q}(\lambda), \mathscr{W}^{\pm}_{p,q}(\lambda')})^{-1}(A)$, for all $\lambda' \geq \lambda$, similarly to the case of germs of local algebras at Cauchy surfaces, defined in

Chapter 3 (see formula (3.6), page 61).

With the above definitions, if $r \in (\lambda^{\bar{g}}_{p,q})^{-1}(\lambda_0) \cap \mathscr{I}$ and $r \ll_{\mathscr{I}} s \leq_{\mathscr{I}} q$, obviously $\mathscr{W}_{r,s} \subset \mathscr{W}^+_{p,q}(\lambda_0)$ and, hence, $\mathscr{W}_{u^\lambda_{p,q}(r),u^\lambda_{p,q}(s)} \subset \mathscr{W}^+_{p,q}(\lambda_0+\lambda)$ for all $\lambda \geq 0$.

Let $p \ll q \in \mathscr{M}in(r)$ for some $r \in \mathscr{I}$, where it's assumed that $\mathscr{M}in(r)$ is is endowed with the vector space operations of $\mathbb{R}^{1,d-2}$. The future scaling algebra $\mathfrak{A}^+(\mathscr{W}_{p,q})$ *(resp.* past scaling algebra $\mathfrak{A}^-(\mathscr{W}_{p,q})$*) in $\mathscr{W}_{p,q}$ consists of the uniformly norm-bounded continuous functions $(0,1] \ni \theta \mapsto \underline{A}^+(\theta) \doteq A^+_\theta \in \mathfrak{A}(\mathscr{W}^+_{p,q}(f^{-1}_{p,+}(\theta)), g)$ (resp. $(0,1] \ni \theta \mapsto \underline{A}^-(\theta) \doteq A^-_\theta \in \mathfrak{A}(\mathscr{W}^-_{p,q}(f^{-1}_{q,-}(\theta)), g)$) satisfying $A^+_1 \doteq A^+ \in \mathfrak{A}(\mathscr{W}_{r,s})$, $r \in \mathscr{W}^+_{p,q}(\lambda_0)$ for some λ_0 implies $A^+_\theta \in \mathfrak{A}(\mathscr{W}^+_{p,q}(\lambda_0 + f^{-1}_{r,+}(\theta)), g)$ (resp. $A^-_1 \doteq A^- \in \mathfrak{A}(\mathscr{W}_{r,s})$, $s \in \mathscr{W}^-_{p,q}(\lambda_0)$ for some λ_0 implies $A^-_\theta \in \mathfrak{A}(\mathscr{W}^-_{p,q}(\lambda_0 + f^{-1}_{s,-}(\theta)), g)$), for all $p \leq_{\mathscr{I}} r \ll_{\mathscr{I}} s \leq_{\mathscr{I}} q$, $\theta \in (0,1]$, where $f_{r,+}$ and $f_{s,-}$ are defined in page 77. The algebraic operations are defined pointwise, and the C*-norm is given by $\|\underline{A}^\pm\| \doteq \sup_{0<\theta\leq 1} \|A^\pm_\theta\|$.*

The more general definition given above copes with the loss of "spatial" localization caused by the action of the *propagators*

$$\alpha^\pm_{\lambda,\lambda'} \doteq (\mathfrak{A}i_{\mathscr{W}^\pm_{p,q}(\lambda'),\mathscr{W}_{p,q}})^{-1} \circ \mathfrak{A}i_{\mathscr{W}^\pm_{p,q}(\lambda),\mathscr{W}_{p,q}}, \tag{4.14}$$

which then can be used to build elements of $\mathfrak{A}^\pm(\mathscr{W}_{p,q})$ (modulo boundary *-ideals) in the same way we've employed $\mathfrak{A}u^\lambda_{p,q}$ in the case of AdS_d. We suppose, accordingly, that the propagators (4.14) are strongly continuous in the norm of the scaling algebras.

We stress now a *very* important point: the images of the elements of the scaling algebras belong to the *intrinsic* algebras, and not to their images in $\mathfrak{A}(\mathscr{W}_{p,q})$, which identify elements which differ by an element of a boundary *-ideal. This eliminates an apparent contradiction between the germs of local algebras at the past and future horizons and the principle of holographic local definiteness. The maps

$$A^+_\theta \mapsto I^+_{p,q}(\underline{A}^+)(\theta) \doteq \mathfrak{A}i_{\mathscr{W}^+_{p,q}(f^{-1}_{p,+}(\theta)),\mathscr{W}_{p,q}}(A^+_\theta) \tag{4.15}$$

and

$$A^-_\theta \mapsto I^-_{p,q}(\underline{A}^-)(\theta) \doteq \mathfrak{A}i_{\mathscr{W}^-_{p,q}(f^{-1}_{q,-}(\theta)),\mathscr{W}_{p,q}}(A^-_\theta) \tag{4.16}$$

define *-morphisms from the scaling algebras to the uniformly bounded continuous functions $(0,1] \ni \theta$ in $\mathfrak{A}(\mathscr{W}_{p,q})$ and preserve the "time" localization. Defining the lift $\omega^\pm_\theta$ of our reference state ω to $I^\pm_{p,q}(\mathfrak{A}^\pm(\mathscr{W}_{p,q}))$ at scale θ in the same manner as in the AdS case, we can finally impose our boundary conditions:

(d)

$$\mathrm{Ann}\mathfrak{S}(\mathscr{W}^{\pm}_{p,q}(\lambda),g) \doteq \bigcap_{\omega'\in\mathfrak{S}(\mathscr{W}^{\pm}_{p,q}(\lambda),g)} \mathrm{Ann}\omega' \supset \mathrm{Ker}\mathfrak{A}i_{\mathscr{W}^{\pm}_{p,q}(\lambda),\mathscr{W}_{p,q}}$$

for all $\lambda \in \mathbb{R}$, and the scaling limits $\omega^{\pm}_{0,\iota}$ define GNS representations linked by net *-isomorphisms with respect to the collection of wedges in $(\mathscr{M},g)$, and *-isomorphic to the *-representation of $\mathfrak{A}(\partial_{\pm}\mathscr{W}_{p,q})$ induced by the quotient modulo $\mathrm{Ann}\mathfrak{S}(\partial_{\pm}\mathscr{W}^{\pm}_{p,q})$ (this state space is defined by employing the contravariance of $\mathfrak{S}$ and the universality of projective limits). The *-automorphic action $(0,1] \ni \mu \mapsto \delta^{\mu,\pm}_{p,q}$ of scaling transformations on $\pi_{0,\iota}(I^{\pm}_{p,q}(\mathfrak{A}^{\pm}(\mathscr{W}_{p,q})))$ is unitarily implemented by the formula

$$\pi_{0,\iota}(\delta^{\exp(\pm\kappa_{\pm}\lambda),\pm}_{p,q}(I^{\pm}_{p,q}(\underline{A}^{\pm}))) = \mathrm{Ad}\Delta^{\frac{i}{2\pi}\lambda}_{p,q}\pi_{0,\iota}(I^{\pm}_{p,q}(\underline{A}^{\pm})), \tag{4.17}$$

where $\Delta_{p,q}$ is the Tomita-Takesaki modular operator associated to $\omega_{0,\iota}$ (which is, hence, essentially unique for all ι), and $\kappa_{\pm} = \mp 1$ is the asymptotic past/future surface gravity of $\mathscr{W}_{p,q}$, obtained in Theorem 2.6 (page 51).

Condition (d), albeit rather complicated, is naturally motivated by Proposition 4.4 and Theorem 2.6 in the case of AdS_d. Together with (c), it's not only common to all states in the local folium of ω but also implies that the von Neumann algebras $\pi_{\omega}(\mathfrak{A}(\mathscr{W}_{p,q}))''$ are type III, analogously to item (iii) of Proposition 4.4. The assumed continuity of the propagators, together with local covariance, guarantees that the scaling limits of different wedges all live in the same Hilbert space, as in item (ii). The geometrical character *at* $\mathscr{I}$ of the action of the modular groups, on its turn, follows from Theorem 4.1. By virtue of the asymptotic value we've found for the surface gravities of wedges and the commutation relations between the different modular groups, resulting from Theorem 4.1, we have that the modular groups act geometrically exactly as conformal transformations in $\mathscr{I}$, and the scaling limit $\omega_{0,\iota}$ defines a conformally invariant vacuum [BS93] over the precosheaf $\mathscr{D}_{p,q} \mapsto \pi_{0,\iota}((I^{\pm}_{p,q}(\mathfrak{A}^{\pm}\circ\rho^{-1}_{(\mathscr{M},g)})(\mathscr{D}_{p,q}))''$. Summing up, the scaling limit we've defined loses *all geometrical information about* $(\mathscr{M},g)$, and could be taken as a quantum theory Rehren-dual to a quantum theory in AdS_d. Moreover, the implementation of asymptotic isometries only occurs in the scaling limit. We notice here the strong kinship of the construction above to the *lightfront holography* proposed by Schroer [Sch06, Sch05].

We've emphasized a *dynamical* definition as Definition 4.9 for the scaling algebras in AAdS wedges, instead of the simpler Definition 4.4 given in the case of AdS_d (which, in principle, also applies here) due to several reasons. First, the boundary condition (d) cast after Definition 4.9 emphasizes the character of *return to equilibrium* dual to the scaling limit, and extends to more general contexts (i.e., relatively compact diamonds in causally simple spacetimes), such as the construction of global time functions in Chapter

2. Second, it opens up the possibility of dynamically probing deviations from thermal equilibrium in the same way we probe deviations from critical (i.e., scale invariant) behaviour by means of, say, operator product expansions [FH81, Wei96, Bos05, BOR02].

4.4 Holographic analysis of superselection sectors

The development which concludes this Thesis consists in showing how the geometrical properties of nontrivial AAdS spacetimes (i.e., different from AdS_d) within the hypotheses employed throughout Chapter 1 modify in a radical way not only the implementation of *geometrical* symmetries, but also of *internal* symmetries, encoded in the structure of superselection sectors. In what follows, we'll make the following set of hypotheses about the spacetimes and the locally covariant quantum theories of our interest:

(i) $(\mathscr{M}, g)$ is an AAdS spacetime satisfying the hypotheses of Proposition 1.8;

(ii) $\mathfrak{A}$ is an extended locally covariant quantum theory, whose realization in $(\mathscr{M}, g)$ satisfies local causality w.r.. $\mathscr{W}(\mathscr{M})$;

(iii) We have at our disposal a state space $\mathfrak{S}_0$, called *reference state space*, such that any $\omega, \omega' \in \mathfrak{S}_0(\mathscr{W}_{p,q})$ satisfy condition (3.8) (page 64) for all $(p, q) \in \mathscr{D}(\mathscr{I})$. In particular, as we've seen, the VON NEUMANN algebras $\pi_\omega(\mathfrak{A}(\mathscr{W}_{p,q}))''$, $\omega \in \mathfrak{S}_0(\mathscr{W}_{p,q})$, are all *-isomorphic.

(iv) $\mathfrak{S}_0(\mathscr{W}_{p,q})$ has a representative ω_0, (also) called *reference state*, which satisfies:

- The REEH-SCHLIEDER property, i.e., the cyclic vector Ω_0 in the GNS HILBERT space $\mathscr{H}_0 \doteq \mathscr{H}_{\omega_0}$ associated to ω_0 is *separating* for $\pi_0(\mathfrak{A}(\mathscr{W}_{p,q}))''$, where $\pi_0 \doteq \pi_{\omega_0}$ is the GNS *-representation associated to ω_0;
- The property of holographic local definiteness;
- The property (III), i.e., $\pi_0(\mathfrak{A}(\mathscr{W}_{p,q}))''$ is type III for all $(p, q) \in \mathscr{D}(\mathscr{I})$ (see Appendix B, Definition B.5, page 133).

Additional conditions will be included from time to time alongside the following developments. Notice that the states ω_0 satisfying the boundary conditions (c)–(d) imposed in the previous Section satisfy the properties listed by hypothesis (iv).

REMARK 4.10 *Hypotheses (iii) and (iv) on the reference state space $\mathfrak{S}_0$ are similar to the ones adopted by* BRUNETTI *and* RUZZI *[BR07] in their locally covariant formulation for the algebraic theory of superselection sectors. We shan't adopt, though, the full-fledged approach of this work for simplicity, since we'll assess only basic elements of this theory in what follows.*

4.4.1 A digression. Haag duality and elements of the algebraic theory of superselection sectors

The fact that the collection $\mathscr{W}(AdS_d)$ of wedges in AdS_d is closed under the operation of taking the causal complement (more precisely, formulae (1.29) are valid) guarantees that not only local causality is preserved by the Rehren bijection, but also the possible maximality of the local algebras with respect to this property. We'll see in the next Subsection that, due to the geometry of the wedges in nontrivial AAdS spacetimes, this maximality acquires contours of a *"no-go"* theorem about the nontriviality of the local algebras in the bulk of such spacetimes, with implications on the structure of superselection sectors. However, before that, we shall formulate in a precise manner this maximality hypothesis, and illustrate its role in the determination of this structure.

Definition 4.11 *Let $\mathfrak{A}$ be an extended locally covariant quantum theory, $(\mathscr{M},g)\in Obj\mathscr{S}tc_d$, π a *-representation of $\mathfrak{A}_{\mathscr{M}}$ in the* Hilbert *space $\mathscr{H}$, and $\mathscr{Q}(\mathscr{M},g)$ a collection of causally complete open sets of $(\mathscr{M},g)$. We say that the realization of $\mathfrak{A}$ in $(\mathscr{M},g)$ is* π-Haag-dual *with respect to $\mathscr{Q}(\mathscr{M},g)$ if the* von Neumann *algebras associated to π and localized at elements of $\mathscr{Q}(\mathscr{M},g)$ are* maximally *locally causal, i.e., $\pi(\mathfrak{A}(\mathscr{O}))''=\pi(\mathfrak{A}(\mathscr{O}'))'$, where $\pi(\mathfrak{A}(\mathscr{O}))'$ denotes the commutant of $\pi(\mathfrak{A}(\mathscr{O}))$ in $\mathscr{H}$ (see in Appendix B the discussion immediately preceding Definition B.4, page 130) – recall that local causality of $\mathfrak{A}$ only implies $\pi(\mathfrak{A}(\mathscr{O}'))''\subset\pi(\mathfrak{A}(\mathscr{O}))'$. More in general, we say that the realization of $\mathfrak{A}$ in $(\mathscr{M},g)$ is* essentially π-Haag-dual *with respect to $\mathscr{Q}(\mathscr{M},g)$ if the* Haag-dual *extension $\mathscr{Q}(\mathscr{M},g)\ni\mathscr{O}\mapsto\pi(\mathfrak{A}(\mathscr{O}))^d\doteq\pi(\mathfrak{A}(\mathscr{O}'))'$ of the realization of $\mathfrak{A}$ in $(\mathscr{M},g)$ in the representation π is* locally causal, *i.e., $\pi(\mathfrak{A}(\mathscr{O}_1))^d\subset(\pi(\mathfrak{A}(\mathscr{O}_2))^d)'$ for all pairs $\mathscr{O}_1,\mathscr{O}_2\in\mathscr{Q}(\mathscr{M},g)$ such that $\mathscr{O}_1\perp\mathscr{O}_2$. A *-representation π of $\mathfrak{A}_{\mathscr{M}}$ is said to be (resp. essentially)* Haag*-dual if the realization of $\mathfrak{A}$ in $(\mathscr{M},g)$ is (resp. essentially) π-*Haag*-dual.*

Remark 4.12 *There is a subtlety in the definition of essential* Haag *duality: it's immediate to see that, if $\mathscr{Q}(\mathscr{M},g)$ is closed under causal complements, i.e., $\mathscr{O}\in\mathscr{Q}(\mathscr{M},g)\Leftrightarrow\mathscr{O}'\in\mathscr{Q}(\mathscr{M},g)$, then* Haag *duality and essential* Haag *duality with respect to $\mathscr{Q}(\mathscr{M},g)$ coincide. Hence, the interesting effects that arise when the realization of $\mathfrak{A}$ in $(\mathscr{M},g)$ is only essentially π-Haag-dual with respect to $\mathscr{Q}(\mathscr{M},g)$ appear only when the latter is* not *closed under causal complements. The typical example is $(\mathscr{M},g)=\mathbb{R}^{1,d-1}$ and $\mathscr{Q}(\mathbb{R}^{1,d-1})=\{I^+(p)\cap I^+(q):p\ll q\in\mathbb{R}^{1,d-1}\}$.*

The breakdown of Haag duality implies, for instance, the spontaneous breaking of internal symmetries in the representation induced by a vacuum in theories of local observables in Minkowski spacetime [Haa96]. The reason, in intuitive terms, is the following: the presence of internal symmetries, manifest in the structure of the particle multiplets which appear in scattering processes, is encoded in the local algebras by

means of intertwiners [10] between the unitary representations of the group of internal symmetries. HAAG duality guarantees that there are enough intertwiners to completely reconstruct the representation theory of this group (= structure of superselection sectors) and, hence, the group itself – this result, considered one of the great triumphs of the algebraic approach to QFT, was demonstrated by DOPLICHER and ROBERTS [Ara99, Haa96], and constitutes an abstract formulation of duality between compact groups and their irreducible representations (TANNAKA-KREĬN theorem).

If only essential HAAG duality holds, it's possible to reconstruct the part of internal symmetries which doesn't go under spontaneous breaking by repeating the procedure of DOPLICHER and ROBERTS for the HAAG-dual extension of the local algebras, according to Definition 4.11. The procedure used to identify the spontaneously broken part of the internal symmetries is, however, a bit more subtle [Haa96].

Let's now make more precise how the structure of superselection sectors is encoded in the realization of $\mathfrak{A}$ in $(\mathscr{M}, g)$, following the treatment in [Ara99] and [GLRV01], to which we refer for details. Let $\mathfrak{A}$, $(\mathscr{M}, g)$ and $\mathscr{Q}(\mathscr{M}, g)$ as in Definition 4.11, and π_0 a HAAG-dual *-representation of $\mathfrak{A}_{\mathscr{M}}$ in the HILBERT space $\mathscr{H}_0$, which we'll adopt as reference. We say that a *-representation π of $\mathfrak{A}_{\mathscr{M}}$ in a HILBERT space $\mathscr{H}$ *satisfies the DHR criterion* [11] with respect to $\mathscr{Q}(\mathscr{M}, g)$ if, for all $\mathscr{O} \in \mathscr{Q}(\mathscr{M}, g)$, $\pi\restriction_{\mathfrak{A}(\mathscr{O}')}$ is unitarily equivalent to $\pi_0\restriction_{\mathfrak{A}(\mathscr{O}')}$. Let, then, V be a unitary operator from $\mathscr{H}$ to $\mathscr{H}_0$ such that $V\pi(A)V^* = \pi_0(A)$ for all $A \in \mathfrak{A}(\mathscr{O}')$. Define the following *-morphism from $\pi_0(\mathfrak{A}_{\mathscr{M}})$ to $\mathscr{B}(\mathscr{H}_0)$:

$$\rho(\pi_0(A)) \doteq V\pi(A)V^*. \tag{4.18}$$

By construction, $\rho\circ\pi_0$ is unitarily equivalent to π. Assuming, for now, that $\mathscr{Q}(\mathscr{M}, g)$ is *directed* under the partial ordering induced by set inclusions, consider $\mathscr{O}_1, \mathscr{O}_2 \in \mathscr{Q}(\mathscr{M}, g)$ such that $\mathscr{O}_2 \supset \mathscr{O}, \mathscr{O}_1$. Then,

$$[\rho(\pi_0(A)), \rho(\pi_0(B))] = [\rho(\pi_0(A)), \pi_0(B)] = 0 \tag{4.19}$$

for all $A \in \mathfrak{A}(\mathscr{O}_1)$ and $B \in \mathfrak{A}(\mathscr{O}_2') \subset \mathfrak{A}(\mathscr{O}')$. Due to HAAG duality, we have $\rho(\pi_0(A)) \in \pi_0(\mathfrak{A}(\mathscr{O}_2'))' = \pi_0(\mathfrak{A}(\mathscr{O}_2))''$. It follows from the definition of quasilocal algebra that $\rho \circ \pi_0(\mathfrak{A}_{\mathscr{M}}) \subset \rho(\pi_0(\mathfrak{A}_{\mathscr{M}})'') \subset \pi_0(\mathfrak{A}_{\mathscr{M}})''$.

[10] Given a group G and two unitary representations $G \ni g \mapsto U_1(g), U_2(g)$ of G respectively in HILBERT spaces $\mathscr{H}_1$ and $\mathscr{H}_2$, a *intertwiner* between U_1 and U_2 is a bounded linear operator $T : \mathscr{H}_1 \to \mathscr{H}_2$ such that $TU_1(g) = U_2(g)T$ for all $g \in G$. In particular, such a definition applies, as it'll be often used, to *-representations of *-algebras.

[11] DHR stands for the names of DOPLICHER, HAAG and ROBERTS.

The argument above is a recurrent line of reasoning in the algebraic theory of superselection sectors. With it, we can define the transport of localized *-endomorphisms of $\pi_0(\mathfrak{A}_{\mathscr{M}})''$ without the aid of the implementation of a translation group. More precisely, we have the following facts:

- Given two *-endomorphisms ρ_1, ρ_2 localized respectively in $\mathscr{O}_1$ and $\mathscr{O}_2$ and such that the representations $\rho_1 \circ \pi_0$ and $\rho_2 \circ \pi_0$ are unitarily equivalent, i.e., $\rho_1 \circ \pi_0(A) = U\rho_2 \circ \pi_0(A)U^*$, then $U \in \pi_0(\mathfrak{A}(\mathscr{O}_3'))'$, where $\mathscr{O}_3 \supset \mathscr{O}_1, \mathscr{O}_2$. That is, *-endomorphisms localized in $\mathscr{O}$ can be transported to any other element of $\mathscr{Q}(\mathscr{M}, g)$ by means of *local* intertwiners.

- The composition $\rho_1 \circ \rho_2$ is clearly localized in $\mathscr{O}_3 \supset \mathscr{O}_1, \mathscr{O}_2$, where ρ_i is localized in $\mathscr{O}_i$. Moreover, if $\mathscr{O}_1 \perp \mathscr{O}_2$, then $\rho_1 \circ \rho_2 = \rho_2 \circ \rho_1$.

A *-endomorphism ρ is said to be *irreducible* if the *-representation $\rho \circ \pi_0$ is. In this case, we say that the set of pure states in the folium of $\rho \circ \pi_0$ is the *(superselection) sector* associated to ρ.

Let us denote the Banach space of (local) intertwiners between localized *-endomorphisms ρ and σ by $(\rho, \sigma) \doteq \{U : U\rho(A) = \sigma(A)U, \forall A\}$. Given $T \in (\rho, \sigma)$, $T' \in (\rho', \sigma')$ we can define the *(tensor) product* $T \otimes T' \doteq T\rho(T') \in (\rho \circ \rho', \sigma \circ \sigma')$. We can prove that, if T and T' are causally disjoint, then $T \otimes T' = T' \otimes T$. It thus follows that, if σ and σ' are localized in causally disjoint regions, then $(T' \otimes T)^* \circ (T \otimes T') \doteq \epsilon(\rho, \rho') \in (\rho \circ \rho', \rho' \circ \rho)$ is *independent* of σ, σ' or T, T'. Generalizing to n localized *-endomorphisms ρ_i, $i = 1, \ldots, n$, we obtain intertwiners $\epsilon_p(\rho_1, \ldots, \rho_n)$ for each permutation p of $\{1, \ldots, n\}$, in such a way that $\epsilon_{p'}(\rho_1, \ldots, \rho_n) \circ \epsilon_p(\rho_1, \ldots, \rho_n) = \epsilon_{p' \circ p}(\rho_1, \ldots, \rho_n)$. In this manner, if $\rho_1 = \cdots = \rho_n = \rho$ the map $S_n \ni p \mapsto \epsilon_p(\rho, \ldots, \rho) \in \rho^n(\pi_0(\mathfrak{A}_{\mathscr{M}}))'$ constitutes a unitary representation of the group S_n of permutations of n elements in $\rho^n \circ \pi_0$. The operators ϵ are called *statistics operators*, for they reflect the spin-statistics relation of the superselection sectors. It's precisely these operators, together with additional structures (conjugates, left inverses), who allow one to rebuild the group of internal symmetries and the particle multiplets.

In the same way that the composition of *-endomorphisms corresponds to the tensor product of representations, we can describe subrepresentations and direct sums by *subobjects* and *direct sums* of *-endomorphisms: we say that σ is a *subobject* of ρ if there exists an *isometry* $W \in (\sigma, \rho)$ (i.e., $W^*W = \mathbb{1}$ and, hence, WW^* is a projection), and τ is a *direct sum* of ρ and σ if there exist isometries $V \in (\rho, \tau)$, $W \in (\sigma, \tau)$ such that $VV^* + WW^* = \mathbb{1}$ and, hence, $\tau(.) = V\rho(.)V^* + W\sigma(.)W^*$. To guarantee that these *-endomorphisms remain localized, we need to invoke the so-called *property B of* Borchers, valid, for instance, for type-III factors: given $\mathscr{O}_1, \mathscr{O}_2 \in \mathscr{Q}(\mathscr{M}, g)$ such that

$\overline{\mathscr{O}_1} \subset \mathscr{O}_2$ and a projection $E \in \pi_0(\mathfrak{A}(\mathscr{O}_1))''$, there exists an isometry $W \in \pi_0(\mathfrak{A}(\mathscr{O}_1))''$ such that $WW^* = E$. In the case of type-III factors, we can take $\mathscr{O}_2 = \mathscr{O}_1$.

When $\mathscr{Q}(\mathscr{M}, g)$ is not directed, the *-morphism ρ defined in (4.18) is defined only in the precosheaf $\mathscr{Q}(\mathscr{M}, g) \ni \mathscr{O} \subset \mathscr{O}_1 \mapsto \pi_0(\mathfrak{A}(\mathscr{O}_1))''$, and here satisfies $A \in \mathfrak{A}(\mathscr{O}_1) \Rightarrow \rho(\pi_0(A)) \in \pi_0(\mathfrak{A}(\mathscr{O}_1))''$.

4.4.2 (Essential) Haag duality in (A)AdS spacetimes

Now, with the adequate language at hand, let us go back to discussing the question raised in the first paragraph of the previous Subsection. Let us consider and AAdS spacetime $(\mathscr{M}, g)$, an extended locally covariant quantum theory $\mathfrak{A}$ realized in $(\mathscr{M}, g)$ and π_0 a reference *-representation of $\mathfrak{A}_{\mathscr{M}}$, which we assume irreducible for simplicity. Once more, $\mathscr{D}(AdS_d)$ as well as $\mathscr{W}(AdS_d)$ are closed under causal complements, due to formulae (1.29) (page 13). Employing the Rehren bijection (1.32) (page 15), it automatically follows that the realization of $\mathfrak{A}$ in AdS_d is π_0-Haag-dual with respect to $\mathscr{W}(AdS_d)$ if and only if the Rehren-dual realization $\mathfrak{A}$ in ESU_{d-1} is with respect to $\mathscr{D}(AdS_d)$ – recall as well that, in both cases, Haag duality and essential Haag duality coincide.

Let1s see now what happens if $(\mathscr{M}, g)$ is a nontrivial AAdS spacetime, satisfying the hypotheses of Theorem 1.3, page 16. In this case, $\mathscr{W}(\mathscr{M}, g)$ is no longer closed under causal complements, there thus being a fundamental difference between Haag duality and essential Haag duality. Nevertheless, still holds the following

Lemma 4.5 *If π_0 is (essentially)* Haag*-dual with respect to $\mathscr{D}(\mathscr{M}, g)$, then it's* Haag*-dual with respect to $\mathscr{W}(\mathscr{M}, g)$ and $\pi_0(\mathfrak{A}(\mathscr{W}_{p,\bar{q}}))'' = \pi_0(\mathfrak{A}(\mathscr{W}'_{q,\bar{p}}))'' = \pi_0(\mathfrak{A}(\mathscr{W}_{\bar{p},q})')'$ for all pairs $p, q \in \mathscr{I}$ such that $(p, \bar{q}) \in \mathscr{D}(\mathscr{I})$, where $\bar{p}$ is the* antipodal *of p (see formulae (1.27)–(1.28) and the discussion preceding them in page 12).*

> Proof. Essential Haag duality with respect to $\mathscr{D}(\mathscr{M}, g)$ is not only equivalent to Haag duality (for the collection of diamonds in $\mathscr{I}$ is closed under causal complements), but also implies Haag duality for wedges, for $\pi_0(\mathfrak{A}(\mathscr{D}_{p,\bar{q}}))'' = \pi_0(\mathfrak{A}(\mathscr{W}_{p,\bar{q}}))'' \subset \pi_0(\mathfrak{A}(\mathscr{W}'_{p,\bar{q}}))' \subset \pi_0(\mathfrak{A}(\mathscr{W}_{q,\bar{p}}))' = \pi_0(\mathfrak{A}(\mathscr{D}_{q,\bar{p}}))' = \pi_0(\mathfrak{A}(\mathscr{D}'_{p,\bar{q}}))'$. The last assertion follows from the identities $\pi_0(\mathfrak{A}(\mathscr{W}_{p,\bar{q}}))'' = \pi_0(\mathfrak{A}(\mathscr{D}_{p,\bar{q}}))'' = \pi_0(\mathfrak{A}(\mathscr{D}'_{q,\bar{p}}))'' = \pi_0(\mathfrak{A}(\mathscr{D}_{q,\bar{p}}))' = \pi_0(\mathfrak{A}(\mathscr{W}_{q,\bar{p}}))' = \pi_0(\mathfrak{A}(\mathscr{W}'_{q,\bar{p}}))''$. □

The last expression of Lemma 4.5 is trivial for AdS_d, but possesses potentially dramatic consequences for nontrivial AAdS spacetimes, for, in this case, $\pi_0(\mathfrak{A}(\mathscr{W}_{p,\bar{q}}))'' \cap \pi_0(\mathfrak{A}(\mathscr{W}_{\bar{p},q}))'' \supset \pi_0(\mathfrak{A}(\mathscr{W}'_{\bar{p},q} \cap \mathscr{W}'_{p,\bar{q}}))''$, and the region $\mathscr{W}'_{\bar{p},q} \cap \mathscr{W}'_{p,\bar{q}}$ is a nonvoid open set if $(\mathscr{M}, g)$ satisfies the hypotheses of Proposition 1.8 (page 20), stronger than those of Theorem 1.3 (see Figure 1.4, page 22). Supposing that the precosheaf of von Neumann

algebras $\mathscr{K}(\mathscr{M},g) \ni \mathscr{O} \mapsto \pi_0(\mathfrak{A}(\mathscr{O}))''$ satisfies the property of *extended locality*

$$\pi_0(\mathfrak{A}(\mathscr{O}_1))'' \cap \pi_0(\mathfrak{A}(\mathscr{O}_1))'' = \mathbb{C}\mathbb{1},\ \forall \mathscr{O}_1, \mathscr{O}_2 \in \mathscr{K}(\mathscr{M},g) : \mathscr{O}_1 \perp \mathscr{O}_2, \tag{4.20}$$

it then follows that $\pi_0(\mathfrak{A}(\mathscr{W}'_{\bar{p},q} \cap \mathscr{W}'_{p,\bar{q}}))'' = \mathbb{C}\mathbb{1}$ and, hence, $\pi_0(\mathfrak{A}(\mathscr{O}_{r,s}))'' = \mathbb{C}\mathbb{1}$ for any $r, s \in \mathscr{M}$ such that $\mathscr{O}_{p,q} \doteq I^+(r,\mathscr{M}) \cap I^-(s,\mathscr{M}) \subset \mathscr{W}'_{\bar{p},q} \cap \mathscr{W}'_{p,\bar{q}}$. As:

- Any sufficiently small diamond in $(\mathscr{M},g)$ satisfies the above condition for some pair $p, q \in \mathscr{I}$ satisfying the conditions of Lemma 4.5, and
- In Minkowski spacetime $\mathbb{R}^{1,d-1}$, any theory of local observables which is additive, translation covariant and locally causal in the GNS representation π_0 associated to a pure vacuum state (i.e., the joint spectrum of the translation generators is contained in $J^+(0,\mathbb{R}^{1,d-1})$ and the vacuum vector is an eigenvector of the generators with zero eigenvalue) satisfies extended locality [Lan69],

we're led to the following "*'no-go'* theorem":

Corollary 4.6 *Let $(\mathscr{M},g)$ be an AAdS spacetime satisfying the hypotheses of Proposition 1.8, $\mathfrak{A}$ an extended locally covariant quantum theory realized in $(\mathscr{M},g)$ and π_0 an irreducible *-representation of $\mathfrak{A}_{\mathscr{M}}$ satisfying the hypotheses of Lemma 4.5. If π_0 satisfies extended locality (4.20), then any* von Neumann *algebra localized in a sufficiently small open set consists only in multiples of $\mathbb{1}$. If the precosheaf of* von Neumann *algebras $\mathscr{K}(\mathscr{M},g) \ni \mathscr{O} \mapsto \pi_0(\mathfrak{A}(\mathscr{O}))''$ is moreover additive, then $\pi_0(\mathfrak{A}_{\mathscr{M}})'' = \mathbb{C}\mathbb{1}$.* □

Remark 4.13 *There is, to the author's knowledge, no proof of the property of extended locality in curved spacetimes under hypotheses similar to the ones employed in* Minkowski *spacetime (i.e., substituting, say, some version of the microlocal spectrum condition [BFK96] for the spectral condition of the vacuum). We conjecture, though, that such a demonstration is possible for real analytic AAdS spacetimes, along lines analogous to the proof of the timelike tube property of* Borchers *(page 81 and footnote 9).*

We conclude from Corollary 4.6 that essential Haag duality with respect to $\mathscr{D}(\mathscr{M},g)$ implies potentially a trivial quantum theory if combined with bulk additivity, and even a non additive bulk quantum theory may end up being trivial at sufficiently small scales – in particular, the procedure of holographic reconstruction of the realization of $\mathfrak{A}$ in $(\mathscr{M},g)$ from the dual theory in the boundary, whose geometrical part was schematized in Subsubsection 1.3.1.3 (page 23), is *useless* in both cases! [12] The following hypothesis, though, does lead to nontrivial results:

[12] On the other hand, the possibility that π_0 is Haag-dual and $\pi_0(\mathfrak{A}(\mathscr{W}'_{p,\bar{q}} \cap \mathscr{W}'_{q,\bar{p}}))'' = \mathbb{C}\mathbb{1}$ is not completely uninteresting from the physical viewpoint – as wedges model, in a certain sense, the exterior of black holes, we can imagine the region $\mathscr{W}'_{p,\bar{q}} \cap \mathscr{W}'_{q,\bar{p}}$ as being "inside the horizon". From this viewpoint,

(v) π_0 is *essentially* HAAG-dual with respect to $\mathscr{W}(\mathscr{M}, g)$, but it's *not* (essentially) HAAG-dual with respect to $\mathscr{D}(\mathscr{M}, g)$.

The second premise allows us to escape from Corollary 4.6. In particular, in this case the precosheaf of VON NEUMANN algebras $\mathscr{D}(\mathscr{M}, g) \ni \mathscr{D}_{p,\bar{q}} \mapsto \pi_0(\mathfrak{A}(\mathscr{D}_{q,\bar{p}}))'$ is not locally causal, but the precosheaf

$$\mathscr{W}(\mathscr{M}, g) \ni \mathscr{W}_{p,q} \mapsto \mathfrak{B}(\mathscr{W}_{p,q}) \doteq \pi_0(\mathfrak{A}(\mathscr{W}'_{p,q}))' \tag{4.21}$$

is, by hypothesis (v).

LEMMA 4.7 *Suppose that π_0 satisfies hypothesis (v). Then, π_0 satisfies holographic local definiteness if and only if the quasilocal algebras $\mathfrak{B}''_{\mathscr{P}oi(r)}$ are* irreducible, *i.e. $\mathfrak{B}'_{\mathscr{P}oi(r)} = \mathbb{C}\mathbb{1}$, for all $r \in \mathscr{I}$. In particular, in this case $\mathfrak{B}''_{\mathscr{M}}$ is irreducible.*

PROOF. It follows immediately from formula

$$\bigcap_{\substack{(p,\bar{q})\in\mathscr{D}(\mathscr{I}),\\ \mathscr{W}_{p,\bar{q}}\ni\bar{r}}} \pi_0(\mathfrak{A}(\mathscr{W}_{p,\bar{q}}))'' = \bigcap_{\substack{(p,\bar{q})\in\mathscr{D}(\mathscr{I}),\\ \mathscr{W}_{p,\bar{q}}\ni\bar{r}}} \pi_0(\mathfrak{A}(\mathscr{W}'_{q,\bar{p}}))'' =$$

$$= \left(\bigcup_{q\ll_{\mathscr{I}}\bar{p}\in\mathscr{M}in(r)} \mathfrak{B}(\mathscr{W}_{q,\bar{p}}) \right)',$$

and the fact that

$$\bigcap_{\substack{(p,\bar{q})\in\mathscr{D}(\mathscr{I}),\\ \mathscr{W}_{p,\bar{q}}\ni\bar{r}}} \mathscr{W}'_{q,\bar{p}} = \bigcap_{\substack{(p,\bar{q})\in\mathscr{D}(\mathscr{I}),\\ \mathscr{W}_{p,\bar{q}}\ni\bar{r}}} \mathscr{W}_{p,\bar{q}} = \{\bar{r}\},$$

by virtue of Proposition 1.8 and Theorem 1.3. □

In particular, internal symmetries of the precosheaf $\mathscr{W}_{p,q} \mapsto \mathfrak{B}(\mathscr{W}_{p,q})$ cannot be spontaneously broken in $\mathscr{H}_0$ if π_0 satisfies holographic local definiteness and hypothesis (v). Notice that this *doesn't* imply that the internal symmetries of the precosheaf $\mathscr{W}_{p,q} \mapsto \pi_0(\mathfrak{A}(\mathscr{W}_{p,q}))''$ are not broken.

Let us consider, now, a *-representation π of $\mathfrak{A}_{\mathscr{M}}$ in the HILBERT space $\mathscr{H}_\pi$ satisfying

$$\forall p, q \in \mathscr{D}(\mathscr{I}), \exists V \in \mathscr{B}(\mathscr{H}_0, \mathscr{H}_\pi) \text{ unitary such that } \pi(A) = V\pi_0(A)V^*, \forall A \in \mathfrak{A}(\mathscr{W}'_{p,q}). \tag{4.22}$$

we see that a quantum theory with the characteristics above is unable to "see" physical events localized "inside the event horizon". As HAAG duality with respect to $\mathscr{D}(\mathscr{M}, g)$ is necessary so that the REHREN-dual quantum theory in the boundary be conformally covariant and π_0 be the GNS representation associated to a vacuum state [BGL93], we conjecture that such a (non additive) quantum theory probably doesn't suffer from the *black hole information paradox*, but is also essentially insensitive to the details of the bulk geometry, in the same way as the scaling limit built in the previous Section.

In particular, π satisfies the DHR criterion with respect to $\mathscr{D}(\mathscr{M},g)$ under the Rehren bijection. The fact that π_0 doesn't satisfy essential Haag duality with respect to $\mathscr{D}(\mathscr{M},g)$ guarantees that the *-representations satisfying (4.22) constitute a *proper* subclass of DHR excitations.

One should notice that, even from the original viewpoint of the original form of Rehren duality, i.e., $(\mathscr{M},g)=AdS_d$, it looks weird at first sight to employ a selection criterion which aims at modelling vacuum excitations corresponding to particles, since conformal quantum field theories are not elementary particle theories. However, it was demonstrated by Buchholz, Mack and Todorov [BMT88] that:

Theorem 4.8 ([BMT88]) *If π_0 is the GNS representation of a conformally invariant vacuum state ω_0 (which is, hence,* Haag*-dual with respect to $\mathscr{D}(AdS_d)$ [BGL93]), any state ω with GNS representation π_ω with positive energy is a DHR excitation of ω_0, i.e., given any $(p,q)\in\mathscr{D}(\mathscr{I})$, we have $\pi_\omega{\restriction}_{\mathfrak{A}(\mathscr{D}'_{p,q})}$ is unitarily equivalent to $\pi_0{\restriction}_{\mathfrak{A}(\mathscr{D}'_{p,q})}$.*

> Proof. Consider the following facts: (i) $(p,\bar{q})\in\mathscr{D}(\mathscr{I})$ if and only if $(q,\bar{p})\in\mathscr{D}(\mathscr{I})$; (ii) $\omega_0{\restriction}_{\pi_0(\mathfrak{A}(\mathscr{D}_{p,q}))''}$ is normal and faithful for all $(p,q)\in\mathscr{D}(\mathscr{I})$ (Reeh-Schlieder property). As the Reeh-Schlieder property also holds if we substitute π for π_0 by virtue of energy-momentum positivity, it follows that the map $\pi_0(A)\mapsto\pi(A)$ for $A\in\mathfrak{A}(\mathscr{D}_{p,q})$ extends to a normal *-isomorphism between $\pi_0(\mathfrak{A}(\mathscr{D}_{p,q}))''$ and $\pi_\omega(\mathfrak{A}(\mathscr{D}_{p,q}))''$. As the cyclic vectors Ω_0 and Ω in the respective GNS Hilbert spaces $\mathscr{H}_0$ and $\mathscr{H}_\omega$ associated to ω_0 and ω are also separating by the Reeh-Schlieder property, it follows from Theorem 2.5.32 in [BR87] (see also Theorem 7.2.9 in [KR86]) that such a *-isomorphism is implemented by a unitary operator between the GNS Hilbert spaces of ω_0 and ω. From (i), we finally see that π satisfies the DHR criterion, with ρ localized in the diamond $\mathscr{D}'_{p,q}$. □

As the Reeh-Schlieder property is valid for the reference representations π_0 we've adopted for AAdS spacetimes, we see that such a criterion is sufficiently general for our purposes. The endomorphism ρ corresponding to π and localized, say, in $\mathscr{D}_{p,\bar{q}}$ has the following properties:

1. By construction, $\rho\circ\pi_0(A)=\pi_0(A)$ for all $A\in\mathfrak{A}(\mathscr{D}_{q,\bar{p}})$;
2. If $(p',q')\in\mathscr{D}(\mathscr{I})$ are such that $p,\bar{q},p',q'\in\mathscr{M}in(r)$ for some $r\in\mathscr{I}$ and $B\in\mathfrak{A}(\mathscr{D}_{p',q'})$, then $\rho\circ\pi_0(B)\in\mathfrak{B}(\mathscr{D}_{p'',q''})$ for $(p'',q'')\in\mathscr{D}(\mathscr{I})$ such that $\mathscr{D}_{p,\bar{q}},\mathscr{D}_{p',q'}\subset\mathscr{D}_{p'',q''}$ (repeat the argument used in Subsection 4.4.1, page 86).

The second property above leads to the following scenario: $\rho\circ\pi_0$ only assumes values in $\pi_0(\mathfrak{A}_{\mathscr{M}})$ for $A\in\mathfrak{A}\left(\bigcup_{(p',q')\in\mathscr{D}(\mathscr{I}):\mathscr{W}_{p',q'}\supset\mathscr{W}_{p,\bar{q}}}\mathscr{W}_{p',q'}\right)$. However, if ρ' is the endomorphism

associated to π and localized in $\mathscr{W}_{q,\bar{p}}$, we have that ρ' can still be obtained from the transport of ρ by means of local intertwiners, for, although there is no $(p',q') \in \mathscr{D}(\mathscr{I})$ such that $\mathscr{W}_{p',q'} \supset \mathscr{W}_{p,\bar{q}} \cup \mathscr{W}_{q,\bar{p}}$, ρ can always be transported to a smaller wedge contained in $\mathscr{W}_{q,\bar{p}}$.

We can imagine, in view of the possibility of spontaneous breaking of internal symmetries allowed by essential Haag duality, that π_0 is *not* irreducible. More precisely, suppose that the (trivially) localized endomorphism $\iota \doteq \mathrm{id}_{\pi_0(\mathfrak{A}_{\mathscr{M}})}$ associated to π_0 has nontrivial subobjects, i.e., for each $(p,\bar{q}) \in \mathscr{D}(\mathscr{I})$ there is a projection $\mathbb{1} \neq E \in \pi_0(\mathfrak{A}(\mathscr{W}_{p,\bar{q}}))'$, a *-endomorphism σ of $\pi_0(\mathfrak{A}_{\mathscr{M}})$ localized in $\mathscr{W}_{p,\bar{q}}$ and an isometry $W \in (\sigma,\iota)$ such that $WW^* = E$ (by the property B of Borchers, $W \in \pi_0(\mathfrak{A}(\mathscr{W}_{p,\bar{q}}))'$). Actually, we only need the existence of E (guaranteed, on its turn, by the non irreducibility of π_0), for the existence of W and, thus, of σ ($\sigma(\pi_0(A)) \doteq W^*\pi_0(A)W$) follows from the property B. Hence, we can also consider the isometry V and the localized endomorphism σ' associated to the projection $\mathbb{1} - E$. $\sigma \circ \pi_0$ as well as $\sigma' \circ \pi_0$ satisfy essential Haag duality with respect to $\mathscr{W}(\mathscr{M}, g)$.

Definition 4.14 *If σ and σ' are subobjects of ι which generate* disjoint *representations, their unitary equivalence classes are then called* phases *of the reference state ω_0. A phase σ is said to be* pure *if the corresponding representation is irreducible.*

Let us take the subobjects σ and σ' of ι constructed in the previous paragraph, localized respectively, say, in $\mathscr{D}_{p,\bar{q}}$ and $\mathscr{D}_{q,\bar{p}}$. As in this case $\sigma(\pi_0(A)) = \sigma'(\pi_0(A)) = \pi_0(A)$ if $A \in \mathfrak{A}(\mathscr{W}'_{p,\bar{q}}) \cap \mathfrak{A}(\mathscr{W}'_{q,\bar{p}}) \supset \mathfrak{A}(\mathscr{W}'_{p,\bar{q}} \cap \mathscr{W}'_{q,\bar{p}})$, we can consider then the following *-endomorphism τ of $\pi_0(\mathfrak{A}_{\mathscr{M}})''$, which satisfies the following properties:

(S1) $\tau \circ \sigma(\pi_0(A)) = \sigma(\pi_0(A))$ for all $A \in \mathfrak{A}(\mathscr{W}_{p,\bar{q}})$;

(S2) $\tau \circ \sigma'(\pi_0(A)) = \sigma'(\pi_0(A))$ for all $A \in \mathfrak{A}(\mathscr{W}_{q,\bar{p}})$.

Obviously, we have $\tau \circ \sigma(\pi_0(A)) = \tau \circ \sigma'(\pi_0(A)) = \tau(\pi_0(A))$ if $A \in \mathfrak{A}(\mathscr{W}'_{p,\bar{q}} \cap \mathscr{W}'_{q,\bar{p}})$. We can, thus, say that τ is "localized" in $\mathscr{W}'_{p,\bar{q}} \cap \mathscr{W}'_{q,\bar{p}}$ and interpolates the phases σ and σ' (recall that, in AAdS spacetimes satisfying, as we've assumed in this Section, the hypotheses of Proposition 1.8 and, hence, of Theorem 1.3, $\mathscr{W}'_{p,\bar{q}} \cap \mathscr{W}'_{q,\bar{p}} \neq \varnothing$! See Figure 1.4, page 1.4).

Definition 4.15 *An endomorphism τ which satisfies (S1) and (S2) for some pair of phases σ, σ' of ω_0 localized respectively in wedges $\mathscr{W}_{p,\bar{q}}$, $\mathscr{W}_{q,\bar{p}}$ is said to be* solitonic. *Then, we say that τ* separates *or* interpolates *σ and σ', and the unitary equivalence class of τ is a* soliton.

Notice that an intertwiner which transports σ from the wedge $\mathscr{W}_{p,\bar{q}}$ to another, sufficiently close wedge $\mathscr{W}_{p',\bar{q}'}$, when applied to σ', it transports the latter from $\mathscr{W}_{q,\bar{p}}$ to $\mathscr{W}_{q',\bar{p}'}$ and, hence, when applied to τ, produces a solitonic endomorphism "localized" in $\mathscr{W}'_{p',\bar{q}'} \cap \mathscr{W}'_{q',\bar{p}'}$ in the sense above. Thus, we're entitled to remove the quotation marks and say that τ is, indeed, *localized* in $\mathscr{W}'_{p,\bar{q}} \cap \mathscr{W}'_{q,\bar{p}}$. The composition of solitonic endomorphisms τ, τ' is not always defined, for it demands that the corresponding endomorphisms representing the interpolated phases τ and τ' "fit together".

4.4.3 Obstructions to the inversion of Rehren duality

It remains to discuss the possibility that, starting from a precosheaf $\mathscr{D}_{p,q} \mapsto \mathfrak{A}(\mathscr{D}_{p,q})$ realizing a locally covariant quantum theory $\mathfrak{A}$ in $(\mathscr{I}, \bar{g}^{(0)})$, we may construct for each AAdS spacetime $(\mathscr{M}, g)$ satisfying the hypotheses of the present Section a precosheaf $\mathscr{O} \mapsto \mathfrak{A}(\mathscr{O})$ indexed by open, globally hyperbolic and causally convex regions $\mathscr{O} \subset \mathscr{M}$, and not only wedges (whose local algebras are obtained by defining $\mathfrak{A}(\mathscr{W}_{p,q}) \doteq \mathfrak{A} \circ \rho_{(\mathscr{M},g)}(\mathscr{W}_{p,q})$).

Let us take a reference state ω in the realization of $\mathfrak{A}$ in $(\mathscr{M}, g)$. If the precosheaf of von Neumann algebras $\mathscr{O} \mapsto \pi_\omega(\mathfrak{A}(\mathscr{O}))''$ is *additive*, we can try to perform this reconstruction by employing the results of Subsubsection 1.3.1.3 (page 23) if we define, for a relatively compact and geodesically convex $\mathscr{O}_{p,q} \subset \mathscr{M}$ sufficiently small,

$$\mathfrak{B}(\mathscr{O}_{p,q}) \doteq \bigcap_{\mathscr{W}_{p,q} \supset \mathscr{O}_{p,q}} \pi_\omega(\mathfrak{A}(\mathscr{W}_{p,q}))''. \tag{4.23}$$

Proposition 4.9 *Suppose that the precosheaf of* von Neumann *algebras* $\mathscr{O} \mapsto \pi_\omega(\mathfrak{A}(\mathscr{O}))''$ *is additive and satisfies* Haag *duality with respect to wedges. Then* $\mathfrak{B}(\mathscr{O}_{p,q}) = \pi_\omega(\mathfrak{A}(\mathscr{O}'_{p,q}))'$.

Proof. Notice that the operation which takes a von Neumann algebra to its commutant satisfies the same properties that of the operation of causal complement in causally complete regions (see Appendix A, end of Subsection A.2.1, page 117), if we substitute the algebra generated by the union for the union of algebras. Then it follows from (4.23) that $\mathfrak{B}(\mathscr{O}_{p,q}) \doteq \bigcap_{\mathscr{W}_{p,q} \supset \mathscr{O}_{p,q}} \pi_\omega(\mathfrak{A}(\mathscr{W}'_{p,q}))' = \left(\bigvee_{\mathscr{W}_{p,q} \supset \mathscr{O}_{p,q}} \pi_\omega(\mathfrak{A}(\mathscr{W}_{p,q})')''\right)' = \pi_\omega\left(\mathfrak{A}\left(\bigcup_{\mathscr{W}_{p,q} \supset \mathscr{O}_{p,q}} \mathscr{W}'_{p,q}\right)\right)' = \pi_\omega(\mathfrak{A}(\mathscr{O}'_{p,q}))'$. □

If the precosheaf $\mathscr{O} \mapsto \pi_\omega(\mathfrak{A}(\mathscr{O}))''$ is Haag-dual, then it's also additive and, thus, the reconstruction procedure based upon (4.23) can be taken to completion in a successful way. If we assume, on the other hand, only *essential* Haag duality, the Haag-dual precosheaf $\mathscr{O} \mapsto \pi_\omega(\mathfrak{A}(\mathscr{O}'))'$ *doesn't need* to be additive. Having in sight the scenario of spontaneous breaking of internal symmetries, we can give a natural physical interpretation for this phenomenon: the Haag-dual net circumvents the spontaneous breaking of

internal symmetries by including precisely those observables corresponding to extended physical procedures, localized, say, around domain walls, just like the soliton sectors built in the previous Subsection. Such a situation is analogous to ordered phases in quantum spin models in Statistical Mechanics, where KRAMERS-WANNIER duality produces a gas of "contours around pure-phase domains", which, on its turn, finds itself in the symmetric (disordered) phase [Kad00, Min00]. If we imagine the region $\mathscr{W}'_{p,\bar{q}} \cap \mathscr{W}'_{q,\bar{p}}$ ($\neq \varnothing$!), sketched in Figure 1.4 (page 1.4) as the rough localization of a *domain wall*, i.e., a maximal (that is, which isn't properly included into any other submanifold of the same codimension), acausal codimension-two submanifolds of $\mathscr{M}$, it's quite tempting to interpret the solitons in Definition 4.15 in the light of the AdS/CFT correspondence as *D-branes*, for, as we've seen, the former share essentially the same properties of the objects of the same name in string theory [Zwi04].

We moreover notice that gravitationally induced superselection sectors have already been previously considered in the literature, in particular related to the superselection rule for electric charge [AS80, Sor79]. Such sectors enjoy a rather large amount of universality, and are induced even in free quantum field theories.

We emphasize that the soliton sectors we've built *don't exist* in AdS_d, for in this case HAAG duality and essential HAAG duality with respect to $\mathscr{W}((\mathscr{M}, g) = AdS_d)$, as we've seen, coincide (this, of course, is directly related to the fact that $\mathscr{W}'_{p,\bar{q}} \cap \mathscr{W}'_{q,\bar{p}} = \varnothing$ in AdS_d), and the direct sum of reference *-representations satisfying HAAG duality doesn't necessarily, but the direct sum of*essentially* HAAG-dual reference *-representations is essentially HAAG-dual [Rob04]. More in general, HAAG duality with respect to wedges leads, by Lemma 4.7, to the irreducibility of π_0, making impossible the existence of soliton sectors even if we manage to escape Corollary 4.6 and have nontrivial observables in $\mathscr{O} \subset \mathscr{M}$ arbitrarily small for $(\mathscr{M}, g) \neq AdS_d$ (in AdS_d, there is no problem in assuming HAAG duality, and in this case the original REHREN duality allows a one-to-one correspondence). Hence, the presence of soliton sectors seems to essentially render useless or impossible (modulo the validity of the property of extended locality) the unequivocal reconstruction of the bulk quantum theory by means of the prescription (4.23) in nontrivial AAdS geometries. We speculate that an alternative, more indirect possibility, for at least a partial reconstruction, albeit beyond the mere specification of the localization of observables in wedges by the REHREN bijection, would be to elaborate an operator product expansion around the scaling limit, as suggested at the end of Section 4.3.

— ¿Cómo puede ser eso? — respondió don Quijote —. ¿Tan de esencia de la historia es saber las cabras que han pasado por estenso, que si se yerra una del número no puedes seguir adelante con la historia?

— No, señor, en ninguna manera — respondió Sancho —; porque así como yo pregunté a vuestra merced que me dijeste cuántas cabras habían pasado, y me respondió que no sabía, en aquel mesmo instante se me fue a mí de la memoria cuanto que quedaba por decir, y a fe que era de mucha virtud y contiento.

— ¿De modo — dijo don Quijote — que ya la historia es acabada?

— Tan acabada es como mi madre — dijo Sancho.

— Dígote de verdad — respondió don Quijote — que tú has contado unas de las más nuevas consejas, cuento o historia que nadie pudo pensar en el mundo, y que tal modo de contarla ni dejarla jamás se podrá ver ni habrá visto en toda la vida, aunque no esperaba yo otra cosa de tu buen discurso; mas no me maravillo, ques quizá estos golpes que no cesan te deben de tener turbado el entendimiento.[xii]

Miguel de Cervantes Saavedra
El ingenioso hidalgo don Quijote de la Mancha, Primero Libro, Cap. XX

Part III

A synthesis

Coda. Concluding remarks

Go, go, go, said the bird: human kind
Cannot bear very much reality.
Time past and time future
What might have been and what has been
Point to one end, which is always present.

T. S. Eliot
"Burnt Norton" (*Four Quartets*, 1943), I

The present work was able to reveal quite a rich structure underlying the AdS/CFT correspondence, of which we've been able only to scratch the surface.

The main contributions of the present work, as a whole, can be listed as follows:

- One has specified the essential aspects underlying Rehren duality. In particular, the geometrical and causal character of the change of localization proposed by Rehren was outlined from the global viewpoint (Chapter 1) as well as the local one (i.e., in a neighbourhood of $\mathscr{I}$ – Chapter 2).

- The elaborate machinery of global Lorentzian geometry developed from two different viewpoints – kinematic (causal structure, conformal infinity) and dynamical (time evolution, return to equilibrium) – allowed one to determine sufficiently robust conditions on the geometry of AAdS spacetimes so as certain basic properties of Rehren duality are preserved and others are modified in a relatively clear way, as seen in Theorem 1.3 (page 16), in Proposition 1.8 (page 20) and in Theorem 1.14 (page 27).

- We proposed a geometrically intrinsic manner of constructing global time functions associated to relatively compact diamonds in causally simple spacetimes and AAdS wedges, which admits a formulation of the (asymptotic) zeroth and second laws of black hole dynamics (Section 2.3, page 40ff.).

- At the quantum level, we implemented an instance of the principle of local covariance, proposed by Brunetti, Fredenhagen and Verch (Chapter 3), in a situation which demands the imposition of boundary conditions (see Definition 4.1, page 70, and the discussion that follows it) adapted to the geometry of (A)AdS spacetimes, with the help of the algebraic formalism developed by Sommer and the prototype case of AdS_d (Proposition 4.4, page 77).

- We proposed a formalism of "scaling algebras" around points at the conformal infinity of AAdS spacetimes which, aided by the powerful structural results of Borchers and Yngvason (Theorems (4.1) to (4.3), page 75ff.) and by the geometrical viewpoint of Section 2.3, not only permitted a precise determination of boundary conditions for physically relevant states, but also established a close relation between return to thermal equilibrium by the action of asymptotic isometries and scaling limits (condition (d), page 84 and the discussion that follows).

- We pointed out how soliton excitations similar to D-branes (Definition 4.15, page 93) may arise in the case of nontrivial AAdS geometries, relating them to a nontrivial "vacuum" structure (Definition 4.14, page 93) and to a potential impossibility of reconstructing the bulk quantum theory from its Rehren dual by means of the prescription (4.23) (page 94).

A problem of great interest in the literature on Lorentzian geometry concerns the stability and rigidity of AAdS spacetimes given conditions similar to the ones assumed in Theorem 1.3 and in Proposition 1.8. An important partial result was recently obtained by Anderson [And06b], which tells us that, given (i) a global time function t in an asymptotically simple, geodesically complete AAdS spacetime $(\mathscr{M}, g)$ satisfying a technical condition of unique continuation of solutions of the linearized Einstein equations across conformal infinity, (ii) a sequence of Cauchy data defined at distinct levels $t^{-1}(t_i)$ of t, such that $t_i \overset{i\to\infty}{\longrightarrow} +\infty$, and tending to *stationary* Cauchy data, then $(\mathscr{M}, g)$ is *globally* stationary. Although this is not a strict rigidity result, it illustrates how the boundary conditions at conformal infinity act so as to prevent the dispersion of perturbations of Cauchy data. We are, however, far from a result about the global nonlinear stability of (A)AdS spacetimes with the same precision as the one due to Christodoulou and Klainerman [CK93] for Minkowski spacetime (see also the Remark 2.2, page 51, for more considerations about this issue). Another open question is the feasibility of demonstrating the rigidity of AdS spacetimes under (absence of) gravitational time delay of null geodesics [PSW02] (see Figure 1.3, page 17). Such questions interest us to the extent that answering them would tell us how natural are our geometrical assumptions from the gravitational viewpoint.

The construction of global time functions for relatively compact diamonds we've obtained can, we believe, be further deepened, in the direction of asymptotic analogs of the other laws of black hole dynamics, thus giving us details on the "fluctuations" of geometrical quantities around the scaling limit at the tips of the diamonds, hence seeking an analog (of a generalization) of the first law of black hole dynamics for diamonds. From the viewpoint of AAdS wedges, it interests us to relate some quantity of this kind with the rescaled electric part E_{ab} of the WEYL tensor, which enters as part of the "initial data" for the FEFFERMAN-GRAHAM expansion of the metric around infinity.

The most glaring limitation of the present work is the total absence of nontrivial examples, concerning geometry as well as (even free) QFT models. The geometrical issue, as seen above, involves finding nontrivial examples of nonsingular AAdS spacetimes, which is very difficult without a general result as in the case of zero cosmological constant. Regarding the quantum part, a future, quite compelling direction of investigation would be to check the boundary condition (4.17), at least for free fields, in terms of more familiar premises. Doing this for general spacetimes involves plunging into the arsenal of microlocal analysis [Hör71, DH72, Dui96, Hör90] in order to build in a precise way the GREEN functions.

The construction of an *operator product expansion* within our extension of REHREN duality represents a challenge of its own, albeit of the highest interest, for it might potentially reveal relations between deviations from thermal equilibrium in AAdS wedges and geometrical quantities such as E_{ab}. An example which motivates such a conjecture is the relation between the *poles* of retarded two-point functions of the dual quantum theory at the boundary, which determine the properties of return to equilibrium, and *quasinormal modes* of AAdS black holes [HH00, SS02]. Such a relation should have an analog for wedges and free fields in AAdS.

Last but not least, we obviously presented in Subsection 4.4.2 (page 89ff.) only the bare bones of the properties of soliton sectors of the bulk quantum theory associated to pairs of phases (satisfying the DHR criterion) of its REHREN dual at the boundary. The results presented above certainly deserve a deeper scrutiny, along the lines of the works of MÜGER [Müg98, Müg96] and SCHLINGEMANN [Sch96] in the general case, and along the lines of the works of ASHTEKAR and SEN [AS80] and SORKIN [Sor79] for the construction of examples, combining the techniques of these papers with the aforementioned tools of microlocal analysis.

Appendices

Appendix A

Elements of Lorentzian geometry

We shall present here the necessary definitions and the statement of some useful results in Lorentzian geometry. Our references are [BEE96, HE73, O'N83, Wal84].

A.1 Local theory

Let $\mathscr{M}$ be a d-dimensional manifold, $d \geq 2$. A *semi-Riemannian* metric of index p is a section g of $T^*\mathscr{M} \otimes T^*\mathscr{M}$ such that $g(r)$ is a nondegenerate symmetric bilinear form of index p, $\forall r \in \mathscr{M}$. The pair $(\mathscr{M}, g)$ is said to be a *semi-Riemannian manifold* – for $p = 0$, $(\mathscr{M}, g)$ is simply a Riemannian manifold. If $p = 1$, we say that $(\mathscr{M}, g)$ is a *Lorentzian manifold*.

For all cases, we'll denote by ∇_a the Levi-Civita connection associated to g – recall that a (Koszul) *connection* in a vector bundle $\mathscr{E} \xrightarrow{p} \mathscr{M}$ is a map $\nabla : \Gamma^\infty(\mathscr{M}, T\mathscr{M}) \times \Gamma^\infty(\mathscr{M}, \mathscr{E}) \to \Gamma^\infty(\mathscr{M}, \mathscr{E})$, $\mathscr{C}^\infty(\mathscr{M})$-linear in the first variable and, for all $X \in \Gamma^\infty(\mathscr{M}, T\mathscr{M})$, $\nabla_X \cdot$ is a *derivation* on the $\mathscr{C}^\infty(\mathscr{M})$-module $\Gamma^\infty(\mathscr{M}, \mathscr{E})$, i.e.,

$$\nabla_X(fS) = \langle df, X\rangle S + f\nabla_X S, \forall S. \tag{A.1}$$

Due to (A.1), the difference between two connections ∇ and $\tilde{\nabla}$ is a $\mathscr{C}^\infty(\mathscr{M})$-bilinear map with values in $\Gamma^\infty(\mathscr{M}, \mathscr{E})$, i.e., is a section of $T\mathscr{M} \otimes \mathscr{E}^* \otimes \mathscr{E}$. A connection ∇ functorially induces a unique connection in all tensor bundles $\mathscr{E}^{(r,s)} = (\otimes^r \mathscr{E}) \otimes (\otimes^s \mathscr{E}^*)$ ($\mathscr{E}*$ is the *dual* bundle to $\mathscr{E}$) by means of the Leibniz rule:

$$\nabla_X(S_1 \otimes S_2) = (\nabla_X S_1) \otimes S_2 + S_1 \otimes (\nabla_X S_2),\ (\nabla_X S')(S) + S'(\nabla_X S) = \langle dS'(S), X\rangle.$$

In the case $\mathscr{E} = T\mathscr{M}$, we say that ∇ is *symmetric* is the *torsion* tensor $T(X, Y) \doteq \nabla_X Y - \nabla_Y X - [X, Y]$ vanishes for all X, Y ($[X, Y] = \pounds_X Y$ denotes the Lie bracket of X with Y). In a semi-Riemannian manifold $(\mathscr{M}, g)$, a Levi-Civita connection is the

(unique!) symmetrical connection ∇ in $T\mathscr{M}$ which satisfies $\nabla_X g = 0$ for all X. We use, for the latter, the notation ∇_a to denote its tensor character.

The Riemann curvature tensor $\mathrm{Riem}(g)$ associated to g (more precisely, associated to ∇_a) is given by $2\nabla_{[a}\nabla_{b]}X_c = \mathrm{Riem}(g)^d_{abc}X_d$, for all X_a. We denote yet the Ricci tensor and the scalar curvature associated to g respectively by $\mathrm{Ric}(g)_{ab} \doteq \mathrm{Riem}(g)^c_{acb}$ and $R(g) = g^{ab}\mathrm{Ric}(g)_{ab}$, where $g^{ab} \doteq (g^{-1})^{ab}$, $g^{ab}g_{bc} = \delta^a_c$. The Weyl tensor $C(g)$ associated to g is given, for $d \geq 3$, by

$$C_{abcd} = \mathrm{Riem}_{abcd} - \frac{2}{d-2}(g_{a[c}\mathrm{Ric}_{d]b} - g_{b[c}\mathrm{Ric}_{a]d}) + \frac{2}{(d-1)(d-2)}R_{a[c}g_{d]b}. \tag{A.2}$$

This expression algebraically vanishes for $d \leq 3$, and, in this case, $\mathrm{Riem}(g)$ is completely determined by $\mathrm{Ric}(g)$, through formula (A.2). We adopt in this formula, as well as in several places throughout the present work, a slight abuse of notation by omitting the specification of g for the Weyl tensor. We'll do it whenever such a practice doesn't cause confusion, to alleviate the notation.

Given a semi-Riemannian manifold $(\mathscr{M}, g)$, $p \in \mathscr{M}$ we say that $X(p) \in T_p\mathscr{M}$ is *timelike*, *spacelike*, *causal* or *lightlike* (*null*) if, respectively, $g(X,X)(p) < 0$, > 0, ≤ 0 or $= 0$. A vector field X is timelike, spacelike, causal or lightlike / null if $X(p)$ is for all $p \in \mathscr{M}$. We define analogously timelike, spacelike, causal and lightlike / null covectors and covector fields by exchanging g with g^{-1}. Such a characteristic of a (co)vector field is said to be its *causal character*. The raising and lowering of indices by g, g^{-1} doesn't modify the causal character.

A piecewise $\mathscr{C}^\infty$ curve γ is said to be *timelike, spacelike, causal* or *null* if, in each $\mathscr{C}^\infty$ component γ_α, $\alpha = 1, \ldots, k$, the tangent vector $\dot\gamma^a$ satisfies respectively $\dot\gamma^a_\alpha\dot\gamma_{\alpha a} < 0$, $\dot\gamma^a_\alpha\dot\gamma_{\alpha a} > 0$, $\dot\gamma^a_\alpha\dot\gamma_{\alpha a} \geq 0$ or $\dot\gamma^a_\alpha\dot\gamma_{\alpha a} = 0$. A causal γ is *past* or *future* directed if, for t^a denoting a timelike vector field which determines the time orientation of $(\mathscr{M}, g_{ab})$, we have respectively $t^a\dot\gamma_{\alpha a} > 0$ or $t^a\dot\gamma_{\alpha a} < 0$ along each component γ_α.

A $\mathscr{C}^\infty$ curve γ is a *geodesic* if its tangent vector is covariantly constant: $\dot\gamma^a\nabla_a\dot\gamma_b = 0$. Thus, a geodesic segment is uniquely determined by the choice of a point p of $\mathscr{M}$ and a tangent vector at p, by the theorem of existence and uniqueness of solutions of ordinary differential equations on manifolds. Given $\mathscr{O}_1 \subset \mathscr{O} \subset \mathscr{M}$, we say that $\mathscr{O}_1$ is *geodesically convex* with respect to $\mathscr{O}$ if, given any $p, q \in \mathscr{O}_1$, there exists a unique geodesic segment contained in $\mathscr{O}$ linking p to q. Any $p \in \mathscr{M}$ has a geodesically convex neighbourhood. More in general, a neighbourhood $\mathscr{U}$ of $p \in \mathscr{M}$ is said to be *geodesically normal* if for all $q \in \mathscr{U}$ there exists a unique geodesic linking p to q (i.e., we don't assume that this is true for $q, q' \neq p$). For such neighbourhoods, we can write the following system of coordinates: the *exponential map*

$$\begin{aligned} &\exp_p : \mathscr{V} \subset T_p\mathscr{M} \to \mathscr{U} \\ &X \mapsto \exp_p(X) = \gamma_X(1), \end{aligned} \tag{A.3}$$

where $\gamma_X : [0,1] \to \mathscr{M}$ is the (unique) geodesic segment such that $\gamma(0) = p$ and $\dot{\gamma}(0) = X$, is a diffeomorphism for $\mathscr{U}$ sufficiently small. In this case, $\exp_p^{-1}(\mathscr{U}) = \mathscr{V}$ is a *star-shaped* neighbourhood of the origin in $T_p\mathscr{M} \cong \mathbb{R}^d$ (i.e., given $q \in \mathscr{V}$, we have $tq \in \mathscr{V}$ for all $t \in [0,1]$), whose Cartesian coordinates $(\exp_p^{-1})^\mu \doteq x^\mu : \mathscr{U} \to \mathscr{V}$ are denominated *(geodesic) normal* coordinates. The metric in $\mathscr{U}$, expressed by means of the latter, becomes [BGM71, Spi79]

$$g_{\mu\nu}(p) = (g \circ x^{-1})_{\mu\nu}(0) = \eta_{\mu\nu},\ \partial_\mu(g \circ x^{-1})(0) = 0; \tag{A.4}$$

$$g_{\mu\nu}(\exp_p(x)) = \eta_{\mu\nu} - \frac{1}{3}\mathrm{Riem}(g \circ x^{-1})(0)_{\mu\rho\nu\sigma}x^\rho x^\sigma + o(x^2); \tag{A.5}$$

$$\Downarrow$$

$$\sqrt{|\det(g \circ x^{-1})|}(\exp_p(x)) = 1 - \frac{1}{6}\mathrm{Ric}(g \circ x^{-1})(0)_{\rho\sigma}x^\rho x^\sigma + o(x^2). \tag{A.6}$$

The formulae (A.4)–(A.5), already present in embryonic form in the celebrated *Habilitationsschrift* of B. Riemann, were proven in complete generality by É. Cartan.

In the case of Lorentzian manifolds, we say that a hypersurface (i.e., a codimension-one submanifold) is *timelike*, *spacelike* or *lightlike / null* if a normal vector n^a at each point is respectively spacelike, timelike or null. Timelike and spacelike hypersurfaces, when endowed with the pullback $g^{(0)}$ of g under the inclusion map (sometimes called *first fundamental form* – see the paragraph after the next), are respectively Riemannian and Lorentzian manifolds by themselves. The null case is exceptional, for then a normal is also tangent to the hypersurface, and the pullback of g results in a *degenerate* symmetric bilinear form, for the one-dimensional subspace of null tangent vectors has eigenvalue zero – the remaining directions are necessarily spacelike, for there are no two linearly independent vectors which are mutually orthogonal with respect to g and such that one is lightlike and the other is causal.

Let Σ be a *nondegenerate* hypersurface in a Lorentzian manifold $(\mathscr{M}, g)$ (i.e., Σ is either timelike or spacelike), with unit normal n^a and induced metric $h_{ab} = g_{ab} - g(n,n)n_a n_b$ (such that $h(X,X) = g(X,X)$ if $X \in T\Sigma$, and $h(n,.) = 0$). We can define, for some $\epsilon > 0$, geodesic normal coordinates (p, ϵ') in a tubular neighbourhood $\mathscr{U}$ of Σ in $\mathscr{M}$ by writing $\Sigma \times \{0\} \doteq \Sigma$ and, for $p \in \Sigma$ and ϵ' satisfying $|\epsilon'| < \epsilon$ (beware that ϵ may depend on p, unless Σ is compact), writing $(p, \epsilon') = \gamma(\epsilon')$, where γ is the unique geodesic with $\gamma(0) = p$ and $\dot{\gamma}(0)^a = n^a(p)$ – with this, we can extend n^a from Σ to $\mathscr{U}$, where it still holds that $g(n,n) = \pm 1$ (we use the same notation for the extension of n,

since there is no confusion). As, in this case, we see that $g(n,.) = d\phi$, where $\phi : \mathscr{U} \to \mathbb{R}$ is given by $\phi((p,\epsilon') = \epsilon'$, it follows that $\nabla_a n_b = \nabla_b n_a$ and $n^a \nabla_a n_b = n^a \nabla_b n_a = 0$.

We'll make use of the following routine to relate the intrinsic (given by h) and extrinsic (given by g) geometries of Σ: let $\nabla^{(0)}$ be the Levi-Civita connection associated to the restriction $g^{(0)}$ of g to Σ (= pullback of g under the natural inclusion map / embedding $i^{\Sigma,\mathscr{M}}$ of Σ into $\mathscr{M}$), let's identify vector fields X, Y in Σ with the vector fields in $\mathscr{U}$ obtained by means of parallel transport of X and Y along the geodesics normal to Σ. Hence, we can identify the connection $\nabla^{(0)}$ with the component of $\nabla_X Y$ *tangential* to Σ, i.e.,

$$\nabla^{(0)}_X Y = (g^{-1}h)\nabla_X Y,$$

where $(g^{-1}h)^a_b = g^{ac}h_{bc} = h^a_b$ is the projector onto the subspace tangent to the foliation. The component of $\nabla_X Y$ *normal* to Σ, given by $g(\nabla_X Y, n) = -X^a Y^b \nabla_a n_b$, defines the *extrinsic curvature* (or *second fundamental form*) $K_{ab} \doteq -\frac{1}{2}(\pounds_n h)_{ab} = -\nabla_a n_b$. Notice that, due to the above considerations, K is symmetric and, indeed, a tensor *in* Σ. The desired relation is, thus, given by the Gauss (A.7) and Codazzi-Mainardi *equations* (A.8)

$$h^q_a h^r_b h^s_c h^t_d \mathrm{Riem}(g)_{qrst} = \mathrm{Riem}(h)_{abcd} - 2g(n,n)K_{a[c}K_{d]b}, \tag{A.7}$$

$$h^c_b n^d \mathrm{Ric}(g)_{cd} = 2h^c_{[a} D_{|c|} K^a_{b]}. \tag{A.8}$$

The lightlike case can be treated in the following manner: taking a lightlike hypersurface $\mathscr{S}$, we see that the null geodesics constructed as in the nondegenerate case above are, at the same time, *normal* and *tangent* to $\mathscr{S}$, and generate the latter, in the sense that it's possible to choose an affine parametrization λ common to all these geodesics (using, for instance, an auxiliary Riemannian metric in $\mathscr{M}$ to normalize the lightlike directions), in such a way that constant-λ submanifolds are spacelike, codimension-two submanifolds of $\mathscr{M}$, with null normal and tangent to $\mathscr{S}$ denoted by $k^a = (\frac{d}{d\lambda})^a$. It's possible, in this situation, to impose the following Riemannian structure in $\mathscr{S}$: at each $p \in \mathscr{S}$, whose affine parameter assumes the value λ_p, we say that $X(p), Y(p) \in T_p\mathscr{S}$ are equivalent if $X(p) - Y(p)$ is lightlike, denoting such an equivalence class by $\hat{X}(p)(= \hat{Y}(p))$. That is, we take at each p the quotient $\widehat{T_p\mathscr{S}} \ni \hat{X}(p)$ of $T_p\mathscr{S}$ modulo its degenerate one-dimensional subspace. The metric $\hat{h}$ induced in $T\mathscr{S}$ by the quotient is a (nondegenerate) Riemannian metric with one dimension less, which is identified at each constant-λ submanifold with the metric induced directly by g. Thus, we can define the *null extrinsic curvature* $\hat{K} = -\frac{1}{2}\frac{d}{d\lambda}\hat{h}$.

We can visualize a nondegenerate Σ hypersurface locally as a parametrization of the family of geodesics normal to Σ. We can, more generally, consider a family (or *congruence*) of geodesics with the same causal character, going through an open set

$\mathscr{O} \subset \mathscr{M}$, such that each $p \in \mathscr{O}$ is crossed by a single geodesic of the family (locally, any vector field X^a generate a family of curves with this property, which are geodesics if and only if $X^a \nabla_a X^b = 0$). The difference with respect to the case of hypersurfaces is that the distribution of subspaces of $T_p\mathscr{O}$ normal to the congruence at each p need not be integrable, that is, it needs not to generate a family of hypersurfaces normal to the congruence. We can, nevertheless, yet in this case define the *first fundamental form* of the congruence in the nondegenerate case $h_{ab} \doteq g_{ab} - g(X,X)X_aX_b$ and in the lightlike case $\hat{h}_{ab} \doteq (g_{ab} - X_aX_b)/^\wedge$, and the *(null) second fundamental form* $\overset{(\wedge)}{K}_{ab} = \nabla_b X_a (/^\wedge)$,[1] where X is the (co)vector field tangent to the congruence, which is parametrized in such a manner that X is normalized at ± 1 resp. in the spacelike / timelike case (resp. normalized at 1 with respect to an auxiliary Riemannian metric in the lightlike case).

It's immediate to see that, in the nondegenerate case, K acts on the subspaces normal to the congruence, for $X^aK_{ab} = \frac{1}{2}\nabla_b X^a X_a = 0$ and $X^bK_{ab} = X^b\nabla_b X_a = 0$. In the null case, K (without hats) acts nontrivially in the spacelike normal subspace *and* in the other null direction normal to this subspace, whose tangent vector field of the same time orientation that of X^a we denote by $\bar{X}^a$. This vector field locally generates another congruence of null geodesics, naturally associated to the first – namely, it's the *only* other congruence which shares the same spacelike normal subspaces. the normalization adopted for $\bar{X}^a$ is given by $g(X,\bar{X}) = -2$.

Unlike as in the case of hypersurfaces, however, K_{ab} is not necessarily a *symmetric* tensor: $\overset{(\wedge)}{\omega}_{ab} \doteq \overset{(\wedge)}{K}_{[ab]}$ (called *(null) twist* of the congruence) expresses precisely the failure of the distribution of subspaces normal to the congruence to be integrable. By the Frobenius theorem [Wal84], the congruence is normal to a family of hypersurfaces if and only if $X_{[a}K_{bc]} \equiv 0$. The symmetric part can be decomposed in terms of the trace with respect to $\overset{(\wedge)}{h}$, called *(null) expansion* (or *(null) mean curvature*) $\theta \doteq \overset{(\wedge)}{K}_{ab}\overset{(\wedge)}{h}^{ab}$ (the hat in the null case is unnecessary for the first member for it's a scalar), and the trace-free part, denominated *(null) shear* $\overset{(\wedge)}{\sigma}_{ab} \doteq \overset{(\wedge)}{K}_{(ab)} - \frac{1}{d-1}\overset{(\wedge)}{\theta}.\overset{(\wedge)}{h}_{ab}$ (a factor $(d-2)$ must be substituted for the factor $(d-1)$ in the null case, because of the dimensional "reduction" due to the quotient modulo the equivalence relation associated to $^\wedge$). Summing up,

$$\overset{(\wedge)}{K}_{ab} = \overset{(\wedge)}{\omega}_{ab} + \overset{(\wedge)}{\sigma}_{ab} + \frac{1}{d-1(2)}\overset{(\wedge)}{\theta h}_{ab}. \tag{A.9}$$

The manner the geometry normal to the congruence changes along the parallel transport is given by the matrix Riccati equation

$$X^c\nabla_c K_{ab} = -K^c_b K_{ac} + \mathrm{Riem}(g)_{cbad}X^cX^d. \tag{A.10}$$

[1] We use, for congruences, a sign convention *opposite* to the one employed for hypersurfaces when we define the second fundamental form.

Let us now take the quotient of (A.10) in the null case, which is the case of interest in the present work, and decompose the result in terms of the expansion, shear and twist in the left hand side. Denoting by λ the common affine parametrization of the congruence, we have:

$$\frac{d\theta}{d\lambda} = -\frac{1}{d-2}\theta^2 - \hat{\sigma}^{ab}\hat{\sigma}_{ab} + \hat{\omega}^{ab}\hat{\omega}_{ab} - R_{cd}X^cX^d; \tag{A.11}$$

$$X^c\nabla_c\hat{\sigma}_{ab} = -\theta\hat{\sigma}_{ab} + C(g)_{cbad}X^cX^d;^2 \tag{A.12}$$

$$X^c\nabla_c\hat{\omega}_{ab} = -\theta\hat{\omega}_{ab}. \tag{A.13}$$

Equation (A.11) is the celebrated Raychaudhuri *equation*.

Any one-parameter subfamily of geodesics γ_λ, $-\epsilon < \lambda < \epsilon$, $\epsilon > 0$ of a congruence produces a vector field $Y \doteq \frac{d}{d\lambda}\restriction_{\lambda=0}$ in γ_0, which denotes the relative displacement a geodesic "infinitesimally close to γ_0". The relative acceleration of two such geodesics is given by the *equation of geodesic deviation* (also called Jacobi *equation*)

$$X^c\nabla_c(X^b\nabla_bY^a) = -R^a_{cbd}X^cX^dY^b. \tag{A.14}$$

More in general, we can define a Jacobi *field* over a geodesic γ as a solution Y of (A.14) with $X = \dot{\gamma}$. We say that a causal geodesic γ *has a pair of conjugate points* $p = \gamma(\lambda_1), q = \gamma(\lambda_2)$, $\lambda_1 < \lambda_2$, if there exists a Jacobi field Y which vanishes nowhere in $\gamma((\lambda_1, \lambda_2))$ but does so at p and q. Geometrically, this means that the geodesics emanating from p and "infinitesimally close" to γ tend to focus at q.

To visualize this phenomenon in the nondegenerate case, let us take an orthonormal frame $e^a_i(\lambda)$, $i = 0, \ldots, d-1$ parallelly propagated along γ, such that $e^a_0(\lambda) = X^a(\lambda)$. Taking $\lambda_1 = 0$ for simplicity, we have $Y^a(0) = 0$ and, hence,

$$Y^i(\lambda) = \sum_{j=1}^{d-1} A^i_j(\lambda)\frac{dY^j}{d\lambda}(0),$$

where the matrix A^i_j expressed in terms of the chosen frame satisfies

$$\frac{d^2A^i_j}{d\lambda^2} = -R^i_{klm}X^kX^mA^l_j.$$

[2] $C(g)_{cbad}X^cX^d$ is a symmetric tensor which assumes a unique value at each equivalence class associated to $\hat{}$.

As

$$\frac{dY^i}{d\lambda} = e^i_b X^a \nabla_a Y^b = e^i_b Y^a \nabla_a X^b = \sum_{j=1}^{d-1} K^i_j Y^j,$$

we have

$$K^i_j = \left[\frac{dA}{d\lambda}\right]^i_k [A^{-1}]^k_j,$$

and thus $\theta = \mathrm{tr}[K] = \frac{1}{\det A}\frac{d}{d\lambda}(\det A)$. Hence, $\det A \to 0$, which implies that A possesses an eigenspace normal to X^a with eigenvalue zero and, thus, there exists a JACOBI field which vanishes at p and q, if and only if $\theta \to -\infty$ at q. The lightlike case is treated by adopting a frame parallelly propagated along γ such that $e^a_0 = X^a$, $e^a_1 = \bar{X}^a$, and $e^a_2, \ldots, e^a_{d-1}$ an orthonormal set such that $g(e_0, e_i) = g(e_1, e_i) = 0$, $i = 2, \ldots, d-1$. The formulae above don't involve e^a_1 in the latter case.

A.2 Global theory

By a *spacetime* one understands a *time orientable* Lorentzian manifold $(\mathscr{M}, g)$, i.e., there exists a timelike vector field $\mathscr{C}^\infty$ T^a in $\mathscr{M}$ which vanishes nowhere (in particular, we can choose $g(T,T) = -1$).

A.2.1 Causal structure, geodesic completeness

Let $(\mathscr{M}, g_{ab})$ be a spacetime, $\mathscr{O} \subset \mathscr{M}$, $p \in \mathscr{O}$. The *chronological* (resp. *causal*) *future* of p with respect to $\mathscr{O}$, denoted by $I^+(p, \mathscr{O})$ (resp. $I^+(p, \mathscr{O})$) is given by

$$\begin{aligned} I^+(p,\mathscr{O}) &\coloneqq \{x \in \mathscr{O} : \exists \gamma : [0,a] \xrightarrow{\mathscr{C}^\infty} \mathscr{O} \text{ future directed,} \\ &\qquad \text{timelike such that } \gamma(0) = p,\ \gamma(a) = x\}; \end{aligned} \tag{A.15}$$

$$\begin{aligned} J^+(p,\mathscr{O}) &\coloneqq \{x \in \mathscr{O} : x = p \text{ or } \exists \gamma : [0,a] \xrightarrow{\mathscr{C}^\infty} \mathscr{U} \text{ future directed,} \\ &\qquad \text{causal such that } \gamma(0) = p,\ \gamma(a) = x\}. \end{aligned} \tag{A.16}$$

Exchanging future with past, one defines in a dual way the *chronological* (resp. *causal*) *past* $I^-(p, \mathscr{O})$ (resp. $J^-(p, \mathscr{O})$) of p with respect to $\mathscr{O}$. It follows from these definitions that $I^\pm(p, \mathscr{O})$ is open and $int(J^\pm(p, \mathscr{O})) = I^\pm(p, \mathscr{O})$. A set $\mathscr{O}_1 \subset \mathscr{O}$ is said to be *causally convex* with respect to $\mathscr{O}$ if, given any $p \leq_{\mathscr{O}} q \in \mathscr{O}_1$, we have $J^+(p, \mathscr{O}) \cap J^-(q, \mathscr{O}) \subset \mathscr{O}_1$.

Using the sets above, we can define chronology and causality relations between two points. Let $p, q \in \mathscr{O} \subset \mathscr{M}$. We say that p *chronologically* (resp. *causally*) *precedes* q

with respect to $\mathscr{O}$ if $p \in I^-(q, \mathscr{O})$ (resp. $p \in J^-(q, \mathscr{O})$). We denote this relation by $p \ll_{\mathscr{O}} q$ (resp. $p \leq_{\mathscr{O}} q$). Equivalently, we say in this case that q *chronologically* (resp. *causally*) *succeeds* p with respect to $\mathscr{O}$, with the notation $q \gg_{\mathscr{O}} p$ (resp. $q \geq_{\mathscr{O}} p$). If $p \leq_{\mathscr{O}} q$ and $p \neq q$, we write $p <_{\mathscr{O}} q$, or, dually, $q >_{\mathscr{O}} p$. If $p \neq q$, $p \not\ll_{\mathscr{O}} q$ and $p \not\gg_{\mathscr{O}} q$ (resp. $p \not\leq_{\mathscr{O}} q$ and $p \not\geq_{\mathscr{O}} q$), we say that p and q are *chronologically* (resp. *causally*) *disjoint* – in this case, we write $p \curlywedge_{\mathscr{O}} q$ (resp. $p \perp_{\mathscr{O}} q$). All chronology and causality relations defined above, as well as the chronological or causal past and future, are defined for nonunit and nonvoid sets in an obvious manner. If $p \neq q \in \mathscr{O} \subset \mathscr{M}$ implies $p \curlywedge_{\mathscr{O}} q$ (resp. $p \perp_{\mathscr{O}} q$), we say that $\mathscr{O}$ is *achronal* (resp. *acausal*) with respect to $\mathscr{O}$.

Some consequences of the definitions of the previous paragraph are:

- $\ll_{\mathscr{O}}$ is an open relation, that is,

$$p \ll_{\mathscr{O}} q \Rightarrow \exists \mathscr{O}_1, \mathscr{O}_2 \subset \mathscr{M} \text{ open such that } p \in \mathscr{O}_1, q \in \mathscr{O}_2 \text{ and } \mathscr{O}_1 \cap \mathscr{O} \ll_{\mathscr{O}} \mathscr{O}_2 \cap \mathscr{O}. \tag{A.17}$$

- $$J^+(J^+(p, \mathscr{O}), \mathscr{O}) = J^+(p, \mathscr{O}), \text{ i.e., } p \leq_{\mathscr{O}} q \text{ and } q \leq_{\mathscr{O}} r \Rightarrow p \leq_{\mathscr{O}} r. \tag{A.18}$$

- $$I^+(I^+(p, \mathscr{O}), \mathscr{O}) = I^+(J^+(p, \mathscr{O}), \mathscr{O}) = J^+(I^+(p, \mathscr{O}), \mathscr{O}) = I^+(p, \mathscr{O}),$$
$$\text{i.e., } (p \ll_{\mathscr{O}} q \text{ and } q \ll_{\mathscr{O}} r) \text{ or } (p \ll_{\mathscr{O}} q \text{ and } q \leq_{\mathscr{O}} r) \text{ or } (p \leq_{\mathscr{O}} q \text{ and } q \ll_{\mathscr{O}} r) \Rightarrow p \ll_{\mathscr{O}} r. \tag{A.19}$$

- $$\overline{I^+(\mathscr{O}_1, \mathscr{O}_2)} = \overline{J^+(\mathscr{O}_1, \mathscr{O}_2)}, \forall \mathscr{O}_1 \subset \mathscr{O}_2, \tag{A.20}$$

  where the closure is taken in the relative topology of $\mathscr{O}_2$.

- $$\partial I^+(\mathscr{O}_1, \mathscr{O}_2) = \partial J^+(\mathscr{O}_1, \mathscr{O}_2), \forall \mathscr{O}_1 \subset \mathscr{O}_2, \tag{A.21}$$

  where the boundary is taken in the relative topology of $\mathscr{O}_2$.

The set defined in (A.21) is called *future achronal boundary* of $\mathscr{O}_1$ with respect to $\mathscr{O}_2$, and constitutes an achronal, topological submanifold of $\mathscr{M}$ such that any $p \in \partial I^+(\mathscr{O}_1, \mathscr{O}_2)$ belongs to a (necessarily unique) geodesic segment, which is achronal

with respect to $\mathscr{O}_2$ and contained in $\partial I^+(\mathscr{O}_1, \mathscr{O}_2)$, which is either past inextendible or has a past endpoint in $\mathscr{O}_1$. Such geodesics are denominated *generators* of $\partial I^+(\mathscr{O}_1, \mathscr{O}_2)$. Notice that (A.17)–(A.21) still hold if we exchange future with past.

One of the most powerful tools in the analysis of the global geometry of Lorentzian manifolds is the *Lorentzian distance*. Let $p \leq_{\mathscr{M}} q \in \mathscr{M}$, and denote by $\Omega_{p,q}$ the space of future directed, causal piecewise $\mathscr{C}^\infty$ curves linking p to q (this space is empty if $p \not\leq_{\mathscr{M}} q$). $\Omega_{p,q}$ inherits the following topology from the space $\mathrm{Path}(\mathscr{M})$ of $\mathscr{C}^0$ curves in $\mathscr{M}$: we say that a sequence of piecewise $\mathscr{C}^\infty$ curves $\{\gamma_n : [0, \lambda] \to \mathscr{M}\}$ converges to $\gamma : [0, \lambda] \to \mathscr{M}$ if $\gamma_n(0) \to \gamma(0)$ and, given any open set $\mathscr{O} \subset \mathscr{M}$ containing $\gamma([0, \lambda])$, there exists $N \in \mathbb{Z}_+$ such that $\gamma_n([0, \lambda]) \subset \mathscr{O}$ for all $n \geq N$. This topology is called $\mathscr{C}^0$ *topology*. Consider $\lambda \in \Omega_{p,q}$ parametrized by $\lambda \in [\lambda_0 = 0, \lambda_k]$, with $\mathscr{C}^\infty$ components γ_i, $i = 1, \ldots, k$, defined respectively in the intervals $[\lambda_{i-1}, \lambda_i]$, $\lambda_0 < \lambda_1 < \cdots < \lambda_k$. The Lorentzian arc length of γ is given by

$$L_g(\gamma) \doteq \sum_{i=1}^{k} \int_{\lambda_{i-1}}^{\lambda_i} \sqrt{-g(\dot{\gamma}_i(\lambda), \dot{\gamma}_i(\lambda))} d\lambda. \tag{A.22}$$

For timelike curves, which correspond to trajectories of observers, (A.22) is also said to be the *proper time* of the trajectory.

Definition A.1 *The* Lorentzian distance *in* $(\mathscr{M}, g)$ *is the function* $d_g : \mathscr{M} \times \mathscr{M} \to \bar{\mathbb{R}}_+ \cup \{+\infty\}$ *given by*

$$(p, q) \mapsto d_g(p, q) = \begin{cases} \sup_{\gamma \in \Omega_{p,q}} L_g(\gamma) & \text{if } p \leq_{\mathscr{M}} q; \\ 0 & \text{otherwise.} \end{cases} \tag{A.23}$$

It follows immediately from the definition that d_g satisfies a *reverse* triangular inequality: if $p \leq_{\mathscr{M}} r \leq_{\mathscr{M}} q$, then

$$d_g(p, q) \geq d_g(p, r) + d_g(r, q).$$

where the value $d_g(p, q) = +\infty$ may be obtained is in the extremal Reissner-Nordström spacetime (see [BEE96]). In the case $p = q$, we have $d_g(p, p) = 0$ or $d_g(p, p) = +\infty$. d_g is lower semicontinuous for all (p, q) where $d_g(p, q) < +\infty$. We say that $\gamma \in \Omega_{p,q}$ is *maximal* if $L(\gamma) = d_g(p, q)$. One can prove [BEE96] that, if γ is maximal, then it's a ($\mathscr{C}^\infty$) causal geodesic, up to reparametrization. In particular, a maximal null geodesic is *achronal*. On the other hand, if a, say, future causal geodesic γ possesses a pair of conjugate points $p \leq q$, then any points p', q' in γ with p' in the past of p and q' in the future of q, we have that the segments of γ which link p' to q and p to q' no longer maximize the Lorentzian arc length between these pairs of points – in particular, if γ is lightlike, it ceases to be achronal if sufficiently extended. The same occurs if a causal

geodesic, *distinct from* γ, links p to q.

Now we recapitulate some causality conditions one can impose on $(\mathscr{M}, g)$. We say that $(\mathscr{M}, g)$ is *chronological* (resp. *causal*) if there is no $p \in \mathscr{M}$ such that $p \ll_{\mathscr{M}} p$ (resp. $p <_{\mathscr{M}} p$) – equivalently, $(\mathscr{M}, g)$ is chronological if and only if $d_g(p, p) = 0$, and causal if and only if $\Omega_{p,p} = \varnothing$, for all $p \in \mathscr{M}$. We say that $(\mathscr{M}, g)$ is *strongly causal* if, for all $p \in \mathscr{M}$, there exist arbitrarily small open neighbourhoods $\mathscr{O}$ of p such that no causal curve has a disconnected intersection with $\mathscr{O}$. This implies that no compact set completely contains a past or future intextendible causal curve (i.e., such that it's not properly contained in any other curve), and, in particular, that $(\mathscr{M}, g)$ is causal.

A particularly interesting class of spacetimes is the one which admits a foliation by acausal hypersurfaces, for these admit a global notion time evolution (possibly supplemented by boundary conditions). Namely, we say that $t \in \mathscr{C}^0(\mathscr{M})$ is a *global time function* if $t \circ \gamma$ is strictly increasing for any future directed causal curve γ. If t is $\mathscr{C}^1$, we equivalently say that t is a global time function if dt is a future directed timelike covector and $t(\mathscr{M}) = \mathbb{R}$. In this case, $t^{-1}(\tau)$ is a spacelike hypersurface for all $\tau \in \mathbb{R}$, and any $p \in \mathscr{M}$ belongs to $t^{-1}(\tau)$ for precisely one value of τ. As dt induces the same time orientation as adopted for $(\mathscr{M}, g)$, it follows that $t^{-1}(\tau)$ is acausal for all $\tau \in \mathbb{R}$. A spacetime which admits a (continuous) global time function is said to be *stably causal*. The name comes from the fact that such spacetimes can be equivalently defined by the following property: there exists a timelike covector field T_a such that the Lorentzian metric $g_{ab} - \lambda T_a T_b$ is chronological for all $\lambda \in [0, 1)$. In our case, $T = dt$; conversely, given T_a as above, it's possible to build a (continuous) global time function [Ger70, Wal84]. The question of how to construct global time $\mathscr{C}^\infty$ functions in this context was partially answered by Seifert [Sei77] and Dieckmann [Die88], and completely recently by Bernal and Sąánchez [BS03, BS05, BS06], thus eliminating the necessity of assuming differentiability of global time functions in the definition of stable causality. Hence, as a differentiable t has no critical points, all constant-t hypersurfaces are diffeomorphic among each other.

Finally, we say that a strongly causal spacetime $(\mathscr{M}, g)$ is *causally simple* if, for all $p \in \mathscr{M}$ one has that $J^{\pm}(p, \mathscr{M})$ is closed, or, equivalently, $J^{\pm}(p, \mathscr{M}) \smallsetminus I^{\pm}(p, \mathscr{M}) = \partial I^{\pm}(p, \mathscr{M})$, and *globally hyperbolic* if, for all $p, q \in \mathscr{V}$, the set $J^{-}(p, \mathscr{M}) \cap J^{+}(q, \mathscr{M})$ is compact if nonvoid.[3] Equivalently, $(\mathscr{M}, g)$ is globally hyperbolic if and only if $\Omega_{p,q}$ is *compact* in the $\mathscr{C}^0$ topology if nonvoid. Thus, it follows that, in a globally hyperbolic spacetime, d_g is continuous and $d_g(p, q) < +\infty$, $\forall p, q \in \mathscr{M}$. Global hyperbolicity clearly implies simple causality, which, on its turn, implies stable causality (for the

[3] Very recently, Bernal and Sąánchez [BS07] showed that we can substitute "causal" for "strongly causal" in the definition of global hyperbolicity given here.

latter assertion, see [BEE96]). All above conditions can be defined for subsets of $\mathscr{M}$.

Let now $\mathscr{S} \subset \mathscr{M}$ be closed and achronal. The *future* (resp. *past*) *domain of dependence* of $\mathscr{S}$, denoted by $D^+(\mathscr{S})$ (resp. $D^-(\mathscr{S})$) consists of the set

$$D^{+/-}(\mathscr{S}) \coloneqq \{p \in \mathscr{M} : \forall \text{ past/future inextendible, causal} \gamma : [0,a) \longrightarrow \mathscr{M} \text{ such that } \gamma(0) = p, \exists b < a \text{ such that } \gamma(b) \in \mathscr{S}\}. \tag{A.24}$$

$D(\mathscr{S}) \doteq D^+(\mathscr{S}) \cup D^-(\mathscr{S})$ is denoted *domain of dependence* or CAUCHY *development* of $\mathscr{S}$. The *edge* of $\mathscr{S}$ (notation: $\dot{\mathscr{S}}$) consists of the points $p \in \mathscr{S}$ such that *any* open neighbourhood of p has points $q \in I^-(p)$, $r \in I^+(p)$ and a timelike curve γ linking q to r with empty intersection with $\mathscr{S}$. If $\dot{\mathscr{S}} = \varnothing$ then $\mathscr{S}$ is a codimension-one embedded topological submanifold of $\mathscr{M}$. A domain of dependence possesses the following properties:

- The set $int(D(\mathscr{S}))$ is globally hyperbolic (in this case, we say that S is a CAUCHY *surface* for $int(D(\mathscr{S}))$), as is $D(\mathscr{S})$ if $\dot{\mathscr{S}} = \varnothing$. Conversely, one can prove that any globally hyperbolic region has a CAUCHY surface – moreover, one can build a global time function t such that $t^{-1}(\tau)$ is a CAUCHY surface for all $\tau \in \mathbb{R}$.
- The achronal, closed set $H^+(\mathscr{S}) \doteq \overline{D^+(\mathscr{S})} \setminus I^-(D^+(\mathscr{S}))$, called *future* CAUCHY *horizon* of $\mathscr{S}$ has the following property: any $p \in H^+(\mathscr{S})$ is contained in a (necessarily unique) achronal, null geodesic segment contained in $H^+(\mathscr{S})$ which is either past inextendible or possesses a past endpoint in $\dot{\mathscr{S}}$. An analogous definition exists for $H^-(\mathscr{S})$, the *past* CAUCHY *horizon* of $\mathscr{S}$;
- The CAUCHY *horizon* $H(\mathscr{S}) \doteq H^+(\mathscr{S}) \cup H^-(\mathscr{S})$ equals $\partial D(\mathscr{S})$.

The main characteristic of a globally hyperbolic spacetime is that it's "dynamically closed", i.e., the CAUCHY problem for any equation of motion which propagates initial data in a locally causal manner (i.e., the support of the solution at each instant grows in time with speed less than that of the light), as, for instance, the wave equation, is *well posed*, i.e., it possesses a unique solution, which depends continuously on the initial data, for the solution associated to initial data with compact support in the time-zero hypersurface has spatially compact support at every other instant. This eliminates the need for boundary conditions, which doesn't necessarily happen in the more general case of stably causal or causally simple spacetimes [Wal80].

On geodesic completeness: we say that a geodesic γ is *past, future complete* or simply *complete* if, respectively, its affine parameter extends to $\mathbb{R}_+$, $\mathbb{R}_-$ or $\mathbb{R}$.[4] $(\mathscr{M}, g)$ is

[4]Here the words "past" and "future", unlike elsewhere, refer only to the affine parameter, rather than to any time orientation whatsoever, for they also apply, as shown afterwards, to spacelike geodesics.

said to be *spacelike, timelike* or *null geodesically complete* if, respectively, any spacelike, timelike or null geodesic is complete. The counterexamples collected and constructed by Geroch in [Ger68] show that the three definitions are logically independent, that is, none of these implies the other. Timelike and/or null geodesic incompleteness are often used criteria to determine the presence of singularities, for they cannot be circumvented by some sort of Cauchy completion with respect to, say, some distance function which generates the manifold topology by means of paracompactness (which implies metrizability [Dug66]).

To conclude, we must consider two important notions for theories of local observables: the *causal complement* of an open set $\mathscr{O}$ with respect to $\mathscr{O} \supset \mathscr{O}_1$ is the set $(\mathscr{O}_1)'_{\mathscr{O}} \doteq \mathrm{int}\{p \perp_{\mathscr{O}} \mathscr{O}_1\}$, and the *causal completion* of $\mathscr{O}_1$ with respect to $\mathscr{O}$ is given by $(\mathscr{O}_1)''_{\mathscr{O}} \equiv (\mathscr{O}_1)'_{\mathscr{O}})'_{\mathscr{O}}$.[5] If $(\mathscr{O}_1)''_{\mathscr{O}} = \mathscr{O}_1$, we say that $\mathscr{O}_1$ is *causally complete* with respect to $\mathscr{O}$. It follows from these definitions that:

- $(\mathscr{O}_1)''_{\mathscr{O}}$ is the smallest causally complete open set with respect to $\mathscr{O}$ which contains $\mathscr{O}_1$. Notice here the importance of our taking $\mathscr{O}_1$ *open* in our definition of causal complement – if, for instance, $\mathscr{O}_1 = \{p\}$ or, more in general, a discrete achronal set, we have $(\mathscr{O}_1)''_{\mathscr{O}} = \varnothing$. See later on how we'll do to define the operations of causal complement for sets which are not necessarily open, for which the property above *doesn't* obligatorily hold.

- $(\mathscr{O}_1)'''_{\mathscr{O}} = (\mathscr{O}_1)'_{\mathscr{O}}$;

- $\mathscr{O}_1 \subset \mathscr{O}_2 \Rightarrow (\mathscr{O}_1)'_{\mathscr{O}} \supset (\mathscr{O}_2)'_{\mathscr{O}}$;

- $(\cup_\alpha \mathscr{O}_\alpha)'_{\mathscr{O}} = \cap_\alpha (\mathscr{O}_\alpha)'_{\mathscr{O}} \;\Rightarrow\; (\cup_\alpha (\mathscr{O}_\alpha)'_{\mathscr{O}})'_{\mathscr{O}} = \cap_\alpha (\mathscr{O}_\alpha)''_{\mathscr{O}}$ and, hence, any intersection of causally complete sets is causally complete – we shall use these formulae to extend the definition of causal complement to arbitrary sets.

- Any *diamond* $\mathscr{O}_{p,q} \doteq I^-(p,\mathscr{O}) \cap I^+(q,\mathscr{O}) = (\{p,q\})''_{\mathscr{O}}$, $p \ll_{\mathscr{O}} q$, is causally complete.

Throughout the present work, we assume that all our spacetimes are *strongly causal*, unless otherwise stated.

Remark A.2 *If $\mathscr{O} = \mathscr{M}$, the part of the notations of this Subsection indicating $\mathscr{O}$ can be omitted.*

[5] Such definitions differ from the ones adopted in [Rib07], where we don't assume $\mathscr{O}$ open. The differences involved, though, don't imply any modification in the proofs of the results of [Rib07].

A.2.2 Conformal infinity

DEFINITION A.3 *The* conformal infinity *or* conformal boundary *of a d-dimensional spacetime $(\mathscr{M}, g)$ is a $d-1$-dimensional spacetime $(\mathscr{I}, \bar{g}^{(0)})$ such that there exists a d-dimensional Lorentzian manifold $(\overline{\mathscr{M}}, \bar{g})$ with boundary (the* conformal closure *or* conformal completion *of $(\mathscr{M}, g)$) satisfying:*

- *$\mathscr{I} \equiv \partial\overline{\mathscr{M}}$; there exists a diffeomorphism Φ of $\mathscr{M}$ onto $\Phi(\mathscr{M}) \doteq \overline{\mathscr{M}} \setminus \partial\overline{\mathscr{M}}$;*
- *$\bar{g}^{(0)}$ is the semi-Riemannian metric (possibly degenerate) induced by $\bar{g}$ in $\mathscr{I}$;*
- *There is a* conformal *or* WEYL factor*, there is, a positive $\mathscr{C}^\infty$ function z in $\mathscr{M}$, which admits a $\mathscr{C}^\infty$ extension to $\overline{\mathscr{M}}$ such that $z\restriction_{\mathscr{I}} \equiv 0$ and $dz\restriction_{\mathscr{I}} \neq 0$ in $\mathscr{I}$ (the* WEYL *factor and its extension to the conformal closure are always denoted by the same symbol, since there is no confusion here) satisfying $\bar{g} = z^2 g$. We denote respectively by $\bar{\nabla}_a$ and ${}^{(0)}\bar{\nabla}_a$ the* LEVI-CIVITA *connections associated to $\bar{g}$ and $\bar{g}^{(0)}$.*

We'll now show what becomes of the formulae for $\bar{\nabla}_a$, $\mathrm{Riem}(\bar{g})$ and $\mathrm{Ric}(\bar{g})$ in $\mathscr{M}$. First, recall that any two linear connections ∇_a and $\bar{\nabla}_a$ differ in its action on the $\mathscr{C}^\infty$-module $\Gamma(\mathscr{M}, T^*\mathscr{M})$ by a tensor C^c_{ab} of rank $(1,2)$, i.e.,

$$\bar{\nabla}_a X_b = \nabla_a X_b - C^c_{ab} X_c. \tag{A.25}$$

Hence, starting from the LEVI-CIVITA conditions

$$\nabla_a g_{bc} = 0,\ \nabla_a \nabla_b f = \nabla_b \nabla_a f,\ \forall f \in \mathscr{C}^\infty(\mathscr{M}),$$

we obtain

$$C^c_{ab} = \frac{1}{2}\bar{g}^{cd}(\nabla_a \bar{g}_{bd} + \nabla_b \bar{g}_{ad} - \nabla_d \bar{g}_{ab}).^6$$

In particular, taking g and $\bar{g}$ as in Definition A.3, it follows $\nabla_a \bar{g}_{bc} = 2z g_{bc} \nabla_a z$ and, hence,

$$C^c_{ab} = \frac{1}{z} g^{cd}(g_{bd}\nabla_a z + g_{ad}\nabla_b z - g_{ab}\nabla_d z). \tag{A.26}$$

Geodesics with respect to ∇_a in general are no longer so with respect to $\bar{\nabla}_a$, for

$$X^a \nabla_a X^b = 0 \ \Rightarrow\ X^a \bar{\nabla}_a X^b = X^a C^b_{ac} X^c = \frac{2}{z} X^b X^c \nabla_c z - \frac{g(X,X)}{z} g^{bd}\nabla_d z.$$

[6] Specializing this formula to a local chart with domain endowed with the MINKOWSKI metric, we thus obtain the usual expression usual in local coordinates for the CHRISTOFFEL symbols.

However, if $X^a = \dot{\gamma}^a(\lambda)$ is null with respect to g (and, thus, $\bar{g}$), where λ is the original affine parametrization of γ, choosing $\bar{\lambda} = e^{2\log z(\gamma(\lambda))}\lambda = z^2\lambda$, it follows that

$$\frac{d^2}{d\bar{\lambda}^2}\gamma(\bar{\lambda}) = 0,$$

i.e., γ is a null geodesic with respect to $\bar{g}$ *up to reparametrization*. The relations involving the Riemann and Ricci tensors become (taking $z^{-1}\nabla_a z = \nabla_a \log z$)

$$\begin{aligned}(\text{A.27})\quad \mathrm{Riem}(\bar{g})^d_{abc} &= \mathrm{Riem}(g)^d_{abc} - 2\nabla_{[a}C^d_{b]c} + 2C^e_{c[a}C^d_{b]e} = \\ &= \mathrm{Riem}(g)^d_{abc} + 2\delta^d_{[a}\nabla_{b]}\nabla_c \log z - 2g^{de}g_{c[a}\nabla_{b]}\nabla_e \log z + \\ &\quad +2(\nabla_{[a}\log z)\delta^d_{b]}\nabla_c \log z - 2(\nabla_{[a}\log z)g_{b]c}g^{df}\nabla_f \log z + \\ &\quad -2g_{c[a}\delta^d_{b]}g^{ef}(\nabla_e \log z)\nabla_f \log z\end{aligned}$$

and (contracting the indices d and b in (A.23))

$$\begin{aligned}(\text{A.28})\qquad \mathrm{Ric}(\bar{g})_{ac} &= \mathrm{Ric}(g)_{ac} - \frac{d-2}{z}\nabla_a\nabla_c z - \frac{1}{z}g_{ac}g^{de}\nabla_d\nabla_e z + \\ &\quad +2\frac{d-2}{z^2}(\nabla_a z)\nabla_c z - \frac{d-3}{z^2}g_{ac}g^{de}(\nabla_d z)\nabla_e z.\end{aligned}$$

It's convenient to have at our disposal the inverse form of (A.28), employed in Chapter 2. Exchanging the roles of g and $\bar{g}$ and taking $z \mapsto z^{-1}$, we arrive at

$$(\text{A.29})\quad \mathrm{Ric}(g)_{ac} = \mathrm{Ric}(\bar{g})_{ac} + \frac{d-2}{z}\bar{\nabla}_a\bar{\nabla}_c z + \bar{g}_{ac}\bar{g}^{de}\left(\frac{1}{z}\bar{\nabla}_d\bar{\nabla}_e z - \frac{d-1}{z^2}(\bar{\nabla}_d z)\bar{\nabla}_e z\right).$$

It follows from formulae (A.2), (A.27) and (A.28) that the Weyl tensor is *invariant* under conformal changes of the metric, i.e.,

$$C(\bar{g})^d_{abc} = C(g)^d_{abc}, \text{ or } C(\bar{g})_{abcd} = z^2 C(g)_{abcd},$$

where it's understood that the raising and lowering of indices of tensors built uniquely from a metric g must be made with g (other cases are treated individually according to the context).

Returning to $\mathscr{I}$, we point out that it's always possible to choose z in such a way that $\bar{\nabla}_a\bar{\nabla}_b z\restriction_{\mathscr{I}} = 0$, with no restrictions on the chosen representative of the conformal class of $\bar{g}^{(0)}$ (a proof of this fact in the case that g satisfies the Einstein equations without matter is presented in Section 2.2). It's customary to call the spacetime $(\mathscr{M}, g)$ simply the *bulk*, and the corresponding conformal infinity $(\mathscr{I}, \bar{g}^{(0)})$, the *boundary*.

An important characteristic of conformal changes of a Lorentzian metric is that the causal structure is an invariant of the conformal structure associated to this metric. In particular, the conformal infinity is also called *null infinity*, due to the invariance of the concept of null geodesics under conformal changes of the metric, as seen above. This is based upon the idea that complete null geodesics may have, from the viewpoint of $(\overline{\mathscr{M}}, \bar{g})$, past and future endpoints (in the sense of "ideal points" of GEROCH, KRONHEIMER and PENROSE [GKP72]) in $\mathscr{I}$. However, one must notice that $\mathscr{I}$ need *not* be the "infinity" for *any* complete null geodesic. A case in which this certainly occurs is when $(\mathscr{M}, g)$ is strongly causal and $(\overline{\mathscr{M}}, \bar{g})$ is compact (example: MINKOWSKI spacetime); however, if $(\overline{\mathscr{M}}, \bar{g})$ is not compact, it may happen that some null geodesics are unable to reach $\mathscr{I}$. This motivates the following

DEFINITION A.4 *Let $(\mathscr{M}, g)$ be a d-dimensional spacetime with conformal infinity $(\mathscr{I}, \bar{g}^{(0)})$. We say that $(\mathscr{M}, g)$ is* asymptotically simple *if any null geodesic in $(\mathscr{M}, g)$ has a unique extension to $(\overline{\mathscr{M}}, \bar{g})$ such that $\mathscr{I}$ contains precisely both endpoints of the latter.*

Obviously, this is only possible if $(\mathscr{M}, g)$ is null geodesically complete. Actually, in the case that $(\mathscr{I}, \bar{g}^{(0)})$ is timelike (and, hence, a spacetime in its own right), one can say more, justifying the name "asymptotically simple":

THEOREM A.1 If $(\mathscr{M}, g)$ is asymptotically simple with timelike conformal infinity, then it's causally simple.

> PROOF. First, notice that if $p, q \in \mathscr{M}$ is such that $p \not\ll_{\overline{\mathscr{M}}} q$, then $p \not\ll_{\mathscr{M}} q$ (a past counterpart holds by an analogous argument), for a timelike curve in $\overline{\mathscr{M}}$ linking p to q can always be slightly deformed so as to result in a timelike curve *contained in* $\mathscr{M}$ and linking p to q. Now, suppose that $p \in \partial I^-(q, \mathscr{M})$ and $p \notin J^-(q, \mathscr{M})$. By the argument above, we have $p \in \partial I^-(q, \overline{\mathscr{M}})$. Moreover, by hypothesis, a null generator γ of $\partial I^-(q, \mathscr{M})$ needs to attain its future endpoint at infinity without crossing q before that. Let r be such an endpoint. Then, $r \in \partial I^-(q, \overline{\mathscr{M}})$ as the latter set is closed. Since the infinity is totally geodesic [Rib07], γ must hit it transversally and, thus, any causal extension of γ must be broken [7]. Hence, if we extend γ slightly towards the future by a null generator segment γ' in $\partial I^-(q, \overline{\mathscr{M}})$ crossing r (by, say, adjusting the affine parameter t of γ' equal to zero at r and extending it up to $t = \epsilon > 0$), then there exists a timelike curve in $\overline{\mathscr{M}}$ linking p to $\gamma'(\epsilon)$ [HE73], which violates the achronality of $\partial I^-(q, \overline{\mathscr{M}})$. Repeat the argument exchanging future with past. □

An asymptotically simple spacetime need not, though, be globally hyperbolic – a prime example is the AdS spacetime.

[7] Recall that a continuous curve segment is said to be *broken* if it's piecewise $\mathscr{C}^\infty$ but not $\mathscr{C}^1$ in a finite set.

Appendix B

Elements of operator algebras

The formalism of operator algebras is a rather powerful language for the discussion of structural aspects of Quantum Physics which are independent of the representation of operators in some fixed Hilbert space, which is absolutely fundamental in the relativistic case (i.e., Quantum Field Theory) and, more in general, in the study of systems with an *infinite* number of degrees of freedom (thermodynamic limits in Quantum Statistical Mechanics, etc.).

C*-algebras and von Neumann algebras (to be seen in Section B.2) are adequate to the discussion of *bounded* operators. We can, when considering unbounded operators, make use of continuous functional calculus (spectral theorem) when the latter applies and treat only bounded functions of these operators, hence avoiding technical problems relative to the determination of domains. When we discuss field operators, as well as their distributional character, it's however convenient to have at our disposal an *algebraic* framework for dealing with such unbounded operators *directly*, albeit in a manner independent of representations. For such, we present a reasonably detailed study of *-algebras in Section B.1, before we introduce C*-algebras or von Neumann algebras, to show that several known results in the context of C*-algebras remain valid or suffer only minor technical modifications in the context of *-algebras. This allows us to treat the cases of algebras of observables and of fields more or less in parallel. For the latter, we'll present the formalism proposed by Wightman, Borchers and Uhlmann in Section B.3.

We limit ourselves to demonstrate results whose proof is simple from the analytical viewpoint and at the same time instructive, and those which are required in the present work in "nonstandard" form. Our main references for this Appendix are [BR87, BR97], and we employ as auxiliary references [Mur90, SW00, Tak01, Tak03a]. For functional analytic aspects, we adopt [RS80, Rud91].

B.1 *-Algebras

This Section is dedicated to purely algebraic aspects. We shall begin by defining the notion of *-algebra and some of its properties.

Definition B.1 *An (associative)* *-algebra *is a vector space* $\mathfrak{F}$ *over* $\mathbb{C}$*, endowed with an associative, distributive* product $x, y \mapsto xy$ *which commute with scalar multiplication, and with an* involution $x \mapsto x^*$ *which satisfies* $x^{**} = x$*,* $(xy)^* = y^* x^*$ *and* $(\alpha x + \beta y)^* = \bar{\alpha} x^* + \bar{\beta} y^*$*, for all* $x, y \in \mathfrak{F}$*,* $\alpha, \beta \in \mathbb{C}$*. We say that* $\mathfrak{F}$ *is* unital *if there exists an* identity *element* $\mathbb{1} \in \mathfrak{F}$ *such that* $\mathbb{1}x = x\mathbb{1} = x$*, for all* $x \in \mathfrak{F}$ *(in this case,* $\mathbb{1}$ *is unique).*

The *centre* of $\mathfrak{F}$ is given by $\mathfrak{Z}(\mathfrak{F}) \doteq \{x \in \mathfrak{F} : [x, y] = 0, \forall y \in \mathfrak{F}\}$, where $[x, y] \doteq xy - yx$ denotes the *commutator* of x and y. A *-algebra $\mathfrak{F}$ is *Abelian* if $\mathfrak{Z}(\mathfrak{F}) = \mathfrak{F}$; at the other extreme, we say that $\mathfrak{F}$ is *factorial* or *primary* if $\mathfrak{Z}(\mathfrak{F}) = \{0\}$. A **subalgebra* $\mathfrak{G} \subset \mathfrak{F}$ is a vector subspace closed under product and involution. Examples of *-subalgebras of $\mathfrak{F}$ are $\mathfrak{Z}(\mathfrak{F})$ and, more in general, the *(relative) commutant* or *normalizer* of a *-subalgebra $\mathfrak{G} \subset \mathfrak{F}$, given by $\mathfrak{G}' = \{x \in \mathfrak{F} : [x, y] = 0, \forall y \in \mathfrak{G}\}$.

We can adjoin an identity $\mathbb{1}$ to a nonunital *-algebra $\mathfrak{F}$ in the following way: let $\widetilde{\mathfrak{F}} = \mathfrak{F} \oplus \mathbb{C}\mathbb{1} \ni (x, \lambda)$ with product $(x, \lambda)(y, \mu) \doteq (xy + \mu x + \lambda y, \lambda\mu)$ and involution $(x, \lambda)^* \doteq (x^*, \bar{\lambda})$. $\mathfrak{F}$ is naturally identified with the *-subalgebra $(\mathfrak{F}, 0)$, and we have $\mathbb{1} = (0, 1)$.

Let $\mathfrak{F}$, $\mathfrak{G}$ be *-algebras. A **morphism* is a linear map $\pi : \mathfrak{F} \to \mathfrak{G}$ which preserves the product and the involution. If π is respectively one-to-one, onto, bijective, we say that π is a **monomorphism, *-epimorphism, *-isomorphism* – we write $\mathfrak{F} \cong \mathfrak{G}$ if there exists a *-isomorphism from $\mathfrak{F}$ to $\mathfrak{G}$. If $\mathfrak{F} = \mathfrak{G}$, we say that π is a **endomorphism*, and a **automorphism* if π, in this case, is bijective. A **representation* of $\mathfrak{F}$ in a Hilbert space $\mathscr{H}$ [1] is a *-morphism $\pi : \mathfrak{F} \to \mathscr{B}(\mathscr{H})$. We say, then that π is *faithful* if π is a *-monomorphism, and *irreducible* (abbreviated *irrep*) if the only subspaces of X invariant under the action of $\pi(\mathfrak{F})$ are $\{0\}$ and X – by the Schur lemma, π is irreducible if and only if $\{a \in \mathscr{B}(\mathscr{X}) : [a, \pi(x)] = 0, \forall x \in \mathfrak{F}\} = \{0\}$.

A *-ideal of $\mathfrak{F}$ is a subspace $\mathfrak{I} \subset \mathfrak{F}$ *invariant under* *, such that $x \in \mathfrak{F}, y \in \mathfrak{I}$ implies $xy \in \mathfrak{I}$. By *-invariance, it follows automatically that $(x^* y^*)^* = yx \in \mathfrak{I}$. Notice that, on the other hand, left *ideals* (not necessarily *-invariant) of $\mathfrak{F}$ are not necessarily right

[1] We demand a Hilbert space structure to avoid worrying about the definition of involution in $\mathrm{End}_{\mathbb{C}}\mathscr{H}$. A definition of *-representation by the latter's elements (i.e., without reference to the inner product) would involve having at our disposal a definition of involution, at least in the image of the *-algebra in $\mathrm{End}_{\mathbb{C}}\mathscr{H}$.

ideals, and vice-versa. $\mathfrak{F}$ is *simple* if it doesn't possess nontrivial *-ideals (i.e., different from $\{0\}$ and $\mathfrak{F}$) – any nontrivial simple representation π of $\mathfrak{F}$ simples is faithful, for $\ker\pi$ is a *-ideal of $\mathfrak{F}$.

Given a *-ideal $\mathfrak{I} \subset \mathfrak{F}$, the *quotient *-algebra* $\mathfrak{F}/\mathfrak{I}$ of $\mathfrak{F}$ modulo $\mathfrak{I}$ has as elements the cosets $[x] = x + \mathfrak{I} \subset \mathfrak{F}$, where $[x] = [y]$ if $x - y \in \mathfrak{I}$. For instance, if $\pi : \mathfrak{F} \to \mathfrak{G}$ is a *-morphism, it follows that $\mathfrak{F}/\ker\pi \cong \pi(\mathfrak{F}) \subset \mathfrak{G}$. The quotient possesses the following universal property: given a *-algebra $\mathfrak{G}$ and a *-morphism $\rho : \mathfrak{F} \to \mathfrak{G}$ such that $\ker\rho \subset \mathscr{I}$, there exists a unique *-morphism $[\rho] : \mathfrak{F}/\mathfrak{I} \to \mathfrak{G}$ such that $\rho(x) = [\rho]([x])$ for all $x \in \mathfrak{F}$. A *-endomorphism π of $\mathfrak{F}$ induces a **-endomorphism* $[\pi] : \mathfrak{F}/\mathfrak{I} \ni [x] \mapsto \pi([x]) = \pi(x) + \pi(\mathscr{I})$ if and only if $\mathfrak{I}$ is *invariant* under π, i.e., $\pi(\mathfrak{I}) \subset \mathfrak{I}$. In particular, $[\pi]$ is an injective *-endomorphism of $\mathfrak{F}/\mathfrak{I}$ if and only if, in addition, $\ker\pi \subset \mathfrak{I}$ and $\pi(\mathfrak{I})supseteqq\mathfrak{I}$.

We state below four fundamental theorems on *-algebras – the latter three are the so-called **-Isomorphism Theorems*, analogous to the Isomorphism Theorems for rings and modules (the incorporation of the operation of involution * is immediate and can be checked directly).

THEOREM B.1 (FUNDAMENTAL THEOREM OF *-MORPHISMS) *Let $\mathfrak{F}$ and $\mathfrak{G}$ be two *-algebras, and $\rho : \mathfrak{F} \to \mathfrak{G}$ a *-morphism. Then the induced *-morphism $\tilde{\rho} : \mathfrak{F}/Ker\rho \to \rho(\mathfrak{F}) \subset \mathfrak{G}$ is a *-isomorphism.*

PROOF. Immediate. □

Theorem B.1 can be generalized, resulting in the

THEOREM B.2 (FIRST *-ISOMORPHISM THEOREM [2]) *Let $\mathfrak{F}$ and $\mathfrak{G}$ two *-algebras, and $\rho : \mathfrak{F} \to \mathfrak{G}$ a *-epimorphism. Then, $\mathfrak{H} \mapsto \rho(\mathfrak{H})$ establishes a one-to-one correspondence between the additive subgroups $\mathfrak{H} \supset Ker\rho$ of $\mathfrak{F}$ and the additive subgroups of $\mathfrak{G}$. Under this correspondence, $\mathfrak{H}$ is a *-subalgebra (resp. *-ideal) of $\mathfrak{F}$ if and only if $\rho(\mathfrak{H})$ is a *-subalgebra (resp.*-ideal) of $\mathfrak{G}$. Moreover, if $\mathfrak{I} \supset Ker\rho$ is a *-ideal, then the map*

$$x + \mathfrak{I} \mapsto \rho(x) + \rho(\mathfrak{I})$$

*is a *-isomorphism of $\mathfrak{F}/\mathfrak{I}$ onto $\mathfrak{G}/\rho(\mathfrak{I})$.*

PROOF. The first assertion follows immediately from Theorem B.2. The remaining one is demonstrated in a manner analogous to the case of rings – see the proof of Theorem 2.6 of [Jac85], pages 107–108. □

Theorem B.3 (Second *-Isomorphism Theorem) *Let $\mathfrak{F}$ be a *-algebra, $\mathfrak{G}$ a *-subalgebra of $\mathfrak{F}$, and $\mathfrak{I}$ a *-ideal of $\mathfrak{F}$. Then $\mathfrak{G}+\mathfrak{I}$ is a *-subalgebra of $\mathfrak{F}$, $\mathfrak{G}\cap\mathfrak{I}$ is a *-ideal of $\mathfrak{G}$, $\mathfrak{I}$ is a *-ideal of $\mathfrak{G}+\mathfrak{I}$, and $(\mathfrak{G}+\mathfrak{I})/\mathfrak{I}\cong\mathfrak{G}/(\mathfrak{G}\cap\mathfrak{I})$.*

Proof. Analogous to the case of rings – see [Jac85], page 108. □

Theorem B.4 (Third *-Isomorphism Theorem) *Let $\mathfrak{F}$ be a *-algebra, and $\mathfrak{I}\subset\mathfrak{J}$ two *-ideals of $\mathfrak{F}$. Then $\mathfrak{I}$ is a *-ideal of $\mathfrak{J}$, and $\mathfrak{F}/\mathfrak{J}\cong(\mathfrak{F}/\mathfrak{I})/(\mathfrak{J}/\mathfrak{I})$.*

Proof. Follows immediately from Theorem B.2. □

The demonstration of the analogs of Theorems B.2 and B.3 for rings consist only in the construction of the isomorphisms and explicit checking of the relevant properties.

Remark B.2 *In the case of topological *-algebras, in particular the examples to be discussed throughout this Appendix, Theorems B.1 to B.4 not only at the algebraic level but also at the topological level, as far as one adds the adjective* "closed" *(in the respective topologies) to the terms "subgroup", "*-subalgebra" and "*-ideal" – the continuity of the *-morphisms is automatic in all cases, by virtue of the C* property of the (semi)norms (see Definitions B.3 and B.8). In this case, given a (semi)norm $\|.\|$ in a *-algebra $\mathfrak{F}$, the corresponding (semi)norm in the quotient *-algebra $\mathfrak{F}/\mathfrak{I}$, where $\mathfrak{I}\subset\mathfrak{F}$ is a *-ideal, is given by*

$$\mathfrak{F}/\mathfrak{I}\ni[x]\mapsto\|[x]\|\doteq\inf_{y\in\mathfrak{I}}\|x+y\|.$$

Completeness of this norm (or separating family of seminorms) can be checked directly; for the verification of the C property in the case of C*-algebras, see Proposition 2.2.19 of [BR87].*

A *state* ω over a unital *-algebra $\mathfrak{F}$ is a *positive* linear functional on $\mathfrak{F}$, i.e., $\omega(x^*x)\geq 0$ for all $x\in\mathfrak{F}$, which satisfies $\omega(\mathbb{1})=1$. [3] We say that ω is *faithful* if $\omega(x^*x)>0$ for all $x\neq 0$ (the reason behind the name will emerge soon). The set of states over $\mathfrak{F}$ is *convex*: if ω_1 and ω_2 are states, then $\lambda\omega_1+(1-\lambda)\omega_2$ is as well, for all $\lambda\in(0,1)$. If ω doesn't admit a decomposition $\omega=\lambda\omega_1+(1-\lambda)\omega_2$ for $\omega_1,\omega_2\neq\omega$, $\lambda\in(0,1)$, we say that ω is *pure* (equivalently, ω is pure if it cannot be written as the sum of two positive linear functionals $\phi_1,\phi_2\notin\mathbb{R}_+\omega$). Otherwise, we say that ω is *mixed*. We denote the set

[2] There seems to be no agreement in the literature on the coining "First Isomorphism Theorem", neither for rings and modules, nor for groups. Some authors attribute this name to the analog of Theorem B.1, and others, to the analog of Theorem B.2, as we do here, for Theorem B.2 can be understood, as said above, as a generalization of Theorem B.1, and more accordingly with the Second and Third (*-)Isomorphism Theorems.

[3] If there is a topology in $\mathfrak{F}$ at our disposal, we can define the normalization condition of a state even in the absence of a unit.

of all states over $\mathfrak{F}$ by $\mathscr{S}_{\mathfrak{F}}$.

The positivity of ω leads to the following, crucial property:

LEMMA B.5 (CAUCHY-SCHWARZ INEQUALITY) *Let ϕ be a positive linear functional over a (not necessarily unital) *-algebra $\mathfrak{F}$. Then, (i) $\phi(x^*y) = \overline{\phi(y^*x)}$ (in particular, if $\mathfrak{F}$ is unital, $\phi(x^*) = \overline{\phi(x)}$) and (ii) $|\phi(x^*y)|^2 \leq \phi(x^*x)\phi(y^*y)$, for all $x, y \in \mathfrak{F}$ (in particular, if $\mathfrak{F}$ is unital, $|\phi(x)|^2 \leq \phi(\mathbb{1})\phi(x^*x)$).*

PROOF (SKETCH). For all $\lambda \in \mathbb{C}$, we have $\phi((\lambda x + y)^*(\lambda x + y)) \geq 0$. By linearity,

$$|\lambda|^2\phi(x^*x) + \bar{\lambda}\phi(x^*y) + \lambda\phi(y^*x) + \phi(y^*y) \geq 0.$$

The reality of the first member of the inequality above implies (i), whence it follows we can now take $\lambda \in \mathbb{R}$. Choosing an appropriate interval for λ, one proves (ii). □

With Lemma B.5 in our possession, we can easily build a representation of $\mathfrak{F}$ by densely defined linear operators in a HILBERT space $\mathscr{H}_\phi$ naturally associated to a positive linear functional ϕ. First, notice that the *annihilator* Annϕ of ϕ, given by Ann$\phi = \{x \in \mathfrak{F} : \phi(x^*x) = 0\}$, is a left ideal of $\mathfrak{F}$. Indeed, Annϕ is a vector subspace of $\mathfrak{F}$, due to the linearity of ϕ and Lemma B.5. Moreover, given $x \in$ Ann$\phi, y \in \mathfrak{F}$, we have $\phi((yx)^*yx)^2 \leq \phi(x^*x)\phi((x^*y^*y)(y^*yx)) = 0$. Consider now the quotient space $\mathfrak{F}/$Ann$\phi \ni [x], [y]$, which admits the scalar product given by $\langle [x], [y] \rangle \doteq \phi(x^*y)$. Completing $\mathfrak{F}/$Ann$\phi \ni [x], [y]$ with respect to the norm $\|.\|$ associated to $\langle ., . \rangle$, we obtain the HILBERT space $\mathscr{H}_\phi \doteq \overline{(\mathfrak{F}/\text{Ann}\phi)}^{\|.\|}$.

Let us consider the operator $\tilde{\pi}_\phi(x)$ associated to x, defined in the dense linear subspace $\mathfrak{F}/$Annϕ by $\tilde{\pi}_\phi(x)[y] \doteq [xy]$. As the adjoint of $\tilde{\pi}_\phi(x)$ is given by $\tilde{\pi}_\phi(x)^* \restriction_{\mathfrak{F}/\text{Ann}\phi} = \tilde{\pi}_\phi(x^*)$ and both are densely defined, it follows that $\tilde{\pi}_\phi(x)$ is closable [RS80], with closure $\overline{\pi_\phi(x)} = \pi_\phi(x)^{**} \subset \pi_\phi(x^*)^*$ (recall that, given densely defined linear operators A, B with respective domains $D(A), D(B)$, $A \subset B$ denotes that B is an *extension* of A, i.e., $D(B) \supset D(A)$ and $B \restriction_{D(A)} = A$) Thus, $\pi_\phi(x) = \overline{\pi_\phi(x)} = \pi_\phi(x)^{**}$ defines a representation of $\mathfrak{F}$ by densely defined, closed linear operators $\mathscr{H}_\phi$. The densely defined antilinear operator given by $S_0 : [x] \mapsto [x*]$ will be crucial in the development of the TOMITA-TAKESAKI modular theory later on.

Notice that the representation π_ϕ is faithful if and only if $\phi(x^*x) > 0$ for all $x \neq 0$. If $\mathfrak{F}$ is *unital* and ω is a *state*, then ϕ_ω is faithful if and only if ω is faithful, justifying the name we've given. For $\mathfrak{F}$ unital, there is yet a special element of $\mathscr{H}_\omega$, given by $\Omega = [\mathbb{1}]$. Obviously, $\|\Omega\| = 1$ if ω is a state; more importantly, it follows that the

set $\pi_\omega(\mathfrak{F})\Omega = \mathfrak{F}/\mathrm{Ann}\phi$ is dense in $\mathscr{H}_\omega$, i.e., Ω is a *cyclic* vector for $\pi_\omega(\mathfrak{F})$. [4] In this case, the representation built above, denoted by the triple $(\mathscr{H}_\omega, \pi_\omega, \Omega)$, is called *cyclic* or Wightman-*GNS representation* of $\mathfrak{F}$ associated to the positive linear functional ω.[5] π_ω is faithful if and only if Ω is not only cyclic but also *separating* for $\mathfrak{F}$, i.e., $\pi_\omega(x)\Omega = 0 \Rightarrow x = 0$.

The Wightman-GNS representation of a *unital* *-algebra $\mathfrak{F}$ associated to a positive linear functional ϕ is unique in the following sense: let π be a representation of $\mathfrak{F}$ by densely defined, closed linear operators in a Hilbert space $\mathscr{H}$, such that:

- There exists a domain $\mathscr{D}$, common to all $\pi(x)$'s, dense in $\mathscr{H}$ and such that $\pi\mathfrak{F}\mathscr{D} \subset \mathscr{D}$;
- Thee exists a vector $\Phi' \in \mathscr{D}$ cyclic for $\pi(\mathfrak{F})$ such that $\pi(\mathfrak{F})\Phi' = \mathscr{D}$ and $\langle \Phi', \pi(x)\Phi' \rangle = \phi(x)$ for all $x \in \mathfrak{F}$.

We can then define the linear operator $U : \mathfrak{F}/\mathrm{Ann}\phi \to \mathscr{H}$ by $U[x] \doteq \pi(x)\Phi'$. U is densely defined and satisfies $\langle U[x], U[y] \rangle = \phi(x^*y) = \langle \pi(x)\Phi', \pi(y)\Phi' \rangle$ for all $x, y \in \mathfrak{F}$, that is, U is an isometry. By the BLT Theorem [RS80], U extends uniquely to an isometry U of $\mathscr{H}_\phi$ into $\mathscr{H}$, injective and with dense range. Invoking once more the BLT Theorem, we have that U is a bijection, and its inverse $U^-1 = U^*$ is also an isometry. That is, the representations π_ϕ and π are unitarily equivalent.[6] Another important consequence is that, if $\Phi = [\mathbb{1}]$ is the cyclic vector of π_ϕ, then $U\Phi = \Phi'$. An immediate consequence of this fact, which we can state as a Corollary, is:

Corollary B.6 *If α is a *-automorphism of a unital *-algebra $\mathfrak{F}$ and ϕ, a positive linear functional such that $\phi \circ \alpha = \phi$, then there exists a unique unitary operator U_α in $\mathscr{H}_\phi$ such that $\pi_\phi(\alpha(x)) = U\pi_\phi(x)U^{-1}$.* □

That is, a sufficient condition for a *-automorphism α to be unitarily implementable in $\mathscr{H}_\phi$ is that ϕ is invariant under the action of α. This is the main mechanism of implementation of symmetries in quantum theory. Finally, there is a close relation between the purity of ω and a "weak" counterpart of the notion of irreducibility for π_ω:

[4] In the case of C*-algebras, it's possible to construct a cyclic vector even in the absence of a unit [Mur90].

[5] GNS calls for the names of Gel'fand and Naimark [GN43] and Segal [Seg47], which proposed the construction above in the case that $\mathfrak{F}$ is a C*-algebra – see Section B.2. Wightman's name comes along as well for the construction above is the core of its Theorem of reconstruction of quantum fields from a hierarchy of k-point functions [SW00]. We shall have more to say about all this in Section B.3.

[6] In the case that $\mathfrak{F}$ is a C*-algebra, the uniqueness of the GNS representation, given by U, still holds even if $\mathfrak{F}$ doesn't have a unit, but in this case the cyclic vector π_ϕ doesn't necessarily belongs to $\mathfrak{F}/\mathrm{Ann}\phi$, and we must analogously impose only that $\pi(\mathfrak{F})\Phi' \subset \mathscr{D}$ is dense in $\mathscr{H}$.

Theorem B.7 *Let ω be a state over a unital *-algebra $\mathfrak{F}$. Define the* weak commutant

$$\pi_\omega(\mathfrak{F})'_w \doteq \{T \in \mathscr{B}(\mathscr{H}_\omega) : \langle T^*[x], \pi_\omega(y)[z]\rangle = \langle \pi_\omega(y^*)[x], T[z]\rangle,\ \forall x, y, z \in \mathfrak{F}\}$$

*of $\pi_\omega(\mathfrak{F})$; $\pi_\omega(\mathfrak{F})'_w$ is a *-invariant linear subspace of $\mathscr{B}(\mathscr{H}_\omega)$. Then $\pi_\omega(\mathfrak{F})'_w = \mathbb{C}\mathbb{1}$ if and only if ω is pure. Moreover, there exists a one-to-one correspondence $\omega_T \leftrightarrow T$ between positive linear functionals ω_T satisfying $\omega_T \leq \omega$ and positive elements T of $pi_\omega(\mathfrak{F})'_w$ satisfying $\|T\| \leq 1$.*

Proof. ($\Rightarrow$) Suppose that ω is mixed. Then there exists a positive linear functional ω' such that $\omega'(x^*x) \leq \omega(x^*x)$, for all x, and such that ω' is not a multiple of ω. Applying the Cauchy-Schwarz inequality, we have $|\omega'(x^*y)|^2 \leq \|\pi_\omega(x)\Omega\|^2\|\pi_\omega(y)\|^2$. Hence, $\pi_\omega(x)\Omega \times \pi_\omega(y)\Omega \mapsto \omega'(x^*y)$ defines a densely defined, bounded sesquilinear form on $\mathscr{H}_\omega$. Thus, there exists a unique bounded operator T in $\mathscr{H}_\omega$ such that $\langle \pi_\omega(x)\Omega, T\pi_\omega(y)\Omega\rangle = \omega'(x^*y)$, and T is not a multiple of $\mathbb{1}$. Moreover, $0 \leq \langle \pi_\omega(x)\Omega, T\pi_\omega(x)\Omega\rangle \leq \|\pi_\omega(x)\|^2$ and thus $0 \leq T \leq \mathbb{1}$. However, in this case,

$$\langle \pi_\omega(x)\Omega, T\pi_\omega(y)\pi_\omega(x)\Omega\rangle = \omega'(x^*yz) = \omega'((y^*x)^*z) = \langle \pi_\omega(y^*)\pi_\omega(x)\Omega, T\pi_\omega(z)\Omega\rangle$$

for all $x, y, z \in \mathfrak{F}$, proving that T is a nontrivial element of $\pi_\omega(\mathfrak{F})'_w$.

($\Leftarrow$) Suppose that $T \in \pi_\omega(\mathfrak{F})'_w$, $T \notin \mathbb{C}\mathbb{1}$. Then, T^* and, thus, $T + T^*$ belong to $\pi_\omega(\mathfrak{F})'_w$, being the latter self-adjoint and nontrivial. Define $S \doteq \lambda(2\|T + T^*\|\mathbb{1} + (T + T^*)) \in \pi_\omega(\mathfrak{F})'_w$, where $\lambda > 0$ is chosen such that $\|S\| < 1$. Then, $0 < S < \mathbb{1}$, and hence there exists a spectral projector P of S such that $0 < P < \mathbb{1}$ and $P \in \pi_\omega(\mathfrak{F})'_w$. Consider the linear functional $\omega'(x) = \langle P\Omega, \pi_\omega(x)\Omega\rangle$. ω' is positive, for

$$\omega'(x^*x) = \langle P\Omega, \pi_\omega(x^*)\pi_\omega(x)\Omega\rangle = \langle \pi_\omega(x)\Omega, P\pi_\omega(x)\Omega\rangle = \langle P\pi_\omega(x)\Omega, \pi_\omega(x)\Omega\rangle \geq 0.$$

Finally,

$$\omega(x^*x) - \omega'(x^*x) = \langle \pi_\omega(x)\Omega, P\pi_\omega(x)\Omega\rangle \geq 0,$$

showing that ω is mixed.

The final assertion final follows automatically from the arguments above. □

B.2 C*-algebras and von Neumann algebras

B.2.1 C*-algebras

Let's now give a sufficient condition so as to make the closed linear operators $\pi_\phi(x)$ built by means of the Wightman-GNS representation in Section B.1 *bounded*. If we can find for all x a $C(x) > 0$ such that $\phi((xa)^*xa) \leq C(x)^2\phi(a^*a)$ for all $a \in \mathfrak{F}$ *and*

any positive linear functional ϕ (i.e., C doesn't depend on ϕ), it then follows that $\|\pi_\phi(x)[a]\| \leq C(x)\|[a]\|$. Hence, by the BLT Theorem, $\pi_\phi(x)$ admits a unique extension to $\mathscr{H}_\phi$, and $\|\pi_\phi(x)\Psi\| \leq C(x)\|\Psi\|$ for all $\Psi \in \mathscr{H}_\phi$. In this case, $\pi_\phi(x^*) = \pi_\phi(x)^*$ and, thus, π_ϕ defines a **-representation* of $\mathfrak{F}$ by bounded operators in $\mathscr{H}_\phi$.

The most obvious manner of obtaining C as in the previous paragraph is to impose a *norm* $\|.\|$ in $\mathfrak{F}$ such that the product and the involution of $\mathfrak{F}$ are continuous with respect to this norm, defining a *normed *-algebra*. In particular, we can choose a norm equivalent to the original such that $\|xy\| \leq \|x\|\|y\|$ and, hence, $\|\mathbb{1}\| = 1$ if $\mathfrak{F}$ is unital (such a property is assumed to hold throughout the whole of the present work). We'll see that $C(x) \doteq \|x\|$ satisfies the condition in the previous paragraph if:

1. $\mathfrak{F}$ is *complete* with respect to $\|.\|$ (then we say that $\mathfrak{F}$ is a Banach **-algebra*), and
2. $\|x^*x\| = \|x\|^2$. In this case, it follows that $\|x\|^2, \|x^*\|^2 \leq \|x\|.\|x^*\|$ and, thus, $\|x\| = \|x^*\|$.

Definition B.3 *A* C*-algebra $\mathfrak{A}$ *is a* Banach **-algebra whose norm satisfies the* C* condition $\|x^*x\| = \|x\|^2$.[7]

A typical example of C*-algebra is the unital *-algebra $\mathscr{B}(\mathscr{H})$ of bounded operators in a complex Hilbert space $\mathscr{H}$. Let's now cite two frequently used results, which moreover show the strength of Definition B.3.

Proposition B.8 *Any *-morphism* ρ *from a unital* Banach **-algebra* $\mathfrak{B}$ *to a C*-algebra* $\mathfrak{C}$ *satisfies* $\|\rho(x)\| \leq \|x\|$ *for all* $x \in \mathfrak{B}$. *If* $\mathfrak{B}$ *is a C*-algebra, then* $\rho(\mathfrak{B})$ *is a C*-subalgebra of* $\mathfrak{C}$, *and* $\|\rho(x)\| = \|x\|$ *for all* x *if and only if* ρ *is a *-monomorphism.*

Proof. See Propositions 2.3.1 and 2.3.3 of [BR87]. □

Proposition B.9 *Any positive linear functional* ϕ *over a C*-algebra* $\mathfrak{A}$ *(not necessarily unital) satisfies the following properties:*

(i) $\phi(x^*) = \overline{\phi(x)}$;

(ii) $|\phi(x)|^2 \leq \|\phi\|\phi(x^*x)$ *(in particular,* $\|\phi\| = \phi(\mathbb{1})$ *if* $\mathfrak{A}$ *is unital);*

(iii) $|\phi(x^*yx)| \leq \phi(x^*x)\|y\|$;

[7] Actually, it's not necessary to impose the condition $\|xy\| \leq \|x\|\|y\|$ or even continuity of the product and of the involution as axioms defining a C*-algebra. Araki and Elliott [AE73] showed that the first condition, and, hence, the condition $\|x\| = \|x^*\|$ follow from the remaining axioms (completeness of the norm, *-algebra structure and the C* condition).

(iv) $\|\phi\| = \sup\{\phi(x^*x) : \|x\| = 1\}$.

Proof. See Proposition 2.3.11 of [BR87]. □

It follows from Proposition B.9(iii) that $\phi((xy)^*xy) \leq \|x\|^2\phi(y^*y)$. Hence, $C(x) = \|x\|$ satisfies the estimate in the first paragraph of this Section, and the GNS representation π_ϕ is, thus, a *-representation of $\mathfrak{A}$ by elements of the C*-algebra $\mathscr{B}(\mathscr{H}_\phi)$. The apparently particular example of C*-algebra given by $\pi_\phi(\mathfrak{A})$, is actually the most general possible, for Gel'fand and Naimark proved [GN43] that any C*-algebra is *-isomorphic to a C*-subalgebra of $\mathscr{B}(\mathscr{H})$ for some Hilbert space $\mathscr{H}$. More precisely, the *universal representation*

$$\mathscr{H}_u = \bigoplus_{\omega\in\mathscr{S}_\mathfrak{A}} \mathscr{H}_\omega,\ \pi_u = \bigoplus_{\omega\in\mathscr{S}_\mathfrak{A}} \pi_\omega$$

is faithful, for one can prove [BR87] that, given $x \in \mathscr{A}$, there exists a pure state ω_x such that $\omega_x(x^*x) = \|x\|^2$. As π_{ω_x} in this case is irreducible (see next paragraph), it follows that π_u has, among its irreducible subrepresentations, π_{ω_x} for all x. Hence, $\pi_u(x) \neq 0$ for all $x \neq 0$.

Finally, let us consider the following class of elements of a C*-algebra $\mathfrak{A}$. We say that $x \in \mathfrak{A}$ is *positive* if it's self-adjoint and its spectrum $\sigma(x)$ satisfies $\sigma(x) \subset \bar{\mathbb{R}}_+$, or, equivalently, can be written as $x = y^*y$, $y \in \mathfrak{A}$.[8] The positive elements of $\mathfrak{A}$ form a convex cone $\mathfrak{A}_+$ which satisfies $\mathfrak{A}_+ \cap (-\mathfrak{A}_+) = \{0\}$, and any self-adjoint element $x \in \mathfrak{A}$ can be written as the orthogonal decomposition $x = x_+ - x_-$, where $x_\pm = \frac{1}{2}(x \pm |x|) \in \mathfrak{A}_+$, $|x| = \sqrt{x^*x}$ and $x_+x_- = 0$. We can, thus, impose the following ordering $\leq$ to the self-adjoint elements of $\mathfrak{A}$, given by $x \leq y \Leftrightarrow y - x \in \mathfrak{A}_+$. We see immediately that $A \leq B$ and $B \leq A$ imply $A = B$, and, by the spectral radius formula [RS80, Rud91], it follows that $\mathfrak{A}_+ \ni x \leq \|x\|\mathbb{1}$. Hence, when we refer ourselves to increasing or decreasing subsets of $\mathfrak{A}_+$ and to upper limits, suprema, lower limits or infima of subsets of $\mathfrak{A}_+$, it'll be always with respect to the ordering $\leq$ defined above.

B.2.2 von Neumann algebras

Let us now consider a state ω over $\mathfrak{A}$. In this case, the weak commutant $\pi_\omega(\mathfrak{A})'_w$ is a C*-algebra, and coincides with the *commutant* $\pi_\omega(\mathfrak{A})' \doteq \{T \in \mathscr{B}(\mathscr{H}_\omega) : [T, \pi_\omega(A)] \doteq$

[8] At first sight, we could define positivity directly in a *-algebra in Section B.1 by employing this condition. We avoid this explicit terminology in this more general case (except, of course, in the definition of positive linear functionals and states over a *-algebra), for the Wightman-GNS representation of of a "positive" element in this sense defines a symmetric and closed operator, but not necessarily a self-adjoint one. Thus, the intuitive notion of positive operator as referring to the non negativity of the spectrum is lost for *-algebras, which could lead to dangerous misunderstandings.

$T\pi_\omega(A) - \pi_\omega(A)T = 0, \forall A \in \mathfrak{A}\}$ of $\pi_\omega(\mathfrak{A})$ in $\mathscr{B}(\mathscr{H}_\omega)$. We can then strengthen the conclusion of Theorem B.7: ω is pure if and only if π_ω is irreducible. Moreover, $\pi_\omega(\mathfrak{A})'$ is a C*-algebra of quite a special kind, for $\pi_\omega(\mathfrak{A})' = (\pi_\omega(\mathfrak{A})')''$.

Definition B.4 *Let $\mathfrak{R}$ be a unital C*-algebra, concretely realized as a C*-subalgebra of $\mathscr{B}(\mathscr{H})$ for some* Hilbert *space $\mathscr{H}$. We say that $\mathfrak{A}$ is a* von Neumann algebra *if $\mathfrak{R} = \mathfrak{R}''$.*

We say that a primary von Neumann algebra $\mathfrak{R}$, i.e., $\mathfrak{Z}(\mathfrak{F}) \doteq \mathfrak{F} \cap \mathfrak{F}' = \mathbb{C}\mathbb{1}$, is simply a *factor*. The *Double Commutant Theorem* of von Neumann tells us that a nontrivial *-subalgebra $\mathfrak{R}$ of $\mathscr{B}(\mathscr{H})$ is von Neumann algebra if and only if $\mathfrak{R}$ is closed in the strong, weak and *σ-weak* (also called *ultraweak*) topologies of $\mathscr{B}(\mathscr{H})$. The latter is given by the seminorms $\sum_{i=1}^\infty |\langle \Psi_i, .\Phi_i\rangle|$, where $\{\Psi_i\}_i, \{\Phi_i\}_i \subset \mathscr{H}$ satisfy $\sum_{i=1}^\infty \|\Psi_i\|^2, \sum_{i=1}^\infty \|\Phi_i\|^2 < \infty$. Moreover, given a C*-subalgebra $\mathfrak{A} \in \mathscr{B}(\mathscr{H})$, we have that $\mathfrak{A}''$ is the closure of $\mathfrak{A}$ in the strong, weak and σ-weak topologies. We cite tho results about the latter topology:

Proposition B.10 *Let $\mathfrak{R}$ be a* von Neumann *algebra, and $\mathfrak{I}$ a σ-weakly closed two-sided ideal in $\mathfrak{R}$. Then $\mathfrak{I}$ is a *-ideal, and there exists a projection $E \in \mathfrak{Z}(\mathfrak{R})$ such that $\mathfrak{I} = E\mathfrak{R}$.*

Proof. See Proposition 2.4.22 of [BR87]. □

Theorem B.11 *Given two* von Neumann *algebras $\mathfrak{R}$ and $\mathfrak{S}$, a *-morphism from $\mathfrak{R}$ to $\mathfrak{S}$ is necessarily σ-weakly continuous.*

Proof. See Theorem 2.4.23 of [BR87]. □

If $\mathfrak{A}$ is a C*-subalgebra of $\mathscr{B}(\mathscr{H})$, one sees that $\mathfrak{A}$ is dense in $\mathfrak{A}''$ in the strong, weak and σ-weak topologies of $\mathscr{B}(\mathscr{H})$.

Notice that, if an element $T \in \mathscr{B}(\mathscr{H})$ commutes with $x = x^* \in \mathfrak{R}$, it commutes with all spectral projections of x. Hence, this projections belong to $\mathfrak{R}$. Employing the decomposition of an element of $\mathscr{B}(\mathscr{H})$ in a linear combination of self-adjoint elements, it follows that the linear hull of the projections of $\mathfrak{R}$ is strongly dense in $\mathfrak{R}$; making use of the decomposition in a linear combination of unitary elements [RS80], one can show that $x \in \mathfrak{R}$ if and only if the partial isometry u and the positive element $|x|$ occurring in the polar decomposition $x = u|x|$ of x, as well as all spectral projectors of $|x|$, belong to $\mathfrak{R}$ [BR87]. More in general, given a densely defined, closed linear operator T in $\mathscr{H}$, we say that T is *affiliated* to $\mathfrak{R}$ if $\mathfrak{R}'D(T) \subseteq D(T)$ and $Tx \supseteq xT$ for all $x \in \mathfrak{R}'$. Equivalently, T is affiliated to $\mathfrak{R}$ if and only if the partial isometry U and the spectral projections of the positive operator $|T|$ occurring in the polar decomposition $T = U|T|$

of T belong to $\mathfrak{R}$.

Given a von Neumann algebra $\mathfrak{R}$, its *predual* $\mathfrak{R}_*$ consists of the σ-weakly continuous linear functionals on $\mathfrak{R}$. This is a Banach subspace of the dual $\mathfrak{R}^*$ of the Banach space $\mathfrak{R}$, and the latter coincides precisely with the dual $(\mathfrak{R}_*)^*$ of $\mathfrak{R}_*$ [BR87]. The typical example if $\mathscr{B}(\mathscr{H})$ ($\mathscr{H}$ separable), whose predual is given by the ideal of trace class operators $\mathscr{L}_1(\mathscr{H})$, and the dual pairing is given by the trace in $\mathscr{H}$. We have, in general, the following structural result, due to Sakai:

Theorem B.12 (Sakai [Tak01]) *A C*-algebra $\mathfrak{A}$ s *-isomorphic to a* von Neumann *algebra if and only if it's the dual of some* Banach *space.* □

Due to Theorem B.12, it's customary to call such "abstract" von Neumann algebras *W*-algebras* (*W* corresponds to *"weak"*, denoting closure in the weak topology), but here we'll use both terms alternately when dealing with such C*-algebras. We see also that the topology most naturally associated to a von Neumann algebra is the *σ-weak* topology, and it's this one we refer to when, for instance, we employ the "topologized" version of Theorems B.1 to B.4 in the context of W*-algebras, according to Remark B.2. However, as one can see from Proposition 2.4.2 of [BR87], the product operation is separately σ-weakly continuous in both variables, but it's *not jointly* σ-weakly continuous is the W*-algebra in question is faithfully realized in an *infinite*-dimensional Hilbert space.

Given a C*-algebra $\mathscr{A}$ and a state ω, we have a von Neumann algebra in $\mathscr{H}_\omega$ naturally associated to this pair, given by $\pi_\omega(\mathfrak{A})''$. More in general, it's possible to show that the double dual $\mathfrak{A}^{**}$ of $\mathfrak{A}$ is a C*-algebra *-isomorphic to $\pi_u(\mathfrak{A})''$, called *(universal) enveloping W*-algebra* of $\mathfrak{A}$. This is the smallest W*-algebra containing $\mathfrak{A}$, and is *-isomorphic to $\pi_\omega(\mathfrak{A})''$ if ω is faithful. A state ω over a von Neumann algebra $\mathfrak{R}$ in $\mathscr{H}$ is called *normal* if there exists a trace class element $\rho \in \mathscr{B}(\mathscr{H})$ such that $\mathrm{Tr}(\rho) = 1$ and $\omega(x) = \mathrm{Tr}(\rho x)$. Any state ω over a C*-algebra $\mathfrak{A}$ admits a unique normal extension to $\mathfrak{A}^{**}$, which is faithful if and only if ω is [Tak01].

Normal states possess the following characterization:

Theorem B.13 *Let ω be a state over a* von Neumann *algebra $\mathfrak{R}$ in the* Hilbert *space $\mathfrak{H}$. Then the following are equivalent:*

(i) *ω is normal;*

(ii) *ω is σ-weakly continuous;*

(iii) *Given any increasing net $\{A_\alpha\} \subset \mathfrak{R}_+$ bounded from above, one has $\omega(\sup_\alpha A_\alpha) = \sup_\alpha \omega(A_\alpha)$.*

Proof. See Theorem 2.4.21 of [BR87]. □

A comment about the condition (iii) of Theorem B.13 is in order. It's not obvious that, given an increasing net $\{A_\alpha\} \subset \mathfrak{R}_+$ bounded from above, there exists $\sup_\alpha A_\alpha \in \mathfrak{R}$. That this is nevertheless true, it can be seen as follows: let $\mathfrak{R}_\alpha$ be the closure of $\{A_\beta : \beta > \alpha\}$ in the weak topology of $\mathscr{B}(\mathscr{H})$. As the closed unit ball $\mathscr{B}(\mathscr{H})_1$ of the latter is weakly compact,[9] it follows from the cone structure of $\mathfrak{R}_+$ that there exists $A \in \cap_\alpha \mathfrak{R}_\alpha$. For all A_α, $\{B \in \mathscr{B}(\mathscr{H})_+ : B \geq A_\alpha\}$ is σ-weakly closed and contains $\mathfrak{R}_\alpha$, hence $A \geq A_\alpha$ for all α. Any element B majorizing $\{A_\alpha\}$ majorizes its weak closure and, thus, $B \geq A$. Hence, $A = \sup_\alpha A_\alpha$. Moreover, $A_\alpha \to A$ in the σ-weak topology, for the latter coincides with the weak topology in $\mathscr{B}(\mathscr{H})_1$ [BR87], thus establishing (ii)⇒(iii) in Theorem B.13. We say in general that a positive linear functional ϕ is a von Neumann algebra $\mathfrak{R}$ is *normal* if it satisfies condition (iii) in Theorem B.13.[10]

We conclude this Subsection with the classification of von Neumann algebras introduced by Murray and von Neumann: let

$$\mathscr{P}(\mathfrak{R}) \doteq \{E \in \mathfrak{R} : E = E^* = E^2\}$$

the set of *projections* of the von Neumann algebra $\mathfrak{R}$. $\mathscr{P}(\mathfrak{R})$ has a partial ordering $\leq$ inherited from $\mathfrak{R}_+$, and given any $E, F \in \mathscr{P}(\mathfrak{R})$, there exists the *complement* $E^\perp \doteq \mathbb{1} - E$ of E, the *infimum* $E \wedge F \doteq s - \lim_{n\to\infty}(EF)^n$ (by $s - \lim$ it's understood the limit in the strong topology of $\mathfrak{R}$) and the *supremum* $E \vee F \doteq (E^\perp \wedge F^\perp)^\perp$ of E and F with respect to the ordering $\leq$ [11] – that is, $(\mathscr{P}(\mathfrak{R}), \leq, \vee, \wedge, ^\perp)$ is an orthocomplemented lattice.

We introduce the following equivalence relation in $\mathscr{P}(\mathfrak{R})$: given $E, F \in \mathscr{P}(\mathfrak{R})$, we say that $E \sim F$ if there exists $W \in \mathfrak{R}$ such that $W^*W = E$ and $WW^* = F$ (i.e., W is a *partial isometry* with source E and target F). We say yet that E is a *subprojection*

[9] Notice that any $\mathscr{B}(\mathscr{H})$ is a von Neumann algebra and, as such, it's the dual of a Banach space – the latter, in this case, is formed by its trace class elements [BR87]. The assertion thus follows from the Banach-Alaoglu theorem [RS80].

[10] It was proved by Kadison [Tak01] that a C*-algebra $\mathfrak{A}$ is a W*-algebra if and only if any increasing net $\{A_\alpha\} \subset \mathfrak{A}_+$ bounded from above has a supremum $\sup_\alpha A_\alpha \in \mathfrak{A}$ and the stated that satisfy condition (iii) in Theorem B.13 separate elements in $\mathfrak{A}_+$, i.e., given $0 \neq x \in \mathfrak{A}_+$, there exists a normal state ω such that $\omega(A) \neq 0$.

[11] Given a partially ordered set $(S, \leq)$, the *infimum* of $x, y \in S$ is an element $x \wedge y \in S$ such that $x \wedge y \leq x, y$ and $z \leq x, y \Rightarrow z \leq x \wedge y$, the *supremum* of $x, y \in S$ is an element $x \vee y \in S$ such that $x, y \leq x \vee y$ and $x, y \leq w \Rightarrow x \vee y \leq w$, and a *complement* operation is a bijection $^\perp : S \to S$ which satisfies $(x^\perp)^\perp = x$, $(x \wedge y)^\perp = x^\perp \vee y^\perp$ and $(x \vee y)^\perp = x^\perp \wedge y^\perp$. If any pair $x, y \in S$ possesses a supremum and an infimum, we say that $(S, \leq, \vee, \wedge)$ is a *lattice* – in particular, any lattice is *directed*, i.e., given $x, y \in S$, there exists $z \geq x, y$. If, moreover, S has a complement operation $^\perp$, then we say that $(S, \leq, \vee, \wedge, ^\perp)$ is an *orthocomplemented lattice*.

of F if $E \leq F$ (E is *proper* if moreover $E \neq F$), and define the following order relation in $\mathscr{P}(\mathfrak{R})/\sim \ni [E], [F]$: $[E] \neq [F]$ if E is equivalent to a subprojection of F (the definition clearly doesn't depend on the choice of representative). Employing a faithful *-representation of $\mathfrak{R}$ in a Hilbert space $\mathscr{H}$, it's easy to prove that $E \leq F$ if and only if $EF = E$, justifying our terminology. Moreover, using the one-to-one correspondence $E \mapsto \mathrm{Ran}E$ which exists between elements of $\mathscr{P}(\mathscr{B}(\mathscr{H}))$ and closed linear subspaces of $\mathscr{H}$, we see that

$$\dim \mathrm{Ran}E \leq +\infty \text{ if and only if there is no } 0 \neq F \lneq E \text{ such that } F \sim E.$$

We can then *define* abstractly that $E \in \mathscr{P}(\mathfrak{R})$ is *finite* if it satisfies the condition above, which obviously doesn't depend on the realization of $\mathfrak{R}$ in $\mathscr{H}$. We say otherwise that E is *infinite*.

Definition B.5 (Murray-von Neumann) *Let $\mathfrak{R}$ be a* von Neumann *algebra. A function $d : \mathscr{P}(\mathfrak{R}) \to \bar{\mathbb{R}}_+ \cup \{+\infty\}$ is said to be a* dimensional functions *if $d(E) = 0 \Rightarrow E = 0$, $E \sim F \Rightarrow d(E) = d(F)$ and $E \leq F^\perp \Rightarrow d(E \vee F) = d(E + F) = d(E) + d(F)$ – such properties determine d up to a multiple $\lambda \in \mathbb{R} \setminus \{0\}$, and $d(E) < +\infty$ if and only if E is finite. We say that $\mathfrak{R}$ is* finite *if $d(\mathbb{1}) < +\infty$, and* infinite *otherwise. $\mathfrak{R}$ is* semifinite *if any $E \in \mathscr{P}(\mathfrak{R})$ possesses a finite subprojection $F \neq 0$, and* purely infinite *if any $E \in \mathscr{P}(\mathfrak{R})$ is infinite.*

If $\mathfrak{R}$ is a factor, we say that $\mathfrak{R}$ is:

- Type I_n *if* $\mathrm{Ran}d = \lambda\{0, 1, \ldots, n\}$, $n = 0, \ldots, +\infty$ *(equivalently, $\mathfrak{R}$ is type I if it has a minimal projection $E \neq 0$, i.e., $F \neq E \Rightarrow F = E$). $\mathfrak{R}$ is finite if and only if $n < +\infty$;*
- Type II *if it's semifinite, but not type I. $\mathfrak{R}$ is, then,* type II_1 *if* $\mathrm{Ran}d = \lambda[0, 1]$ *(equivalently, if $\mathfrak{R}$ is finite) and* type II_∞ *if* $\mathrm{Ran}d = \bar{\mathbb{R}}_+ \cup \{+\infty\}$*;*
- Type III *if it's purely infinite, i.e.,* $\mathrm{Ran}d = \{0, +\infty\}$.

Several of the more advanced results about von Neumann algebras make use of the concept of *weight*, which generalizes the concept of positive linear functional. Given a W*-algebra $\mathfrak{R}$, a *weight* in $\mathfrak{R}$ is a function $\omega : \mathfrak{R}_+ \to \bar{\mathbb{R}}_+ \cup \{+\infty\}$ (where we set $0. + \infty = 0$) which satisfies (i) $\omega(A) + \omega(B) = \omega(A + B)$ and (ii) $\omega(\alpha A) = \alpha\omega(A)$, for all $A, B \in \mathfrak{R}_+$, $\alpha \in \bar{\mathbb{R}}_+$. A weight τ is said to be *tracial* (or simply a *trace*) if $\omega(A^*A) = \omega(AA^*)$, for all A. Writing $\mathfrak{R}_{\omega+} \doteq \{A \in \mathfrak{R}_+ : \omega(A) < +\infty\}$ and $\mathfrak{L}_\omega \doteq \{A \in \mathfrak{R} : \omega(A^*A) < +\infty\}$, we have immediately that $\mathfrak{R}_{\omega+}$ is an *hereditary* cone in $\mathfrak{R}_+$, i.e., $0 \leq A \leq B \in \mathfrak{R}_{\omega+} \Rightarrow A \in \mathfrak{R}_{\omega+}$, and $\mathfrak{R}_\omega \doteq \mathrm{span}_{\mathbb{C}}\mathfrak{R}_{\omega+} = \mathfrak{L}_\omega^*\mathfrak{L}_\omega$. Hence, the weight ω extends by linearity to a positive linear functional in the *-algebra $\mathfrak{R}_\omega$ – in

particular, if ω is a trace, then $\omega(AB) = \omega(BA)$ for all $A, B \in \mathfrak{R}_\omega$. We say that a weight ω is *normal* if it satisfies condition (iii) of Theorem B.13, *faithful* if $\omega\restriction_{\mathfrak{R}_\omega}$ is faithful, and *semifinite* if $\mathfrak{R}_\omega$ is σ-weakly dense in $\mathfrak{R}$. It's possible to prove that *any* von Neumann algebra $\mathfrak{R}$ admits a normal, faithful semifinite weight, and that $\mathfrak{R}$ is semifinite if and only if it admits a normal, faithful semifinite *trace* τ [Tak01, Tak03a] – the reciprocal of the last assertion is immediate, for if E is a projection in $\mathfrak{R}_\tau$, then E is necessarily finite.

σ-finite von Neumann algebras $\mathfrak{R}$, i.e., such that any mutually orthogonal subset $S \in \mathscr{P}(\mathscr{R})$ (i.e., which satisfies $E \leq F^\perp$ for all $E, F \in S$) is *countable*, can be decomposed in a "direct integral" of factors, in terms of the set of central projections $\mathfrak{Z}(\mathfrak{R}) \cap \mathscr{P}(\mathscr{R})$. $\mathfrak{R}$ is σ-finite if and only if it admits a normal, faithful state ω [BR87]. Thus, one can say that, for such algebras, the classification task is reduced to the classification of factors. A refinement of the classification given by Definition B.5, for type-III factors, will be presented succinctly at the end of Section B.2.3.

B.2.3 Tomita-Takesaki modular theory

We're now in a position to discuss a structure typical of von Neumann algebras, extremely useful and powerful. Consider a C*-algebra $\mathfrak{A}$ and a *faithful* state ω. We then identify the enveloping W*-algebra $\mathfrak{A}^{**} = \pi_\omega(\mathfrak{A})'' \doteq \mathfrak{R}$ in the Hilbert space $\mathscr{H}_\omega \doteq \mathscr{H}$, which possesses, on its turn, a cyclic and separating vector Ω. Let's now define the densely defined antilinear operators S_0 (already mentioned *en passant* in Section B.1) and F_0 as

$$D(S_0) = \mathfrak{R}\Omega,\ S_0 A\Omega \doteq A^*\Omega;\ D(F_0) = \mathfrak{R}'\Omega,\ S_0 A'\Omega \doteq A'^*\Omega$$

(F_0 is densely defined, for a vector Φ is cyclic for a von Neumann algebra $\mathfrak{S}$ if and only if it's separating for $\mathfrak{S}'$). It follows that $S_0 \subseteq F_0^*$ and $F_0 \subseteq S_0^*$, and, thus, S_0 as well as F_0 are closable. One can yet prove that [BR87]

$$F_0^* = \bar{S}_0 \doteq S \text{ and } S_0^* = \bar{F}_0 \doteq F. \tag{B.1}$$

Definition B.6 *Let J be the* anti-unitary *operator (i.e., J is antilinear, invertible and such that $\langle J\Phi, J\Psi\rangle = \overline{\langle \Phi, \Psi\rangle} = \langle \Psi, \Phi\rangle$ for all $\Phi, \Psi \in \mathscr{H}$) and Δ the positive operator of the polar decomposition $S = J\Delta^{\frac{1}{2}}$ of S. We say that Δ is the* modular operator*, and J, the* modular conjugation*, associated to* $(\mathfrak{R}, \Omega)$.

The name has its origin in harmonic analysis: when we define the involution of the convolution *-algebra of compactly supported continuous functions on a locally compact topological group, we need to introduce the Radon-Nikódym derivative of the right-invariant Haar measure with respect to the left-invariant Haar measure,

called *modular function* (this function is equal to 1 for Abelian or compact groups). It follows from (B.1) that

(B.2)

$$\Delta = FS,\ \Delta^{-1} = SF,\ F = J\Delta^{-\frac{1}{2}},\ J = J^*\ (\text{and, hence, } J^2 = \mathbb{1})\ \text{and}\ \Delta^{-\frac{1}{2}} = J\Delta^{\frac{1}{2}}J.$$

The most important result involving J and Δ is the

THEOREM B.14 (TOMITA-TAKESAKI [BR87, Tak03a]) *Let $\mathfrak{R}$ be a* VON NEUMANN *algebra in $\mathscr{H}$ with cyclic and separating vector Ω, and Δ and J respectively the modular operator and the modular conjugation associated to $(\mathfrak{R}, \Omega)$. Then,*

$$J\mathfrak{R}J = \mathfrak{R}' \text{ and } \Delta^{it}\mathfrak{R}\Delta^{-it} = \mathfrak{R},\ \forall t \in \mathbb{R}.$$

□

In particular, $t \mapsto \Delta^{it}$ defines a strongly continuous, one-parameter groups of unitary operators in $\mathscr{H}$, with generator $\log\Delta$, defined by the spectral theorem, which, by Theorem B.14, implements a σ-weakly continuous group of *-automorphisms of $\mathfrak{R}$ and of $\mathfrak{R}'$ (in the latter, the sign of t is changed), called *modular group*. For $\mathfrak{R} = \pi_\omega(\mathfrak{A})''$, where $\mathfrak{A}$ is a C*-algebra, and Ω the cyclic and separating GNS vector associated to a state ω, we write, then, for each $x \in \mathfrak{A}$:

$$j(x) \doteq \pi_\omega^{-1}(J\pi_\omega(x)J),\ \text{and} \tag{B.3}$$

$$\sigma_t^\omega(x) \doteq \pi_\omega^{-1}(\Delta^{it}\pi_\omega(x)\Delta^{-it}). \tag{B.4}$$

Notice that the formulae (B.3) and (B.4) assume values in $\mathfrak{A}^{**}$, i.e., π_ω^{-1} denotes, in a slight but innocuous abuse of notation, its unique σ-weakly continuous extension to $\pi_\omega(\mathfrak{A})''$. A formula of the highest importance which follows from the definition of the modular objects is:

$$\langle\Delta^{\frac{1}{2}}\pi_\omega(x)\Omega, \Delta^{\frac{1}{2}}\pi_\omega(y)\Omega\rangle = \langle J\pi_\omega(x^*)\Omega, J\pi_\omega(y^*)\Omega\rangle = \langle\pi_\omega(y^*)\Omega, \pi_\omega(x^*)\Omega\rangle$$

$$\Downarrow$$

$$\omega(\sigma_{i/2}^\omega(x)\sigma_{-i/2}^\omega(y)) = \omega(yx). \tag{B.5}$$

The TOMITA-TAKESAKI modular theory can be extended to arbitrary VON NEUMANN algebras if we employ a normal, faithful semifinite weight instead of a normal, faithful state. The proofs, however, are considerably more complicated.

Finally, there is a rigid relation between the modular groups associated to two normal, faithful states (or, more in general, semifinite weights) ϕ, ω over a VON NEUMANN algebra $\mathfrak{R}$, given by the

Theorem B.15 (Connes) *There exists a one-parameter family $t \mapsto \Gamma_t \doteq (D\phi : D\omega)_t$ of unitary elements of $\mathfrak{R}$ which satisfies $\sigma_t^{\phi}(A) = \Gamma_t \sigma_t^{\omega}(A)\Gamma_t^*$ and the cocycle relation $\Gamma_{t+s} = \Gamma_t \sigma_t^{\omega}(\Gamma_s)$. In particular, if $\phi = \omega(U.U^*)$, where $U \in \mathfrak{R}$ is a unitary element, then necessarily $\Gamma_t = U^* \sigma_t^{\omega}(U)$.*

Proof. See Theorem 2.7.16 of [BR87] and Theorem 5.3.34 of [BR97]. □

The expression $S(\phi|\omega) \doteq \frac{d}{dt}\omega((D\phi : D\omega)_t)\restriction_{t=+0}$ also defines the *relative entropy* of ϕ with respect to ω [Ara76, BR97, OP93]. The cocycles defined in Theorem B.15 are of fundamental importance in the classification of type-III factors performed by Connes [BR87, Tak03a]. Let $\mathfrak{R}$ be a factor (σ-finite, for simplicity), and $\mathbb{R} \ni u \mapsto \alpha_g$ a one-parameter, σ-weakly continuous group of *-automorphisms of $\mathfrak{R}$, so that it's meaningful to talk about the *σ-weak integral*

$$L^1(\mathbb{R}) \ni f \mapsto \alpha_f(A) \doteq \int f(t)\alpha_t(A)dt,\ A \in \mathfrak{R}.$$

That is, the functional $\mathfrak{R}_* \ni \phi \mapsto \phi(\alpha_f(A)) \doteq \int f(t)\phi(\alpha_t(A))dt$ defines an element of $(\mathfrak{R}_*)^* = \mathfrak{R}$ for each $A \in \mathfrak{R}$, $f \in L^1(\mathbb{R})$. Define the *fixed-point subalgebra* $\mathfrak{R}^{\alpha} \doteq \{A \in \mathfrak{R} : \alpha_t(A) = A, \forall t \in \mathbb{R}\}$. The Arveson *spectrum* of $\alpha_{\cdot}$ is given by the intersection of the supports of the $\mathfrak{R}$-valued tempered distributions $f \mapsto \alpha_f(A)$ through $A \in \mathfrak{R}$, or, equivalently,

$$\sigma(\alpha) \doteq \mathbb{R} \setminus \bigcup \{U \subset \mathbb{R} \text{ open: } \alpha_f(A) = 0,\ \forall A \in \mathfrak{R}, f \in L^1(\mathbb{R}), \text{supp} f \subset U\},$$

and the Connes *spectrum* is given by

$$\Gamma(\alpha) \doteq \bigcap_{E \in \mathscr{P}(\mathfrak{R}) \cap \mathfrak{R}^{\alpha}} \sigma(\alpha\restriction_{\mathfrak{R}E}).$$

Let's now consider the modular group $\alpha_{\cdot} = \sigma_{\cdot}^{\omega}$ associated to the normal, faithful state ω. As the spectrum of a self-adjoint operator is invariant under the adjoint action of a unitary operator [RS80], it follows from Theorem B.15 that $\Gamma(\mathfrak{R}) \doteq \Gamma(\sigma^{\omega})$ doesn't depend on ω. One can yet prove [BR87, Tak03a] that $\exp\Gamma(\mathfrak{R}) = S(\mathfrak{R}) \setminus \{0\}$ is a multiplicative subgroup of $\mathbb{R}_+$, where $S(\mathfrak{R}) = \bigcap_{\omega \in \mathfrak{R}_*,\, \text{Ann}\omega = \{0\}} \sigma(\Delta_{\omega})$ and Δ_{ω} is the Tomita-Takesaki modular operator associated to ω, and that $\Gamma(\mathfrak{R}) = \{0\}$ if and only if $\mathfrak{R}$ is *semifinite*. Hence it follows that the remaining possibilities for $S(\mathfrak{R})$ by virtue of its multiplicative structure

- $S(\mathfrak{R}) = \{0, 1\}$;
- $S(\mathfrak{R}) = \{0\} \cup \{\lambda^n : n \in \mathbb{Z}\}$, $\lambda \in (0, 1)$;
- $S(\mathfrak{R}) = \bar{\mathbb{R}}_+$,

can only be realized for $\mathfrak{R}$ type III.

Definition B.7 (Connes) *We say that* $\mathfrak{R}$ *is* type III_0 *if* $S(\mathfrak{R}) = \{0,1\}$, type III_λ, $\lambda \in (0,1)$, *if* $S(\mathfrak{R}) = \{0\} \cup \{\lambda^n : n \in \mathbb{Z}\}$, $\lambda \in (0,1)$, *and* type III_1 *if* $S(\mathfrak{R}) = \bar{\mathbb{R}}_+$.

Of particular interest in Quantum Field Theory are the hyperfinite factors (a von Neumann algebra $\mathfrak{R}$ is *hyperfinite* or *injective* if it's generated by the union of a sequence of matrix algebras of increasing dimension) of type III_1. These typically appear as thermodynamic limits of local algebras associated to a net of local observables, such as the Araki-Woods factor, given by the thermodynamic limit of algebras of observables of spin systems at finite temperature [BR97]. It was proven by Haagerup, completing former partial results of Connes, that the Araki-Woods factor is the *unique* hyperfinite, type-III_1 factor up to *-isomorphism [Tak03b]. We finalize this Section with two useful criteria to determine if $\mathfrak{R}$ is type III.

Theorem B.16 *Given a normal, faithful state* ω, $\mathfrak{R}$ *is type III if and only if* $\sigma^\omega_\cdot\restriction_{\mathfrak{R}E}$ *is* not *a group of* internal *automorphisms (i.e., there is no* σ*-weakly continuous group* $\mathbb{R} \ni t \mapsto U_t$ *of unitaries of* $\mathfrak{R}$ *such that* $\sigma^\omega_t(.) = U_t^*.U_t$, $\forall t \in \mathbb{R}$*) for* any $E \in \mathscr{P}(\mathfrak{R}) \cap \mathfrak{Z}(\mathfrak{R})$.

> Proof. See Theorem 2.7.17 of [BR87] and the discussion preceding Theorem 5.3.35 of [BR97]. □

The second criterion is due to Driessler [Dri77]:

Theorem B.17 ([Dri77]) *Let* $\mathscr{H}$ *be a separable* Hilbert *space,* $\mathfrak{R}$ *a* von Neumann *subalgebra of* $\mathscr{B}(\mathscr{H})$ *and* Ω *a cyclic and separating for* $\mathfrak{R}$. *Suppose there exists an infinite* von Neumann *subalgebra* $\mathscr{R}_1 \subset \mathfrak{R}$ *and* $\{\alpha_n\}_{n\in\mathbb{Z}_+}$ *a sequence of* *-*automorphisms of* $\mathscr{B}(\mathscr{H})$ *satisfying:*

1. $\alpha_n(\mathfrak{R}_1) \subset \mathfrak{R}_1$, *for all* n;
2. $w - \lim_{n\to\infty} \alpha_n(A_1) = \omega(A_1)\mathbb{1}$, *for all* $A_1 \in \mathfrak{R}_1$, $0 \not\equiv \omega \in \mathfrak{R}_{1*}$;
3. $s - \lim_{n\to\infty}[A, \alpha_n(A_1)] = \{0\}$, *for all* $A \in \mathfrak{R}$, $A_1 \in \mathfrak{R}_1$.

Then $\mathfrak{R}$ *is type III.*

> Proof. Suppose that $P \neq 0$ is a finite projection. In this case, $P\mathfrak{R}P$ is finite and, as we've seen at the end of Subsection B.2.2, possesses a normal, faithful semifinite weight τ. As $P\Omega$ is cyclic and separating for $P\mathfrak{R}P$ in $P\mathscr{H}$, it follows from the analog of Theorem 2.5.31 in [BR87] for weights [Tak03a] that τ is implemented by a single tracial vector $\xi \in P\mathscr{H}$. We obtain, then, that
>
> $$\lim_n \langle \xi, \alpha_n(A)P\alpha_n(B)\xi\rangle \overset{3.)}{=} \lim_n \langle \xi, \alpha_n(AB)\xi\rangle \overset{2.)}{=} \langle \xi, AB\xi\rangle = \langle \xi, BA\xi\rangle.$$
>
> The limit above thus defines a normal, faithful and *finite* trace τ_1 in $\mathfrak{M}_1$, i.e., $\tau_1(A) < +\infty$ for all $A \in \mathfrak{R}_1$, implying that $\mathfrak{R}_1$ is finite, which is absurd. □

B.2.4 C*- e W*-dynamical systems. KMS condition

We shall now arrive at a physical interpretation for the formula (B.5). First, notice that, given $A \in \mathfrak{R}$, the σ-weak integrals

$$A_n = \sqrt{\frac{n}{\pi}} \int \sigma_t^\omega(A) e^{-nt^2} dt,\ n \in \mathbb{Z}_+ \tag{B.6}$$

$$\left(\text{i.e., } \phi(A_n) = \sqrt{\frac{n}{\pi}} \int \phi(\sigma_t^\omega(A)) e^{-nt^2} dt \text{ for all } \phi \in \mathfrak{F}_*\right)$$

define uniquely a sequence of elements (A_n) which σ-weakly converge to A (see Proposition 2.5.18 of [BR87]), and such that each A_n is an *analytic element* for σ_t^ω, i.e., there exists $\lambda > 0$ and a function $f_{A_n} : \mathbb{R} + i(-\lambda, \lambda) \to \mathfrak{R}$ such that $f(t) = \sigma_t^\omega(A)$ for all $t \in \mathbb{R}$ and $z \mapsto \phi(f(z))$ is analytic in $\mathbb{R} + i(-\lambda, \lambda)$ for all $\phi \in \mathfrak{R}_*$. Hence, σ_t^ω has a σ-weakly dense set of analytic elements B in $\mathfrak{R}$. For such, we write $f_B(z) = \sigma_z^\omega(B)$.

Let's return to (B.5). Choosing $A = \mathbb{1}$ and B an analytic element of $\mathfrak{R}$ in the strip $\mathbb{R} + i(-1/2, 1/2)$ (this can always be obtained by an appropriate rescaling of f_B), we have that the function $F(z) = \omega(\sigma_{z+i/2}^\omega(B))$ is continuous and bounded in $\mathbb{R} + i[-1, 0]$, analytic in $\mathbb{R} + i(-1, 0)$, and *periodic* with period 1 in $\Im z$, due to (B.5). Thus, F admits a *bounded* analytic extension to all $\mathbb{C}$ (inclusive in $\mathbb{R} + i\mathbb{Z}$, by Schwarz's reflection principle). Hence, by the Liouville theorem, F must be constant, i.e., $F(z) = \omega(B)$ for all $z \in \mathbb{C}$, in particular for $z = t \in \mathbb{R}$. Since the analytic elements are dense in $\mathfrak{R}$, it follows that ω is *invariant* under σ_t^ω.

More in general, given $A, B \in \mathfrak{R}$, we can choose sequences $(A_n), (B_n)$ such that $\|A_n\| \leq \|A\|$, $\|B_n\| \leq \|B\|$ and $A_n^{(*)}\Omega \to A^{(*)}\Omega$, $B_n^{(*)}\Omega \to B^{(*)}\Omega$ (this is always possible by Kaplansky's density theorem – see [BR87], Theorem 2.4.16). In this case, it follows from (B.5) and Phragmén-Lindelą̈f's Three Line Theorem (see Theorem 6.4 in [Lan99]) that $\omega(A_n f_{B_n}(z))$ converges uniformly in $\mathbb{R} + i[-1/2, 1/2]$ to a function $F'_{A,B}$, continuous and bounded in $\mathbb{R} + i[-1/2, 1/2]$ and analytic in $\mathbb{R} + i(-1/2, 1/2)$ (see Proposition 5.3.7 in [BR97] for a more detailed proof), such that, for all $t \in \mathbb{R}$,

$$F'_{A,B}(t + i/2) = \omega(A\sigma_t^\omega(B)) \text{ and } F'_{A,B}(t - i/2) = \omega(\sigma_t^\omega(B)A). \tag{B.7}$$

We can recast the assertion above in in the following form: let $f \in \mathscr{S}(\mathbb{R})$ such that $\hat{f} \in \mathscr{C}_c^\infty(\mathbb{R})$. In this case, there exists $R > 0$ such that $\text{supp}\hat{f} \subset [-R, R]$ and, hence, by the Paley-Wiener theorem [Hör90], f is an entire function which satisfies $|f(x + iy)| = O(x^{-\infty} e^{R|y|})$ for all $x, y \in \mathbb{R}$ (indeed, this Theorem also tells us that this estimate completely characterizes the f's which satisfy the hypothesis above). We can then rewrite (B.7) as

$$\int f(t + i/2)\omega(A\sigma_t^\omega(B))dt = \int f(t - i/2)\omega(\sigma_t^\omega(B)A)dt,\ \forall f : \hat{f} \in \mathscr{C}_c^\infty(\mathbb{R}). \tag{B.8}$$

Let $\mathfrak{A}$ be a C-algebra and $\mathbb{R} \ni t \mapsto \alpha_t$ a one-parameter group of *-automorphisms of $\mathfrak{A}$, which we assume strongly continuous for simplicity, i.e., $\|\alpha_t(x) - x\| \stackrel{t\to 0}{\longrightarrow} 0$ for all $x \in \mathfrak{A}$. The pair $(\mathfrak{A}, \alpha_t)$ is said to be a* C*-dynamical system. *If $\mathfrak{A}$ is a* von Neumann *algebra and α is, instead, σ-weakly continuous, we say that the pair $(\mathfrak{A}, \alpha_t)$ is a* W*-dynamical system.

Notice that, if $\mathfrak{A}$ is a C*-subalgebra of $\mathscr{B}(\mathscr{H})$ for some $\mathscr{H}$, the C*-dynamical system $(\mathfrak{A}, \alpha_t)$ extends uniquely to the W*-dynamical system $(\mathfrak{A}'', \alpha_t)$, due to Theorem B.11.

Definition B.9 *Let $(\mathfrak{A}, \alpha_t)$ be a C*- (resp. W*-)dynamical system, $\beta \in \mathbb{R} \cup \{\pm\infty\}$. A (resp. normal) state ω over $\mathfrak{A}$ is said to be a* (α, β)-KMS state *if it satisfies the* condition of Kubo, Martin and Schwinger*: given $x, y \in \mathfrak{A}$, there exists a function $F_{x,y}$, continuous and bounded in $\mathbb{R} + (sgn\beta)i[0, |\beta|]$ and analytic in $\mathbb{R} + (sgn\beta)i(0, |\beta|)$, such that*

$$F_{x,y}(t) = \omega(x\alpha_t(y)) \text{ and } F_{x,y}(t + i\beta) = \omega(\alpha_t(y)x).$$

In the case $\beta = 0$, ω satisfies $\omega(xy) = \omega(yx)$ for all x, y; then we say that ω is a tracial *state. In the case $\beta = +\infty$, we say that ω is a* ground *state.*

It follows from considerations similar to the ones made for the modular group that a (α, β)-KMS state ω is invariant under α_t, and, by the uniqueness of the GNS representation, there exists a unique implementation of α_t by unitary operators $U_\alpha(t)$ in $\mathscr{H}_\omega$. A consequence of this is that the (unique) normal extension of a (α, β)-KMS state over $\mathfrak{A}$ to $\mathfrak{A}^{**}$ is also a (α, β)-KMS state for the W*-dynamical system $(\mathfrak{A}^{**}, \alpha_t)$, where α_t denotes its σ-weakly continuous extension.

The importance of this concept owes to the fact that the KMS condition characterizes *thermodynamic equilibrium* of the state ω at temperature $T = (k_B\beta)^{-1}$, where k_B is the Boltzmann constant. Indeed, in quantum systems with a finite number of degrees of freedom, the KMS condition characterizes the Gibbs states $\omega_\beta = \frac{\mathrm{Tr}(e^{-\beta H}.)}{\mathrm{Tr}(e^{-\beta H})}$, where H is the Hamiltonian operator which generates α_t. The KMS condition, however, survives the thermodynamic limit, which doesn't happen to the characterization of equilibrium based upon Gibbs states. Its original formulation, given by Kubo, Martin and Schwinger, was based upon Green functions associated to creation and annihilation operators, being later rephrased for C*-dynamical systems in the version given by Definition B.9 by Haag, Hugenholtz and Winnink [HHW67].

Equivalently, by the considerations above, ω is a (α, β)-KMS state if and only if, for all f such that $\hat{f} \in \mathscr{C}_c^\infty(\mathbb{R})$,

$$\int f(t)\omega(x\alpha_t(y))dt = \int f(t + i\beta)\omega(\alpha_t(y)x)dt, \forall x, y \in \mathfrak{A}. \tag{B.9}$$

The case $\beta = +\infty$ is illuminated by the following consideration: the left hand side of (B.9) is nothing more than the distributional Fourier transform of $\omega(x\alpha_t(y))$. If $\mathrm{supp}\hat{f} \subset \mathbb{R}_-$, it follows that $f(t+i\beta)$ decays exponentially as $\beta \to +\infty$, and, thus, the right hand side of (B.9) tends to zero in this limit. Hence, we conclude that the support of the distributional Fourier transform of $\pi_\omega(\alpha_t(x))\Omega$ is contained in $\bar{\mathbb{R}}_+$, justifying the name *"ground state"* for ω.

For $0 < \beta < +\infty$, a (α, β)-KMS state ω satisfies the following "fluctuation-dissipation relation": $\omega(x^*\alpha_t(x))$ and $\omega(\alpha_t(x)x^*)$ are positive distributions; by the Bochner-Schwartz theorem [RS75], the distributional Fourier transforms

$$\hat{\mu}_x(\hat{f}) \doteq \int f(t)\omega(x^*\alpha_t(x))dt,\ \hat{\nu}_x(\hat{f}) \doteq \int f(t)\omega(\alpha_t(x)x^*)dt$$

define Radon measures over $\mathbb{R}$. A manner of seeing this is to use the spectral decomposition of the unitary operators $U_\alpha(t)$ which implement α_t. We can, then, see that $\hat{\mu}_x$ and $\hat{\nu}_x$ are absolutely continuous one with respect to the other, and the Radon-Nikódym derivative of $\hat{\mu}_x$ with respect to $\hat{\nu}_x$ is given by

$$\frac{d\hat{\mu}_x}{d\hat{\nu}_x}(p) = e^{-\beta p},\ \text{or, formally, } d\hat{\mu}_x(p) = e^{-\beta p}d\hat{\nu}_x(p). \tag{B.10}$$

If we consider the signed measures $\hat{\delta}_{x,y}$ and $\hat{\phi}_{x,y}$ given respectively by the Fourier transforms of $\omega([x, \alpha_t(y)])$ and $\omega(\{x, \alpha_t(y)\})$ ($\{a, b\} \doteq ab + ba$ denotes the *anticommutator* of $a, b \in \mathfrak{A}$), we have

$$d\hat{\delta}_{x,y}(p) = \tanh\left(\frac{\beta p}{2}\right) d\hat{\phi}_{x,y}(p), \tag{B.11}$$

which is precisely the fluctuation-dissipation relation which characterizes the KMS condition, relating bounded perturbations of the dynamics to fluctuations of measurements of observables along time. Other relations, equivalent to (B.10) and (B.11) can be analogously obtained among other Green functions associates to $\omega(x^*\alpha_t(y))$.

Finally, we see that any normal, faithful state ω over a von Neumann algebra $\mathfrak{R}$ is a $(\sigma^\omega_{-\beta t}, \beta)$-KMS state.[12] This example, in a certain sense, is the most general case possible: conversely, given a (α, β)-KMS state ω over a C*-algebra $\mathfrak{A}$, it follows that

Lemma B.18 *Ω is* separating *for $\pi_\omega(\mathfrak{A})''$.*

[12] Amusingly, the work of Haag, Hugenholtz and Winnink [HHW67] preceded by little the original, unpublished work of Tomita [Tom67], and, in a way, motivated the inception of modular theory, as it can be seen by the pioneering exposition of Takesaki [Tak70].

Proof. First, notice that, if $\mathfrak{A}$ is a C*-subalgebra of $\mathscr{B}(\mathscr{H})$ for some Hilbert space $\mathscr{H}$ and $\omega = \langle \Omega, .\Omega \rangle$ is a state over $\mathfrak{A}$ such that Ω is cyclic for $\mathfrak{A}$ and Annω is *-invariant, then Ω is separating for $\mathfrak{A}$: given x such that $\omega(x^*x) = 0$, we have $yx\Omega = 0$ for all $y \in \mathfrak{A}$. However, in this case, $(yx)^*\Omega = x^*y^*\Omega = 0$. By the cyclicity of Ω, it follows that $x = 0$. Returning to our hypotheses, consider now the C*-subalgebra $\mathfrak{R} = \pi_\omega(\mathfrak{A})''$ of $\mathscr{B}(\mathscr{H}_\omega)$. Obviously, Ω is cyclic for $\mathfrak{R}$. Taking $A \in \mathfrak{R}$ such that $\omega(A^*A) = 0$, consider the function $F_{A^*,A}(z)$ as in Definition B.9. Then, $F_{A^*,A}(t) = \omega(A^*\alpha_t(A)) = \langle A\Omega, U_\alpha(t)A\Omega \rangle = 0$. By the Edge-of-the-Wedge Theorem [SW00], it follows that $F_{A^*,A}(z) = 0$ for $z \in \mathbb{R} + i[0, \beta]$. But, in this case, $F_{A^*,A}(i\beta) = \omega(AA^*) = 0$. Hence, by the argument above, $A = 0$. □

Consider now the following situation: given a W*-dynamical system $(\mathfrak{R}, \alpha_t)$ and a normal, (α, β)-KMS state, it follows from Lemma B.18 that ω is faithful in $\pi_\omega(\mathfrak{R})$. However, Annω is then a σ-weakly closed two-sided ideal. Invoking Proposition B.10, there exists a projection $E \in \mathfrak{R} \cap \mathfrak{R}'$ such that Ann$\omega = (\mathbb{1} - E)\mathfrak{R}$. Hence, $\omega(\mathbb{1} - E) = \{0\}$ and ω is faithful in $\mathfrak{R}E$. Moreover,

$$\omega(AE) = \omega(A) = \langle \Omega, \pi_\omega(A)\Omega \rangle.$$

We conclude this Subsection with the following, surprising result of Takesaki, which follows form the modular theory presented in Subsection B.2.3:

Theorem B.19 (Takesaki [BR97, Tak03a]) *Let $\mathfrak{R}$ be a von Neumann algebra and ω a normal state over $\mathfrak{R}$. Then the following are equivalent:*

1. *ω is faithful in $\pi_\omega(\mathfrak{R})$, i.e., there exists a projection $E \in \mathfrak{R} \cap \mathfrak{R}'$ such that $\omega(\mathbb{1} - E) = 0$ and $\omega\restriction_{\mathfrak{R}E}$ is faithful.*

2. *There exists a σ-weakly continuous group α_t of *-automorphisms of $\mathfrak{R}$ such that ω is a KMS state for the W*-dynamical system $(\mathfrak{R}, \alpha_t)$.*

Moreover, if the conditions above are satisfied, it follows that $\alpha_t(E) = E$, and $\alpha_t\restriction_{\mathfrak{R}E}$ coincides with the modular group of $\mathfrak{R}E$ associated to ω. In particular, $\alpha_t\restriction_{\mathfrak{R}E}$ is uniquely determined by ω.

Proof. See Theorem 5.3.10 in [BR97]. □

B.3 Locally convex *-algebras and Borchers-Uhlmann algebras

We recall, following [Rud91] and [RS80], that a topological vector space (always here assumed Hausdorff) X (always over $\mathbb{C}$, unless otherwise stated) is *locally convex* if it admits a fundamental system of open neighbourhoods $\mathscr{V} = \{V_\alpha \ni 0\}_{\alpha \in I}$ such

that each of these neighbourhoods is convex, i.e., $tx + (1-t)y \in V_\alpha$ for all $x, y \in V_\alpha$, $t \in (0,1)$. Equivalently, a topological vector space is locally convex if it admits a family $\{\|.\|_\beta\}_{\beta\in J}$ of *seminorms* (i.e., maps $\|.\|_\alpha : X \to \bar{R}_+$ which satisfy the same properties of a norm, except possibly for $\|x\| = 0 \Rightarrow x = 0$) which *separates points* in X, i.e., given $x \neq y \in X$, there exists $\beta \in J$ such that $\|x\|_\beta \neq \|y\|_\beta$. A system of convex open neighbourhoods of the origin corresponding to this family of seminorms is given, for instance, by $V_{\beta_1,\ldots,\beta_k,n} = \{x \in X : \|x\|_{\beta_i} < n_i^{-1}, \forall i = 1,\ldots,k,\, n_i \in \mathbb{Z}_+\}$. Conversely, notice that any locally convex vector space admits a fundamental system of open, convex and *balanced* neighbourhoods (i.e., $e^{it}V_\alpha = V_\alpha$, for all $t \in [0, 2\pi)$). Defining $\|x\|_\beta \doteq \inf\{t > 0 : t^{-1}x \in V_\beta\}$, it follows from the continuity of scalar multiplication and the fact that V_β is balanced that $\|.\|_\beta$ defines a continuous seminorm – notice that $\|V_\beta\|_\beta \subset [0,1)$, for V_β is open. Moreover, the family $\{\|.\|_\beta\}_{\beta\in J}$ of seminorms separates points in X, since for each $x \neq 0$ there exists β such that $x \notin V_\beta$ and, in particular, $\|x\|_\beta \geq 1$.

Our representative example of a locally convex topological vector space will be $\mathscr{D}(\mathscr{O}) = \mathscr{C}_c^\infty(\mathscr{O})$, $\mathscr{O} \subset R^d$ open, whose locally convex topology is defined in the following way: this space is given by the union of subspaces

$$\bigcup_{n=1}^{\infty} \mathscr{D}_{K_n}(\mathscr{O}),$$

where $\mathscr{D}_{K_n}(\mathscr{O}) \doteq \{f \in \mathscr{D}(\mathscr{O}) : \mathrm{supp} f \subset K_n\}$ has a locally convex topology given by the seminorms

$$\|f\|_{k,K_n} = \sum_{|\alpha|\leq k} \sup_{K_n} |\partial^{(\alpha)} f|,$$

and $K_n = \bar{K}_n \Subset \mathscr{O}$ satisfies $K_n \Subset K_{n+1}$ and$\cup_n K_n = \mathscr{O}$ (i.e., the collection $\{K_n\}_{n\in\mathbb{Z}_+}$ is an *exhaustion* of $\mathscr{O}$), implying $\mathscr{D}_{K_n}(\mathscr{O}) \subseteq \mathscr{D}_{K_{n+1}}(\mathscr{O})$ in a way that the topology of $\mathscr{D}_{K_n}(\mathscr{O})$ is clearly induced by the topology of $\mathscr{D}_{K_{n+1}}(\mathscr{O})$. The topology of $\mathscr{D}(\mathscr{O})$ is the finest topology such that the inclusions $\mathscr{D}_{K_n}(\mathscr{O}) \hookrightarrow \mathscr{D}(\mathscr{O})$ are continuous for all n, called *inductive limit* of (the topologies of) the collection $(\mathscr{D}_{K_n}(\mathscr{O}))_{n\in\mathbb{Z}_+}$.

A separating collection of seminorms which defines the topology of an inductive limit is not (notice!) necessarily countable, though the collection of seminorms given above is (i.e., the latter is not always sufficient to generate the topology of an inductive limit). If, however, $\mathscr{O}$ is replaced by K *compact* and/or the support requirements are abandoned, we enumerate the family of seminorms given above as $\|.\|_k$, $k = 1, 2, \ldots$. Defining

$$d(x,y) \doteq \sum_{k=1}^{\infty} \frac{\|x-y\|_k}{2^k(1+\|x-y\|_k)},$$

we have that $d(.,.)$ defines a translation-invariant metric which generates the topology of the space. We have, nevertheless, that $\mathscr{C}^\infty_{(c)}(\mathscr{O})$ and $\mathscr{C}^\infty_{(c)}(K)$ are *sequentially complete*, i.e., any Cauchy sequence (for general topological vector spaces, a sequence (x_n) is said to be Cauchy if, for any neighbourhood V of the origin, there exists $N \in \mathbb{Z}_+$ such that $x_n - x_m \in V$ for all $n, m > N$) converges. In particular, $\mathscr{C}^\infty(\mathscr{O})$ and $\mathscr{C}^\infty_{(c)}(K)$ are complete metric spaces (Fréchet *spaces*). The topology of the respective topological distribution duals is equivalent to the topology of pointwise convergence of sequences, being that this topology is also sequentially complete.

We can now give the

Definition B.10 *A* locally convex *-algebra *is a *-algebra $\mathfrak{F}$ which, as a vector space, is locally convex, and such that the operations of product and involution are continuous in this topology. We demand that the seminorms $\|.\|_\alpha$ of $\mathfrak{F}$ satisfy the C* property, i.e., $\|x^*x\|_\alpha = \|x\|^2_\alpha$, for all α.*

Let's now build the example which interests us the most, the Borchers-Uhlmann algebra [Bor62, Uhl62, Bor65]. The space $\mathscr{D}(\mathscr{O})$ is *nuclear*: there exists a unique locally convex topology in the algebraic tensor product $\mathscr{D}(\mathscr{O}) \otimes \mathscr{D}(\mathscr{O}) \subset \mathscr{D}(\mathscr{O} \times \mathscr{O})$ which extends the topology of $\mathscr{D}(\mathscr{O})$. The Cauchy completion of $\mathscr{D}(\mathscr{O}) \otimes \mathscr{D}(\mathscr{O})$ in this topology is precisely $\mathscr{D}(\mathscr{O} \times \mathscr{O})$. Dually, the *Kernel Theorem* of L. Schwartz [Hör90] asserts that to any continuous linear map $K : \mathscr{D}(\mathscr{O}) \to \mathscr{D}'(\mathscr{O})$ corresponds a unique $u \in \mathscr{D}'(\mathscr{O} \times \mathscr{O})$ such that $(K\phi)(\psi) = u(\phi \otimes \psi)$.

Definition B.11 *The* Borchers-Uhlmann *algebra associated to $\mathscr{O}$ is the unital, locally convex *-algebra $\mathfrak{F}(\mathscr{O})$ given by*

$$\mathfrak{F}(\mathscr{O}) \doteq \sum_{k=0}^{\infty} \otimes^k \mathscr{C}^\infty_c(\mathscr{O}),$$

where $\otimes^0 \mathscr{C}^\infty_c(\mathscr{O}) = \mathbb{C}\mathbb{1}$ denotes the adjunction of the unit $\mathbb{1}$. More precisely, it's the $\bar{\mathbb{Z}}_+$-graded algebra composed of elements $f = (f_{(0)}, f_{(1)}, \ldots)$, where $f_{(k)} \in \otimes^k \mathscr{C}^\infty_c(\mathscr{O})$ and $f_{(k)} = 0$ except for a finite set of values of k. The algebraic operations are given by:

- $\alpha f + \beta g = (\alpha f_{(0)} + \beta g_{(0)}, \alpha f_{(1)} + \beta g_{(1)}, \ldots,)$;
- $fg = f \otimes g = (f_{(0)} g_{(0)}, f_{(1)} g_{(0)} + f_{(0)} g_{(1)}, \ldots, \sum_{i+j=k} f_{(i)} \otimes g_{(j)}, \ldots)$;
- $f^* = (f^*_{(0)}, f^*_{(1)}, \ldots)$, *where* $f^*_{(0)} = \overline{f_{(0)}}$ *and, for* $f_{(k)} = f^1_{(k)}(x_1) \cdots f^k_{(k)}(x_k)$, $x_i \in \mathscr{O}$, $i = 1, \ldots, k$, *we define* $f^*_{(k)} = \overline{f^k_{(k)}(x_1)} \cdots \overline{f^1_{(k)}(x_k)}$, *extending the operation to* $\otimes^k \mathscr{C}^\infty_c(\mathscr{O})$ *by linearity.*

The seminorms of $\mathfrak{F}(\mathscr{O})$ are the ones inherited from $\mathscr{C}_c^\infty(\mathscr{O})$. It follows immediately from the Leibniz *rule that these seminorms satisfy the C* property.*

A state ω over $\mathfrak{F}(\mathscr{O})$ is given by a hierarchy of distributions $\omega_n \in \mathscr{D}'(\mathscr{O}^n)$, $\omega_0(f_{(0)}\mathbb{1}) = f_{(0)}$, subject to the requirement of *positivity*: $\sum_{n,m=0}^{k} \omega_{n+m}(f_{(n)}^* \otimes f_{(m)}) \geq 0$, $\forall f_{(n)} \in \otimes^n \mathscr{C}_c^\infty(\mathscr{O})$, $n = 0, \ldots, k$, $k \in \bar{\mathbb{Z}}_+$. Notice that positivity is enough to guarantee the weak continuity of ω_n, and, indeed, a good deal of regularity is guaranteed *a priori* for ω_n. One demands directly that the ω_n's define distributions to include the possibility of ω not being a positive functional (a situation in which this occurs is the case of gauge fields).

The Wightman-GNS representation of $\mathfrak{F}(\mathscr{O})$ associated to ω produces a distribution ϕ in $\mathscr{C}_c^\infty(\mathscr{O})$ with values in the set of closed linear operators (densely) defined in $\mathfrak{F}(\mathscr{O})/\mathrm{Ann}\omega \subset \mathscr{H}_\omega$. ϕ satisfies the hermiticity condition $\phi(\bar{f})^* = \phi(f)$.

Everything we've done above goes without essential changes from $\mathbb{R}^d$ to any d-dimensional manifold $\mathscr{M}$. Moreover, this example is the base for us to construct the Borchers-Uhlmann algebra associated to fields with arbitrary spin, which will be sketchly done as follows. Consider a vector bundle $\mathscr{E} \xrightarrow{p} \mathscr{M}$ with fiber $E \cong \mathbb{R}^D$. Given $\mathscr{O} \subset \mathscr{M}$ open, we take

$$\mathfrak{F}(\mathscr{O}) \doteq \sum_{k=0}^{\infty} \otimes^k \Gamma_c^\infty(\mathscr{O}, \mathscr{E}).$$

In this case, the k-point function ω_k associated to a state ω assumes values in $\otimes^k E$, and the field obtained is a D-tuple of operator-valued distributions.

We call here attention to the extremely important fact that the Borchers-Uhlmann algebra contains only *kinematic* information about a theory, determining the support, the tensor character and the global internal symmetries of the field (multiplet). *All* the *dynamical* information of the theory (equations of motion, commutation relations, spectrum, etc.) is contained in ω. We can alternatively define ϕ by taking the quotient of $\mathfrak{F}(\mathscr{O})$ modulo some *-ideal $\mathfrak{I}$ (we treat several instances of this scenario in Chapters 3 and 4). In this case, a *bona fide* state ω over the field ϕ must satisfy $\mathrm{Ann}\omega \supset \mathfrak{I}$, for it follows from the Third *-Isomorphism Theorem (Theorem B.4), supplemented by Remark B.2, that $(\mathfrak{F}(\mathscr{O})/\mathfrak{I})/(\mathrm{Ann}\omega/\mathfrak{I}) \cong \mathfrak{F}(\mathscr{O})/\mathrm{Ann}\omega$.

One can construct, under special circumstances (as, for instance, in the case of a quantum field theory satisfying the Wightman axioms), a version of the Tomita-Takesaki modular theory for *-algebras such as the Borchers-Uhlmann algebra. This problem, addressed for the first time by Bisognano and Wichmann [BW75, BW76], is attacked in a systematic fashion in [Ino98].

Appendix C

Categories and functors

Here, we shall introduce in a minimalistic way the needed concepts from the theory of categories and functors, a language devised with the purpose of organizing the structures that appear in the different areas of mathematics. Our reference for this Appendix is the classic text [Lan98] of Mac Lane, one of the fathers of the concept.

A brief comment about the use of set theory made when one speaks about categories is in order here. Strictly speaking, the concept of category [1] (Definitions C.1 and C.2) makes no mention whatsoever to any of the axioms of set theory. Indeed, it's common (and quite useful) to consider "large" categories, i.e., such that the objects and arrows (morphisms) constitute aggregates more general than sets, such as *classes* [2]. According to the system of set-theoretical axioms of Gödel and Bernays, the notion of class is realized by all positive instances of a certain propositional function that defines the properties of the objects of the class. For our purposes, it'll suffice to say that a class is a *set* if each one of the positive instances which forms the former reduces itself to a *single* element of the universe. Summing up, sets are "small" aggregates (i.e., elements of the universe) and classes, "large" aggregates.[3] We won't need more general set-theoretical concepts.

C.1 Rudiments

Definition C.1 *A* metacategory *consists of:*

[1] More precisely, of *metacategory* – see Definition C.1.

[2] The concept of class makes itself necessary every time one tries to build "sets" with cardinality bigger than the one of the universe of the set theory adopted (naïve or following the axioms of Zermelo-Fraenkel). The paradigmatic example, pointed by the first time by B. Russell, is the class of subsets of the universe.

[3] There are certain categories whose objects may not constitute a class, as in the case of categories of functors (see Section C.2).

1. objects $a, b, c, \ldots$;

2. arrows *(or* morphisms*)* $f, g, h, \ldots$;

3. domain *(or* source*)* $D(.)$ *and* counterdomain *(or* target*)* $CD(.)$ *operations on arrows, resulting in the objects* $D(f), CD(f), D(g), CD(g), \ldots$ *(it's customary to employ the notation*

$$f : a \longrightarrow b \quad or \quad a \xrightarrow{f} b, \quad for \quad a = D(f), b = CD(f));$$

4. *an* identity $id_{\cdot} \equiv \mathbb{1}_{\cdot}$ *operation on objects, resulting in the arrows* $\mathbb{1}_a, \mathbb{1}_b, \ldots$ *with* $D(\mathbb{1}_a) = CD(\mathbb{1}_a)$;

5. *and a* composition law $.\circ.$ *on ordered pairs of arrows* (f, g) *with* $D(f) = CD(g)$, *resulting in the arrows* $f \circ g$ *with* $D(f \circ g) = D(g)$ *and* $CD(f \circ g) = CD(f)$,

with the operations above subject to the following axioms:

Cat1 (associativity): *For any objects and arrows arranged in the form*

$$a \xrightarrow{f} b \xrightarrow{g} c \xrightarrow{h} d,$$

we have $h \circ (g \circ f) = (h \circ g) \circ f$;

Cat2 (unitary law): *For any arrow* f, *we have* $\mathbb{1}_{CD(f)} \circ f = f \circ \mathbb{1}_{D(f)} = f$.[4]

For our purposes, it'll be enough to consider a category as the following realization of a metacategory within set theory (see footnote 3):

Definition C.2 *A* category $\mathscr{C}$ *consists of a* class *of objects* $Obj\mathscr{C}$ *(sometimes identified, in abuse of notation, with* $\mathscr{C}$*), and a* class *of arrows* $Arr\mathscr{C}$ [5]*, endowed with operations satisfying the axioms of a metacategory,* such that the subclasses $Hom_{\mathscr{C}}(a, b) \doteq \{f : D(f) = a, CD(f) = b\}$ are sets, for any pair $a, b \in Obj\mathscr{C}$.

A *subcategory* $\mathscr{D}$ of $\mathscr{C}$ consists of a category whose objects form a subclass $Obj\mathscr{D}$ of $Obj\mathscr{C}$ and the arrows form a subclass $Arr\mathscr{D}$ of $Arr\mathscr{C}$, such that the operations of $\mathscr{D}$ are inherited from $\mathscr{C}$. $\mathscr{D}$ is a *full* subcategory if $Hom_{\mathscr{D}}(a, b) = Hom_{\mathscr{C}}(a, b)$, $\forall a, b \in \mathscr{D}$.

[4] Since this axiom determines $\mathbb{1}_a$ completely for any object a, it's sometimes convenient to identify a with the arrow $\mathbb{1}_a$. In particular, this makes possible to define a metacategory just by means of arrows (see [Lan98] for a detailed discussion).

[5] This, by virtue of the discussion in footnote 4, can be considered as a superclass of $Obj\mathscr{C}$. Under this point of view, it would be more adequate to identify $\mathscr{C}$ with $Arr\mathscr{C}$, but here we surrender to current practice.

We also say that a category $\mathscr{C}$ is *small* if $Obj\mathscr{C}$ (and, thus, $Arr\mathscr{C}$) is a set. We can associate $\mathscr{C}$ to the *opposite category* $\mathscr{C}^{op}$ such that $Obj\mathscr{C}^{op} = Obj\mathscr{C}$ and $Hom_{\mathscr{C}^{op}}(a,b) = Hom_{\mathscr{C}}(b,a)$, with the *inverse* composition law $\circ$ with respect to $\mathscr{C}$.

Since the arrows abstract from the category $\mathscr{S}et$ of sets the notion of functions between sets, it's natural to seek the categorical abstraction injective, surjective and bijective functions. We say that an arrow f is *surjective*, *epic* or simply an *epi* (resp. *injective*, *monic* or a *mono*) if, given two arrows $g_1, g_2 \in Hom_{\mathscr{C}}(CD(f), b)$ (resp. $h_1, h_2 \in Hom_{\mathscr{C}}(a, D(f))$) such that $g_1 \circ f = g_2 \circ f$ (resp. $f \circ h_1 = f \circ h_2$), we have $g_1 = g_2$ (resp. $h_1 = h_2$) [6]. An arrow e is *right* (resp. *left*) *invertible* if there exists an arrow e' such that $D(e') = CD(e)$, $CD(e') = D(e)$ and satisfying $e \circ e' = \mathbb{1}_{CD(e)}$ (resp. $e' \circ e = \mathbb{1}_{D(e)}$), and simply *invertible* (or an *isomorphism*) if both properties are satisfied *and any right inverse is a left one and vice-versa*. In this case, the inverse is unique, which does not happen to the previous cases. We also say in this case that $D(e)$ and $CD(e)$ are *isomorphic*. A *groupoid* is a small category such that any arrow is an isomorphism.

Let f be an epi and g a mono. If $f \circ g = \mathbb{1}_{D(g)}$, it's customary to say that g is a *section* of f (in analogy with sections of fiber bundles), and f a *retraction* of g. We also here say that f is a *split epi* and g, a *split mono*. In this case, $h = g \circ f : D(f) \longrightarrow D(f)$ is an *idempotent* arrow (or a *projection*), i.e., $h^2 \doteq h \circ h = h$. Conversely, an idempotent arrow $h \in Hom_{\mathscr{C}}(a,a)$ is said to be *split* by an object b if there exist an epi $f : a \longrightarrow b$ and a mono $g : b \longrightarrow a$ such that $f \circ g = \mathbb{1}_b$ and $g \circ f = h$.

Finally, let us move towards some examples, which will also be used to fix the notation for some standard categories which shall be frequently used:

1: Category endowed with a single object a and a single arrow $\mathbb{1}_a$;

2: Category endowed with only two objects a, b and three arrows $\mathbb{1}_a$, $\mathbb{1}_b$, $a \xrightarrow{f} b$;

U**:** (Small) *discrete* category, where $ObjU = U$ is a set, $\mathrm{Hom}_U(a,b) = \varnothing$ if $a \neq b$, and $\mathrm{Hom}_U(a,a)\{\mathbb{1}_a\}$ (summing up, we can also write $U = ObjU = ArrU$);

$\mathscr{C} \times \mathscr{D}$**:** *Product* of the categories $\mathscr{C}$ and $\mathscr{D}$ – $Obj\mathscr{C} \times \mathscr{D} \ni (a,b)$, where $a \in Obj\mathscr{C}$ and $b \in Obj\mathscr{D}$, $\mathrm{Hom}_{\mathscr{C}\times\mathscr{D}}((a,b),(c,d)) \ni (f : a \to c, g : b \to d)$, $\mathbb{1}_{(a,b)} = (\mathbb{1}_a, \mathbb{1}_b)$ and $(f,g) \circ (f',g') = (f \circ f', g \circ g')$;

[6] One haven't pursued a definition of epic (resp. monic) arrows by means of right (resp. left) inverses, for, although such definition applies to $\mathscr{S}et$, the same fails to happen, for instance, to the category $\mathscr{G}rp$ of groups – epimorphisms possesses a *set-theoretical* right inverse, for it may not be possible to choose an *homomorphism* with that property. The definition we gave, on the other hand, embraces all cases of interest.

$\mathscr{C}^I$: *Cartesian power* of a category $\mathscr{C}$ to the set I – formally, $Obj\mathscr{C} \ni a : I \to Obj\mathscr{C}$ (a is then said to be the *product* of the $a(\alpha)$'s), $\mathrm{Hom}_{\mathscr{C}^I}(a,b) = \{f : I \ni \alpha \mapsto f(\alpha) \in \mathrm{Hom}_{\mathscr{C}}(a(\alpha), b(\alpha))\}$, $\mathbb{1}_a : I \ni \alpha \mapsto \mathbb{1}_{a(\alpha)}$ and $(f \circ g) : I \ni \alpha \mapsto f(\alpha) \circ g(\alpha)$;

$(U, \leq)$: Partially ordered set – $Obj(U, \leq) = U$ is the underlying set, and $\mathscr{H}_{(U,\leq)}(a,b)$, if nonvoid, possesses a single arrow $\leq$, also denoted by $a \leq b$, satisfying the usual properties of an order relation, here suitably rephrased in categorical language: $\mathbb{1}_a = a \leq a$ and $(b \leq c) \circ (a \leq b) = a \leq c$. In this case, $Arr(U, \leq) \doteq \leq$ is said to be a *partial order* in U. Moreover, we say that $(U, \leq)$ is: *directed* if, given $a, b \in U$, there exists $c \in U$ such that $a \leq c$ and $b \leq c$; and *totally ordered* if for any $a, b \in U$ necessarily $a \leq b$ or $b \leq a$. The **Zorn** lemma here is stated in the following way: any full, totally ordered subcategory $\mathscr{C}$ of $(U, \leq)$ possesses a *terminal object* u, i.e., for any $a \in \mathscr{C}$ there exists an *unique* arrow $a \leq u$ (dually, $a \in Ob\mathscr{C}$ is an *initial* object if for any $b \in Ob\mathscr{C}$ there exists an *unique* arrow $a \to b$). An example of a directed set is the power set, ordered by inclusions $(U, \leq) = (P(X), \subseteq)$, of the set X – in particular, this directed set possesses a terminal object X and initial objects $\{x\}$, $x \in X$;

$\mathscr{S}et$: Sets, whose arrows are functions;

$\mathscr{G}rp$: Groups, whose arrows are homomorphisms;

$\mathscr{T}op$: Topological spaces, whose arrows are continuous functions;

$\mathscr{T}vs$: Topological vector spaces (assumed in the text Hausdorff and locally convex – see Section B.3), whose arrows are linear continuous maps.

Other categories shall be introduced in the text as needed.

C.2 Functors and natural transformations

Definition C.3 *Let $\mathscr{A}$ and $\mathscr{B}$ be categories. A* covariant *(resp.* contravariant*)* functor *from $\mathscr{A}$ to $\mathscr{B}$ is a rule $\mathfrak{F}$ which associates to each $a \in Ob\mathscr{A}$ an object $\mathfrak{F}a \in Ob\mathscr{B}$ and to each $f \in Arr\mathscr{A}$ an arrow $\mathfrak{F}f \in Arr\mathscr{B}$ satisfying $D(\mathfrak{F}f) = \mathfrak{F}D(f)$ (resp. $D(\mathfrak{F}f) = \mathfrak{F}CD(f)$), $CD(\mathfrak{F}f) = \mathfrak{F}CD(f)$ (resp. $CD(\mathfrak{F}f) = \mathfrak{F}D(f)$) and such that, for $a, b, c \in Ob\mathscr{A}$ and $f, g \in Arr\mathscr{A}$ satisfying $a \xrightarrow{f} b \xrightarrow{g} c$, we have*

$$\mathfrak{F}(g \circ f) = \mathfrak{F}g \circ \mathfrak{F}f \ (\textit{resp.}\ \mathfrak{F}(g \circ f) = \mathfrak{F}f \circ \mathfrak{F}g).$$

It's customary to write a functor as an "arrow between categories", i.e., $\mathfrak{F} : \mathscr{A} \to \mathscr{B}$ or even $\mathscr{A} \xrightarrow{\mathfrak{F}} \mathscr{B}$.

We can say, equivalently, that a contravariant functor $\mathfrak{F} : \mathscr{A} \to \mathscr{B}$ is a covariant functor $\mathfrak{F} : \mathscr{A} \to \mathscr{B}^{op}$. Hence, when we say functor, we just mean "covariant functor".

$\mathfrak{F}$ is said to be an *equivalence of categories* if it's invertible, i.e., there exists a functor $\mathfrak{F}^{-1}$ such that $\mathfrak{F}^{-1} \circ \mathfrak{F} = \mathfrak{id}_{\mathscr{A}}$ and $\mathfrak{F} \circ \mathfrak{F}^{-1} = \mathfrak{id}_{\mathscr{B}}$, where $\mathscr{C} \xrightarrow{\mathfrak{id}_{\mathscr{C}}} \mathscr{C}$ is the *identity* functor associated to the category $\mathscr{C}$ (i.e., $\mathfrak{id}_{\mathscr{C}} a = a$, $\mathfrak{id}_{\mathscr{C}} f = f$, $\forall a \in Obj\mathscr{C}$, $f \in Arr\mathscr{C}$). Another example is the functor $\mathfrak{i}_{\mathscr{C},\mathscr{D}}$ given by the inclusion of a category $\mathscr{C}$ into a supercategory $\mathscr{D}$ of $\mathscr{C}$.

We shall give now some examples of functors which will be of great use to us:

- Let $\mathscr{C}$ be a category and $(U, \leq)$ a partially ordered set. A functor $\mathfrak{F} : (U, \leq) \to \mathscr{C}^{op}$ (resp. $\mathfrak{G} : (U, \leq) \to \mathscr{C}$) is said to be a *pre(co)sheaf* (of objects of $\mathscr{C}$) in U.[7] If $(U, \leq)$ is directed, we say that $\mathfrak{F}$ is an *inverse system* in $\mathscr{C}$ indexed by U, and $\mathfrak{G}$, a *net* (of objects of $\mathscr{C}$) in U, or a *direct system* in $\mathscr{C}$ indexed by U;

- The Cartesian power of $\mathscr{C}$ to I can be understood as a functor $\mathfrak{F} : I \to \mathscr{C}$, where I is here understood as a (discrete) category;

- Let I be a set and $\mathscr{C}$ a category. The *diagonal functor* $\Delta : \mathscr{C} \to \mathscr{C}^I$ is given by $Obj\mathscr{C} \ni a \mapsto (\Delta a : I \ni \alpha \mapsto a)$ and $Arr\mathscr{C} \ni f \mapsto (\Delta f : I \ni \alpha \mapsto f)$;

- Given a functor $\mathfrak{F} : \mathscr{C} \to \mathscr{D}$ and $d \in Obj\mathscr{D}$, an *universal arrow* from d to $\mathfrak{F}$ (resp. from $\mathfrak{F}$ to d) consists in a pair (u, f), where $u \in Obj\mathscr{C}$ and $f : d \to \mathfrak{F}u$ (resp. $f : \mathfrak{F}u \to d$) which satisfy the following *universal property:* given any $c \in Obj\mathscr{C}$, $g : d \to \mathfrak{F}c$ (resp. $g : \mathfrak{F}c \to d$), there exists *a single arrow* $g' : u \to c$ (resp. $g' : c \to u$) such that $\mathfrak{F}g' \circ f = g$ (resp. $f \circ \mathfrak{F}g' = g$). In this case, we say that (u, f) is a universal element of the functor $\mathrm{Hom}_{\mathscr{D}}(d, \mathfrak{F}.) : \mathscr{C} \to \mathscr{S}et$ (the action of this functor on arrows is given by $\mathrm{Hom}_{\mathscr{D}}(d, \mathfrak{F}.) : (g : a \to b) \mapsto (\mathrm{Hom}_{\mathscr{D}}(d, \mathfrak{F}a) \ni h \mapsto \mathfrak{F}g \circ h \in \mathrm{Hom}(d, \mathfrak{F}b))$) – more generally, if $\mathfrak{H} : \mathscr{C} \to \mathscr{S}et$, an *universal object* of $\mathfrak{H}$ consists of the pair (u, e), where $u \in Obj\mathscr{C}$ and $e \in \mathfrak{H}u$ are such that for each $a \in Obj\mathscr{C}$, $x \in \mathfrak{H}a$ there exists *a single arrow* $f : u \to a$ such that $(\mathfrak{H}f)(e) = x$. Conversely, interpreting e as $e : \{*\} \ni * \mapsto (\mathfrak{H}u)(*) \doteq e$, we see that (u, e) is a universal arrow of $*$ in $\mathfrak{H}$. The dual case cited parallelly above consists only in exchanging $\mathscr{D}$ with $\mathscr{D}^{op}$. Examples of universal arrows comprise:

  - *Generators* of free groups ($\mathscr{C} = \mathscr{G}rp$, $\mathscr{D} = \mathscr{S}et$ e $\mathfrak{F}$ identifies a group with the underlying set): given $X \in Obj\mathscr{S}et$, let u be the free group generated by X and f identifying $x \in X$ with the corresponding element of u;

[7] Actually, the usual definition of pre(co)sheaf in U consists of a functor $\mathscr{F} : (P(U), \subseteq) \to \mathscr{C}$ or, more generally, $\mathfrak{F} : (\tau(U), \subseteq) \to \mathscr{C}$, where τ is a topology (possibly discrete, as in the previous case) in U. It's not difficult to see that such a definition is superseded by the one we've given above.

- *Coproducts* ($\mathscr{D} = \mathscr{C}^I$, where I is a set, and $\mathfrak{F} = \Delta$): given $a : I \ni \alpha \mapsto a(\alpha)$ and (u, f) and universal arrow from a to Δ, we then say that u is the *coproduct* or *direct sum* of the $a(\alpha)$'s, sometimes denoted $u \doteq \coprod_{\alpha\in I} a(\alpha)$ or $u \doteq \bigoplus_{\alpha\in I} a(\alpha)$, and $f(\alpha)$, the *injection* of $a(\alpha)$ in u;
- *Products* ($\mathscr{D} = \mathscr{C}^I$, where I is a set, and $\mathfrak{F} = \Delta$): given $a : I \ni \alpha \mapsto a(\alpha)$ and (u, f) a universal arrow from Δ to a, we then say that u is the *product* of the $a(\alpha)$'s, sometimes denoted $u \doteq \prod_{\alpha\in I} a(\alpha)$, and $f(\alpha)$, the *projection* of u in $a(\alpha)$;
- *Completion* of topological spaces, free tensor algebras, etc.,

and of universal objects:

- *Quotient* $S/\sim$ of a set S modulo an equivalence relation $\sim$ ($\mathscr{C} = \mathscr{S}et$, $\mathfrak{H} : X \mapsto \mathfrak{H}X = \{f : S \to X : s, s' \in S,\, s \sim s \Rightarrow f(s) = f(s')\}$);
- *Tensor product* of two vector spaces (particular case of the previous example), etc..

- Given a category $\mathscr{C}$, a directed set $(U, \leq)$ and a direct (resp. inverse) system $\mathfrak{F}$ in $\mathscr{C}$ indexed by U, the *inductive* (resp. *projective*) limit, or simply *colimit* (resp. *limit*) of $a \in Obj\mathscr{C}^U$ is simply a universal arrow (u, f) from a to Δ (resp. from Δ to a), such that $f(\alpha') = f(\alpha) \circ \mathfrak{F}(\alpha' \leq \alpha)$ (resp. $f(\alpha') = \mathfrak{F}(\alpha' \leq \alpha) \circ f(\alpha)$) – we denote

$$u \doteq \varinjlim_{\alpha,\mathfrak{F}} a(\alpha) \text{ (resp. } u \doteq \varprojlim_{\alpha,\mathfrak{F}} a(\alpha)). \tag{C.1}$$

Definition C.4 *Let $\mathscr{A}$ and $\mathscr{B}$ be categories, and the functors* [8] $\mathfrak{F}, \mathfrak{G} : \mathscr{A} \to \mathscr{B}$. *A* natural transformation Φ *from $\mathfrak{F}$ to $\mathfrak{G}$ is a rule that associates to each $a \in Obj\mathscr{A}$ an arrow $\Phi_a \in Arr\mathscr{B}$ satisfying $D(\Phi_a) = \mathfrak{F}a$, $CD(\Phi_a) = \mathfrak{G}a$ and such that, for each $f : a \to b$, $a, b \in Obj\mathscr{A}$, we have*

$$\Phi_a \circ \mathfrak{F}f = \mathfrak{G}f \circ \Phi_b.$$

A natural transformation ϕ between functors $\mathfrak{F}, \mathfrak{G}$ is said to be a *natural equivalence* between $\mathfrak{F}$ and $\mathfrak{G}$ if there exists a natural transformation ϕ^{-1} from $\mathfrak{G}$ to $\mathfrak{F}$ such that $\phi_a^{-1} \circ \phi_a = \phi_a \circ \phi_a^{-1} = \mathbb{1}_a$. For instance, a functor $\mathfrak{F}$ from a category $\mathscr{C}$ to a supercategory $\mathscr{D}$ of $\mathscr{C}$ is said to be *forgetful* id it's naturally equivalent to the inclusion functor $\mathscr{C}$ into $\mathscr{D}$, i.e., $\mathfrak{F}$ "forgets" part of the structure of $\mathscr{C}$.

Since the composition of natural transformations is clearly associative, and there always exists, for each functor $\mathfrak{F}$, an identity $\mathbb{1}_{\mathfrak{F}}$ given by $(\mathbb{1}_{\mathfrak{F}})_a \doteq \mathbb{1}_{\mathfrak{F}a}$, we can formally

[8] Using the notion of opposite category, we can extend the definition to contravariant functors.

consider the *functor category* $\mathscr{C}^{\mathscr{D}}$, with $Obj\mathscr{C}^{\mathscr{D}} = \{\mathfrak{F} : \mathscr{D} \to \mathscr{C}\}$ and $Arr\mathscr{C}^{\mathscr{D}}$ consisting of all natural transformations between functors from $\mathscr{D}$ to $\mathscr{C}$, *modulo* the set-theoretical caveats discussed at the beginning of this Appendix, in particular footnote 3. An example of such a category is, given a category $\mathscr{A}$, the *category of arrows* $\mathscr{A}^{\mathbf{2}}$.

Appendix D

Rudiments of homotopy

We introduce here some basic definitions and results which are employed at several points of Chapter 1. Our reference for the matter is Part I (Sections 1–7) of [GH81]. For applications related to differentiable manifolds (isotopies, etc.), we follow [Hir76].

D.1 Homotopic maps

Definition D.1 *Given two topological spaces X, Y and two continuous maps $f, g : X \to Y$, we say that f is* homotopic *to g if there exists a (jointly) continuous map $F : X \times [0,1] \to Y$ such that $F(x,0) = f(x)$ and $F(x,1) = g(x)$, for all $x \in X$ (such a F is said to be a* homotopy *between f and g). In particular, if $A \subset X$ and $f\restriction_A = g\restriction_A$, we say that f is* homotopic to g relative to A *if we can choose the homotopy F in a way that $F(x,t) = f(x) = g(x)$ for all $t \in [0,1]$ – in this case, we say that F is a* homotopy between f and g relative to A.

We say that X is *contractible* if the identity map $\mathrm{id}_X : X \ni x \mapsto x$ is homotopic to the constant map $f_p(x) : X \ni x \mapsto p$ for some $p \in X$. X is contractible if and only if, given any topological space Y, then any two continuous maps $f, g : Y \to X$ are homotopic – the direct implication is immediate, and the inverse implication is obtained through the homotopy $G : Y \times [0,1] \to X$ given by $G(y,t) = F(f(y), 2t)$ $(t \in [0, \frac{1}{2}])$ and $G(y,t) = F(g(y), 1-2t)$ $(t \in [\frac{1}{2}, 1])$, where F is a homotopy between id_X and f_p. As a consequence of this, a contractible topological space X is always *pathwise connected*, i.e., given any two points $x, y \in X$, there exists a continuous curve linking x to y – namely, $\gamma : [0,1] \to X$ given by $\gamma(t) = F(x, 2t)$ $(t \in [0, \frac{1}{2}])$ and $\gamma(t) = F(y, 1-2t)$ $(t \in [\frac{1}{2}, 1])$.

A contractible topological space X is also *simply connected*, i.e., any two continuous curves $\gamma, \gamma' : [0,1] \ni s \to X$ such that $\gamma(0) = \gamma'(0)$ and $\gamma(1) = \gamma'(1)$ are *fixed-endpoint*

homotopic, i.e., homotopic relative to $\{0,1\}$.

Let us now consider two differentiable manifolds $\mathscr{M},\mathscr{N}$. Given two maps $\mathscr{C}^\infty$ $f,g:\mathscr{M}\to\mathscr{N}$ and a homotopy F between f and g, it's always possible to regularize F in such a way that F is a $\mathscr{C}^\infty$ map from $\mathscr{M}\times[0,1]$ to $\mathscr{N}$. The same holds if $f=g$ in a subset $U\subset\mathscr{M}$ and F is a homotopy between f and g relative to U. Hence, once the concept of homotopy is transported from the $\mathscr{C}^0$ category to the $\mathscr{C}^\infty$ category, we can refine it in the following way:

- If $F(.,t)$ is an embedding of $\mathscr{M}$ into $\mathscr{N}$ for all $t\in[0,1]$ (and, in particular, f and g are as well), we say that F is an *isotopy* between f and g. A typical example of isotopy is given by a foliation of $K\subset\mathscr{N}$ by diffeomorphic copies of $\mathscr{M}$ – in this case, $K=F(\mathscr{M},[0,1])$ and $F(\mathscr{M},t_1)\cap F(\mathscr{M},t_2)=\varnothing$ if $t_1\neq t_2$.
- If $\mathscr{M}=\mathscr{N}$, $F(.,t)$ is a diffeomorphism for all $t\in[0,1]$ and $f=\mathrm{id}_{\mathscr{M}}$, we say that F is a *diffeotopy* or *ambient isotopy*. An example of diffeotopy is the restriction of an one-parameter group of diffeomorphisms $\mathscr{M}\times\mathbb{R}\ni(x,t)\mapsto\phi_t(x)$ (i.e., $\phi_{t_1+t_2}=\phi_{t_1}\circ\phi_{t_2}$ and $\phi_0=\mathrm{id}_{\mathscr{M}}$) to $\mathscr{M}\times[0,1]$ – in this case, $g=\phi_1$.

D.2 The fundamental group and covering spaces

Definition D.2 *Let X be a topological space and $x_0\in X$, called* base. *The* loop space *$\Omega(X,x_0)$ of X based on x_0 consists of the continuous curves $\gamma:[0,1]\to X$ such that $\gamma(0)=\gamma(1)=x_0$. Consider: the equivalence relation $\sim$ in $\Omega(X,x_0)$*

$$\gamma_1\sim\gamma_2\Leftrightarrow\gamma_1 \textit{ is fixed-endpoint homotopic to }\gamma_2,$$

the operations of

- Product: $\gamma_1,\gamma_2\in\Omega(X,x_0)\to\gamma_1\gamma_2$ *given by*

$$\gamma_1\gamma_2(t)=\begin{cases}\gamma_1(2t) & (t\in[0,\frac{1}{2}])\\ \gamma_2(2t-1) & (t\in[\frac{1}{2},1])\end{cases},e \tag{D.1}$$

- Inversion: $\gamma^{-1}(t)=\gamma(1-t)$

and the identity loop $\gamma_0(t)\equiv x_0$ *in* $\Omega(X,x_0)$. *The* fundamental group *(or* first homotopy group*) of X based on x_0 is set-theoretically given by $\pi_1(X,x_0)=\Omega(X,x_0)/\sim\ni[\gamma]$, with group operations $[\gamma_1][\gamma_2]\doteq[\gamma_1\gamma_2]$, $[\gamma]^{-1}\doteq[\gamma^{-1}]$ and identity $\mathbb{1}\doteq[\gamma_0]$.*

Clearly, the pathwise connected component X to which x_0 belongs is simply connected if and only if $\pi_1(X,x_0)=\{\mathbb{1}\}$. If X is pathwise connected, then $\pi_1(X,x_1)$ is isomorphic to $\pi_1(X,x_2)$ for all $x_1,x_2\in X$, and then we can define the *fundamental group* $\pi_1(X)\doteq\pi_1(X,x_0)$ of X.

DEFINITION D.3 *Given two topological spaces X, Y, a map $\phi : Y \to X$ is said to be a* covering map *if each $x \in X$ possesses an open neighbourhood $U \ni x$ such that $\phi^{-1}(U) = \dot{\cup}_\alpha V_\alpha$, where $\{V_\alpha\}$ is a collection of* disjoint *open subsets ($\dot{\cup}$ denotes disjoint union) of Y such that $\phi\restriction_{V_\alpha}$ is a homeomorphism such that $F(V_\alpha) = U$, for all α. In this case, we say that Y is a* covering (space) *of X – we also employ the condensed notation $Y \xrightarrow{\phi} X$. Y is said to be* universal *if it's pathwise and simply connected.*

If $Y \xrightarrow{\phi} X$ is a covering of the topological space X, it follows immediately from Definition D.3 that:

(i) $\phi^{-1}(x)$ is discrete for all $x \in X$,

(ii) ϕ is a local homeomorphism, and

(iii) ϕ is surjective, and the topology of X is precisely the quotient topology of E modulo the equivalence relation $y_1 \sim y_2 \Leftrightarrow \phi(y_1) = \phi(y_2)$.

The importance of the universal covering follows from the fact that, despite having quite a simple global structure, it encodes in a rather convenient way the fundamental group of the topological space it covers. Defining the group of *covering* (or *deck*) *transformations* $G(Y, X)$ of an universal covering Y of X as the group of homeomorphisms f from Y to itself such that $\phi \circ f = \phi$, one can prove [GH81] that $G(Y, X) \cong \pi_1(X)$.

Moreover, the universal covering Y of X is essentially *unique*: given any other universal covering $Y' \xrightarrow{\phi'} X$, there exists a homeomorphism $f : Y' \to Y$ such that $\phi \circ f = \phi'$. We can then speak about *the* universal covering of X. Conversely, given a group G of homeomorphisms of a pathwise and simply connected topological space Y which acts *properly discontinuously*, i.e., for all $y \in Y$ there exists an open neighbourhood $V \ni y$ such that $f(V) \cap V = \varnothing$ for all $f \in G$, it follows that $\pi_1(Y/G) \cong G$ – notice that, above, $G(Y, X)$ acts properly discontinuously in Y. We say then that $X = Y/G$ is the *fundamental domain* of Y with respect to G.

Bibliography

[AE73] H. Araki, G. Elliott. *On the Definition of C*-Algebras*. Publ. RIMS **9** (1973) 93–112.

[AGH98a] L. Andersson, G. J. Galloway, R. Howard. *A Strong Maximum Principle for Weak Solutions of Quasi-Linear Elliptic Equations with Applications to Lorentzian and Riemannian Geometry*. Commun. Pure Appl. Math. **L 1** (1998) 581–624. `arXiv:dg-ga/9707015`.

[AGH98b] —. *The Cosmological Time Function*. Clas. Quantum Grav. **15** (1998) 309–322. `arXiv:gr-qc/9709084`.

[And05] M. T. Anderson, *Unique Continuation Results for Ricci Curvature and Applications* (2005). Preprint, `arXiv:math.DG/0501067`.

[And06a] —, *Boundary Value Problems for Einstein Metrics, I* (2006). Preprint, `arXiv:math.DG/0612647`.

[And06b] —. *On the Uniqueness and Global Dynamics of AdS Spacetimes*. Clas. Quantum Grav. **23** (2006) 6935–6954. `arXiv:hep-th/0605293`.

[And06c] —, *Topics in Conformally Compact Einstein metrics*. In: *Perspectives in Riemannian Geometry, CRM Proc. Lecture Notes*, vol. **40**, 2006, 1–26. `arXiv:math.DG.0503243`.

[Ara76] H. Araki. *Relative Entropy of States of von Neumann Algebras*. Publ. RIMS **11** (1976) 809–833.

[Ara99] —, *Mathematical Theory of Quantum Fields*. Oxford University Press, 1999.

[Arn89] V. I. Arnold, *Mathematical Methods of Classical Mechanics*. 2nd. ed., Springer-Verlag, 1989.

[AS80] A. Ashtekar, A. Sen. *On the Role of Space-Time Topology in Quantum Phenomena: Superselection of Charge and Emergence of Nontrivial Vacua*. J. Math. Phys. **21** (1980) 526–533.

[AS02] M. Arnsdorf, L. Smolin, *The Maldacena Conjecture and Rehren Duality* (2002). Preprint, `arXiv:hep-th/0106073`.

[BBMS00] M. Bertola, J. Bros, U. Moschella, R. Schaeffer. *AdS/CFT Correspondence for n-point Functions*. Nucl. Phys. **B 587** (2000) 619–644. arXiv:hep-th/9908140.

[BCH73] J. M. Bardeen, B. Carter, S. W. Hawking. *The Four Laws of Black Hole Mechanics*. Commun. Math. Phys. **31** (1973) 161–170.

[BEE96] J. K. Beem, P. E. Ehrlich, K. L. Easley, *Global Lorentzian Geometry*. 2nd. ed., Marcel Dekker, 1996.

[Bek74] J. D. Bekenstein. *Generalized Second Law of Thermodynamics in Black-Hole Physics*. Phys. Rev. **D 9** (1974) 3292–3300.

[BF07] R. Brunetti, K. Fredenhagen, *Towards a Background Independent Formulation of Perturbative Quantum Gravity*. In: *Proceedings of the "Workshop on Mathematical and Physical Aspects of Quantum Gravity"*, Birkhäuser, Blaubeuren, 2007, 151–159. arXiv:gr-qc/0603079.

[BFK96] R. Brunetti, K. Fredenhagen, M. Köhler. *The Microlocal Spectrum Condition and Wick Polynomials of Free Fields on Curved Spacetimes*. Commun. Math. Phys. **180** (1996) 633–652. arXiv:gr-qc/9510056.

[BFS00] D. Buchholz, M. Florig, S. J. Summers. *The Second Law of Thermodynamics, TCP, and Einstein Causality in Anti-de Sitter Space-Time*. Clas. Quantum Grav. **17** (2000) L31–L37. arXiv:hep-th/9905178.

[BFV03] R. Brunetti, K. Fredenhagen, R. Verch. *The Generally Covariant Locality Principle – A New Paradigm for Local Quantum Physics*. Commun. Math. Phys. **237** (2003) 31–68. arXiv:math-ph/0112041.

[BGL93] R. Brunetti, D. Guido, R. Longo. *Modular Structure and Duality in Conformal Quantum Field Theory*. Commun. Math. Phys. **156** (1993) 201–219. arXiv:funct-an/9302008.

[BGM71] M. Berger, P. Gauduchon, E. Mazet, *Le Spectre d'une Varieté Riemannienne*, *Lecture Notes in Mathematics*, vol. **194**. Springer-Verlag, 1971.

[Bla85] B. Blackadar. *Shape Theory for C*-Algebras*. Math. Scand. **56** (1985) 249–275.

[Bla06] —, *Operator Algebras – Theory of C*-Algebras and von Neumann Algebras*, *Encyclopaedia of Mathematical Sciences*, vol. **122**. Springer-Verlag, 2006.

[BMT88] D. Buchholz, G. Mack, I. Todorov. *The Current Algebra on the Circle as a Germ of Local Field Theories*. Nucl. Phys. B (Proc. Suppl.) **5B** (1988) 20–56.

[Bor62] H.-J. Borchers. *On Structure of the Algebra of Field-Operators*. Nuovo Cimento **24** (1962) 214–236.

[Bor65] —. *On the Structure of the Algebra of Field-Operators II*. J. Math. Phys. **1** (1965) 49–56.

[BOR02] D. Buchholz, I. Ojima, H. Roos. *Thermodynamic Properties of Non-equilibrium States in Quantum Field Theory*. Ann. Phys. (N.Y.) **297** (2002) 219–242. arXiv:hep-ph/0105051.

[Bos05] H. Bostelmann. *Operator Product Expansions as a Consequence of Phase Space Properties*. J. Math. Phys. **46** (2005) 082.304.

[BR87] O. Bratteli, D. W. Robinson, *Operator Algebras and Quantum Statistical Mechanics I – C*- and W*-Algebras, Symmetry Groups, Decomposition of States*. 2nd. ed., Springer-Verlag, 1987.

[BR97] —, *Operator Algebras and Quantum Statistical Mechanics II – Equilibrium States, Models in Quantum Statistical Mechanics*. 2nd. ed., Springer-Verlag, 1997.

[BR02] R. Bousso, L. Randall. *Holographic Domains of Anti-de Sitter Space*. JHEP **0204** (2002) 057. arXiv:hep-th/0112080.

[BR07] R. Brunetti, G. Ruzzi. *Superselection Sectors and General Covariance. I*. Commun. Math. Phys. **270** (2007) 69–108.

[BS93] D. Buchholz, S. J. Summers. *An Algebraic Characterization of Vacuum States in Minkowski Space*. Commun. Math. Phys. **155** (1993) 449–458.

[BS03] A. N. Bernal, M. Sánchez. *On Smooth Cauchy Hypersurfaces and Geroch's Splitting Theorem*. Commun. Math. Phys. **243** (2003) 461–470. arXiv:gr-qc/0306108.

[BS05] —. *Smoothness of Time Functions and the Metric Splitting of Globally Hyperbolic Spacetimes*. Commun. Math. Phys. **257** (2005) 43–50. arXiv:gr-qc/0401112.

[BS06] —. *Further Results on the Smoothability of Cauchy Hypersurfaces and Cauchy Time Functions*. Lett. Math. Phys. **77** (2006) 183–197. arXiv:gr-qc/0512095.

[BS07] —. *Globally Hyperbolic Spacetimes can be Defined as "Causal" instead of "Strongly Causal"*. Clas. Quantum Grav. **24** (2007) 745–749. arXiv:gr-qc/0611138.

[BV95] D. Buchholz, R. Verch. *Scaling Algebras and Renormalization Group in Algebraic Quantum Field Theory*. Rev. Math. Phys. **7** (1995) 1195–1240. arXiv:hep-th/9501063.

[BW75] J. J. Bisognano, E. H. Wichmann. *On the Duality Condition for a Hermitian Scalar Field*. J. Math. Phys. **16** (1975) 985–1007.

[BW76] —. *On the Duality Condition for Quantum Fields*. J. Math. Phys. **17** (1976) 303–321.

[BY99] H.-J. Borchers, J. Yngvason. *Modular Groups of Quantum Fields in Thermal States*. J. Math. Phys. **40** (1999) 601–624. arXiv:math-ph/9805013.

[CK93] D. Christodoulou, S. Klainerman, *The Global Nonlinear Stability of the Minkowski Space*. Princeton University Press, 1993.

[DH72] J. J. Duistermaat, L. Hörmander. *Fourier Integral Operators II*. Acta Math. **128** (1972) 183–269.

[Die88] J. Dieckmann. *Cauchy Surfaces in a Globally Hyperbolic Space-Time*. J. Math. Phys. **29** (1988) 578–579.

[DR02] M. Dütsch, K.-H. Rehren. *A Comment on the Dual Field in the AdS-CFT Correspondence*. Lett. Math. Phys. **62** (2002) 171–184. arXiv:hep-th/0204123.

[DR03] —. *Generalized Free Fields and the AdS-CFT Correspondence*. Ann. Henri Poincaré **4** (2003) 613–635. arXiv:math-ph/0209035.

[Dri77] W. Driessler. *On the Type of Local Algebras in Quantum Field Theory*. Commun. Math. Phys. **53** (1977) 295–297.

[Dug66] J. Dugundji, *Topology*. Allyn and Bacon, 1966.

[Dui96] J. J. Duistermaat, *Fourier Integral Operators*, *Progress in Mathematics*, vol. **130**. Birkhäuser, 1996.

[EG92] L. C. Evans, R. F. Gariepy, *Measure Theory and Fine Properties of Functions*. CRC/Longman Press, 1992.

[ES00] P. E. Ehrlich, M. Sánchez. *Some Semi-Riemannian Volume Comparison Theorems*. Tohoku Math. J. **52** (2000) 331–348. arXiv:math.9811166.

[FG85] C. Fefferman, C. R. Graham, *Conformal Invariants*. In: *Élie Cartan et les mathématiques d'aujourd'hui*, *Astérisque*, vol. **hors série**, Societé Mathématique de France, Lyon, 1985, 95–116.

[FH81] K. Fredenhagen, J. Hertel. *Local Algebras of Observables and Point-like Localized Fields*. Commun. Math. Phys. **80** (1981) 555–561.

[FH90] K. Fredenhagen, R. Haag. *On the Derivation of the Hawking Radition Associated with the Formation of a Black Hole*. Commun. Math. Phys. **127** (1990) 273.

[Fri75] F. G. Friedlander, *The Wave Equation on a Curved Space-Time*. Cambridge University Press, 1975.

[Gal00] G. J. Galloway. *Maximum Principles for Null Hypersurfaces and Null Splitting Theorems*. Ann. Henri Poincaré **1** (2000) 543–567. arXiv:math.DG/9909158.

[Ger68] R. Geroch. *What is a Singularity in General Relativity?* Ann. Phys. (N.Y.) **48** (1968) 526–540.

[Ger70] —. *Domain of Dependence*. J. Math. Phys. **11** (1970) 437–449.

[GH81] M. J. Greenberg, J. R. Harper, *Algebraic Topology – A First Course*. Addison-Wesley, 1981.

[GKP72] R. Geroch, E. H. Kronheimer, R. Penrose. *Ideal Points in Space-Time*. Proc. Roy. Soc. Lond. **A 327** (1972) 545–567.

[GKP98] S. S. Gubser, I. R. Klebanov, A. M. Polyakov. *Gauge Theory Correlators from Non-Critical String Theory*. Phys. Lett. **B 428** (1998) 105–114. arXiv:hep-th/9802109.

[GL91] C. R. Graham, J. M. Lee. *Einstein Metrics with Prescribed Conformal Infinity on the Ball*. Adv. Math. **87** (1991) 186–225.

[GLRV01] D. Guido, R. Longo, J. E. Roberts, R. Verch. *Charged Sectors, Spin and Statistics in Quantum Field Theory on Curved Spacetimes*. Rev. Math. Phys. **13** (2001) 125–198. arXiv:math-ph/9906019.

[GN43] I. M. Gel'fand, M. A. Naimark. *On the Imbedding of Normed Rings into the Ring of Operators in Hilbert Space*. Mat. Sb. **12** (1943) 197–213.

[Gra00] C. R. Graham. *Volume and Area Renormalizations for Conformally Compact Einstein Metrics*. Rend. Circ. Mat. Palermo Ser. II Suppl. **63** (2000) 31–42. arXiv:math.DG/9909042.

[GSWW99] G. Galloway, K. Schleich, D. M. Witt, E. Woolgar. *Topological Censorship and Higher Genus Black Holes*. Phys. Rev. **D 60** (1999) 104.039. arXiv:gr-qc/9902061.

[GT98] D. Gilbarg, N. S. Trudinger, *Elliptic Partial Differential Equations of Second Order*. 2nd. ed., Springer-Verlag, 1998.

[GW00] S. Gao, R. M. Wald. *Theorems on Gravitational Time Delay and Related Issues*. Clas. Quantum Grav. **17** (2000) 4999–5008. arXiv:gr-qc/0007021.

[Haa96] R. Haag, *Local Quantum Physics – Fields, Particles, Algebras*. 2nd. ed., Springer-Verlag, 1996.

[Had03] J. Hadamard, *Lectures on Cauchy's Problem in Linear Partial Differential Equations*. Dover, 1923, 1952, 2003.

[Haw71] S. W. Hawking. *Gravitational Radiation from Colliding Black Holes*. Phys. Rev. Lett. **26** (1971) 1344–1346.

[Haw75] —. *Particle Creation by Black Holes*. Commun. Math. Phys. **43** (1975) 199–220.

[HE73] S. W. Hawking, G. F. R. Ellis, *The Large Scale Structure of Space-Time*. Cambridge University Press, 1973.

[HH00] G. T. Horowitz, V. E. Hubeny. *Quasinormal Modes of AdS Black Holes and the Approach to Thermal Equilibrium*. Phys. Rev. **D 62** (2000) 024.027. arXiv:hep-th/9909056.

[HHW67] R. Haag, N. M. Hugenholtz, M. Winnink. *On the Equilibrium States in Quantum Statistical Mechanics*. Commun. Math. Phys. **5** (1967) 215–236.

[HIM05] S. Hollands, A. Ishibashi, D. Marolf. *Comparison between Various Notions of Conserved Charges in Asymptotically AdS-spacetimes*. Clas. Quantum Grav. **22** (2005) 2881–2920. arXiv:hep-th/0503045.

[Hir76] M. W. Hirsch, *Differential Topology*. Springer-Verlag, 1976.

[HK64] R. Haag, D. Kastler. *An Algebraic Approach to Quantum Field Theory*. J. Math. Phys. **5** (1964) 848–861.

[Hoo93] G. 't Hooft, *Dimensional Reduction in Quantum Gravity*. In: *Salam-Festschrift* (A. Ali et al., ed.), World Scientific, 1993, 284–296. arXiv:gr-qc/9310026.

[Hör71] L. Hörmander. *Fourier Integral Operators I*. Acta Math. **127** (1971) 79–183.

[Hör90] —, *The Analysis of Linear Partial Differential Operators I – Distribution Theory and Fourier Analysis*. 2nd. ed., Springer-Verlag, 1990.

[HS98] M. Henningson, K. Skenderis. *The Holographic Weyl Anomaly*. JHEP **9807** (1998) 023. arXiv:hep-th/9806087.

[HS00] —. *Holography and the Weyl Anomaly*. Fortsch. Phys. **48** (2000) 125–128. arXiv:hep-th/9812032.

[HSS01] S. de Haro, K. Skenderis, S. Solodukhin. *Holographic Reconstruction of Spacetime and Renormalization in the AdS/CFT Correspondence*. Commun. Math. Phys. **217** (2001) 595–622. arXiv:hep-th/0002230.

[Ino98] A. Inoue, *Tomita-Takesaki Theory in Algebras of Unbounded Operators*, *Lecture Notes in Mathematics*, vol. **1699**. Springer-Verlag, 1998.

[Isr86] W. Israel. *Third Law of Black Hole Dynamics: A Formulation and Proof*. Phys. Rev. Lett. **57** (1986) 397–399.

[Jac85] N. Jacobson, *Basic Algebra I*. 2nd. ed., W. H. Freeman Co., 1985.

[Joh82] F. John, *Partial Differential Equations*. 4th. ed., Springer-Verlag, 1982.

[Kad00] L. P. Kadanoff, *Statistical Physics – Statics, Dynamics and Renormalization*. World Scientific, 2000.

[KF75] A. N. Kolmogorov, S. V. Fomin, *Introductory Real Analysis (Translation by R. A. Silverman)*. Dover, 1975.

[Kic04] S. Kichenassamy. *On a Conjecture of Fefferman and Graham*. Adv. Math. **184** (2004) 268–288.

[KR86] R. V. Kadison, J. R. Ringrose, *Fundamentals of the Theory of Operator Algebras – vol. 2 (Advanced Theory)*. Academic Press, 1986.

[Lan69] L. J. Landau. *A Note on Extended Locality*. Commun. Math. Phys. (1969) 246–253.

[Lan98] S. Mac Lane, *Categories for the Working Mathematician*. 2nd. ed., Springer-Verlag, 1998.

[Lan99] S. Lang, *Complex Analysis*. 4th. ed., Springer-Verlag, 1999.

[Lie67a] E. H. Lieb. *Exact Solution of the Problem of the Entropy of Two-Dimensional Ice*. Phys. Rev. Lett. **18** (1967) 692–694.

[Lie67b] —. *Residual Entropy of Square Ice*. Phys. Rev. **162** (1967) 162–172.

[LM75] M. Lüscher, G. Mack. *Global Conformal Invariance in Quantum Field Theory*. Commun. Math. Phys. **41** (1975) 203–234.

[Mal98] J. Maldacena. *The Large N Limit of Superconformal Field Theories and Supergravity*. Adv. Theor. Math. Phys. **2** (1998) 231–252. `arXiv:hep-th/9711200`.

[Min00] R. A. Minlos, *Introduction to Mathematical Statistical Mechanics*, *University Lecture Series*, vol. **19**. American Mathematical Society, 2000.

[MR03] P. Martinetti, C. Rovelli. *Diamond's Temperature: Unruh Effect for Bounded Trajectories and Thermal Time Hypothesis*. Clas. Quantum Grav. **20** (2003) 4919–4931.

[Müg96] M. Müger, *Disorder Operators, Quantum Doubles, and Haag Duality in 1+1 Dimensions* (1996). Seminar given at NATO ASI "Quantum Fields and Quantum Spacetime", Cargèse, `arXiv:hep-th/9611131`.

[Müg98] —. *Quantum Double Actions on Operator Algebras and Orbifold Quantum Field Theories*. Commun. Math. Phys. **191** (1998) 137–181. arXiv:hep-th/9606175.

[Mun71] M. E. Munroe, *Measure and Integration*. 2nd. ed., Addison-Wesley, 1971.

[Mur90] G. J. Murphy, *C*-algebras and Operator Theory*. Academic Press, 1990.

[O'N83] B. O'Neill, *Semi-Riemannian Geometry*. Academic Press, 1983.

[OP93] M. Ohya, D. Petz, *Quantum Entropy and its Use*. Springer-Verlag, 1993.

[PSW93] R. Penrose, R. D. Sorkin, E. Woolgar, *A Positive Mass Theorem Based on the Focusing and Retardation of Null Geodesics* (1993). Preprint, arXiv:gr-qc/9301015.

[PSW02] D. N. Page, S. Surya, E. Woolgar. *Positive Mass from Holographic Causality*. Phys. Rev. Lett. **89** (2002) 121.301. arXiv:hep-th/0204198.

[Reh00] K.-H. Rehren. *Algebraic Holography*. Ann. Henri Poincaré **1** (2000) 607–623. arXiv:hep-th/9905179.

[Reh05] —, *QFT Lectures on AdS-CFT*. In: *Proceedings of the 3rd Summer School in Modern Mathematical Physics, Institue of Physics, Belgrade*, 2005, 95–118. arXiv:hep-th/0411086.

[Rib04] P. L. Ribeiro, *Renormalization Group Flow in Algebraic Holography*. In: *Proceedings of the International Conference on Renormalization Group and Anomalies in Gravity and Cosmology* (M. Asorey, I. L. Shapiro, eds.), *Nucl. Phys. B (Proc. Suppl.)*, vol. **127**, Ouro Preto, Brazil, 2004, 193–195. arXiv:hep-th/0306024.

[Rib07] —, *Algebraic Holography in Asymptotically Simple, Asymptotically AdS Spacetimes*. In: *Rigorous Quantum Field Theory – a Festschrift in the Honour of Jacques Bros* (A. Boutet de Monvel, D. Buchholz, D. Iagolnitzer, U. Moschella, eds.), *Progress in Mathematics*, vol. **251**, Birkhäuser, SPhT-CEA Saclay, Paris, 2007, 253–270. arXiv:hep-th/0502096.

[Rob04] J. E. Roberts, *More Lectures in Algebraic Quantum Field Theory*. In: *Noncommutative Geometry – C.I.M.E. Lectures* (S. Doplicher, R. Longo, eds.), *Lecture Notes in Mathematics*, vol. **1831**, Springer-Verlag, Martina Franca, Italy, 2004, 263–342.

[RS75] M. Reed, B. Simon, *Methods of Modern Mathematical Physics II – Fourier Analysis, Self-Adjointness*. Academic Press, 1975.

[RS80] —, *Methods of Modern Mathematical Physics I – Functional Analysis*. Revised ed., Academic Press, 1980.

[Rud87] W. Rudin, *Real and Complex Analysis*. 3rd. ed., McGraw-Hill, 1987.

[Rud91] —, *Functional Analysis*. 2nd. ed., McGraw-Hill, 1991.

[Sch96] D. Schlingemann. *On the Algebraic Theory of Soliton and Antisoliton Sectors*. Rev. Math. Phys. **8** (1996) 301–326. `arXiv:hep-th/9601015`.

[Sch05] B. Schroer, *Area Density of Localization-Entropy II: Double Cone-Holography, Holographic Symmetry and its Relation to the Classical BMS Group* (2005). Preprint, `arXiv:hep-th/0511291`.

[Sch06] —. *Area Density of Localization-Entropy I: the Case of Wedge-Localization*. Clas. Quantum Grav. **23** (2006) 5227–5248. `arXiv:hep-th/0507038`.

[Seg47] I. E. Segal. *Postulates for General Quantum Mechanics*. Ann. Math. **48** (1947) 930–948.

[Sei77] H.-J. Seifert. *Smoothing and Extending Cosmic Time Functions*. Gen. Rel. Grav. **8** (1977) 815–831.

[Sew82] G. L. Sewell. *Quantum Fields on Manifolds: PCT and Gravitationally Induced Thermal States*. Ann. Phys. (N.Y.) **141** (1982) 201–224.

[Som06] C. Sommer, *Algebraische Charakterisierung von Randbedingungen in der Quantenfeldtheorie* (2006). Diplomarbeit, Universität Hamburg, `http://www.desy.de/uni-th/lqp/psfiles/dipl-sommer.ps.gz`.

[Sor79] R. Sorkin. *The Quantum Electromagnetic Field in Multiply Connected Space*. J. Phys. **A**: Math. Gen. **12** (1979) 403–421.

[Spi79] M. Spivak, *A Comprehensive Introduction to Differential Geometry – Volume 2*. 2nd. ed., Publish or Perish, 1979.

[SS02] D. T. Son, A. O. Starinets. *Minkowski-Space Correlators in AdS/CFT Correspondence: Recipe and Applications*. JHEP **0209** (2002) 042. `arXiv:hep-th/0205051`.

[Str00] A. Strohmaier. *On the Local Structure of the Klein-Gordon Field on Curved Spacetimes*. Lett. Math. Phys. **54** (2000) 249–261. `arXiv:math-ph/0008043`.

[Sus95] L. Susskind. *The World as a Hologram*. J. Math. Phys. **36** (1995) 6377–6396. `arXiv:hep-th/9409089`.

[SV96] A. Strominger, C. Vafa. *Microscopic Origin of the Bekenstein-Hawking Entropy*. Phys. Lett. **B 379** (1996) 99–104.

[SW00] R. F. Streater, A. S. Wightman, *PCT, Spin and Statistics, and All That*. Revised ed., Princeton University Press, 2000.

[Sán05] M. Sánchez, *Causal Hierarchy of Spacetimes, Temporal Functions and Smoothness of Geroch's Splitting. A Revision*. In: *Proceedings of the XIII School of Differential Geometry – Part II* (P. Piccione, ed.), *Matemática Contemporânea*, vol. **29**, Universidade de São Paulo, São Paulo, Brazil, 2005, 127–155. `arXiv:gr-qc/0411143`.

[Tak70] M. Takesaki, *Tomita's Theory of Modular Hilbert Algebras and its Applications*, *Lecture Notes in Mathematics*, vol. **128**. Springer-Verlag, 1970.

[Tak01] —, *Theory of Operator Algebras I*, *Encyclopaedia of Mathematical Sciences*, vol. **124**. Springer-Verlag, 2001.

[Tak03a] —, *Theory of Operator Algebras II*, *Encyclopaedia of Mathematical Sciences*, vol. **125**. Springer-Verlag, 2003.

[Tak03b] —, *Theory of Operator Algebras III*, *Encyclopaedia of Mathematical Sciences*, vol. **127**. Springer-Verlag, 2003.

[Tom67] M. Tomita, *Quasi-Standard von Neumann Algebras* (1967). Preprint.

[Uhl62] A. Uhlmann. *Über die Definition der Quantenfelder nach Wightman und Haag*. Wiss. Z. Karl-Marx-Univ. Leipzig Math.-Nat. Reihe **11** (1962) 213–217.

[Ver96] R. Verch, *Scaling Algebras, the Renormalization Group and the Principle of Local Stability in Algebraic Quantum Field Theory*. In: *Proceedings of the Conference on Operator Algebras and Quantum Field Theory*, Academia dei Lincei, Roma, 1996, 564–577. `arXiv:funct-an/9611004`.

[Wal80] R. M. Wald. *Dynamics in Nonglobally Hyperbolic, Static Space-Times*. J. Math. Phys. **21** (1980) 2802–2805.

[Wal84] —, *General Relativity*. Chicago University Press, 1984.

[Wal94] —, *Quantum Field Theory in Curved Spacetime and Black Hole Thermodynamics*. Chicago University Press, 1994.

[Wal97] —. *"Nernst Theorem" and Black Hole Thermodynamics*. Phys. Rev. **D 56** (1997) 6467–6474.

[Wei96] S. Weinberg, *The Quantum Theory of Fields II: Modern Applications*. Cambridge University Press, 1996.

[Whi34] H. Whitney. *Analytic Extension of Differentiable Functions Defined in Closed Sets*. Trans. Amer. Math. Soc. **36** (1934) 63–89.

[Wit98] E. Witten. *Anti de Sitter Space and Holography*. Adv. Theor. Math. Phys. **2** (1998) 253–291. `arXiv:hep-th/9802150`.

[Woo94] E. Woolgar. *The Positivity of Energy for Asymptotically Anti-de Sitter Spacetimes*. Clas. Quantum Grav. **11** (1994) 1881–1900. `arXiv:gr-qc/9404019`.

[Zwi04] B. Zwiebach, *An Introduction to String Theory*. Cambridge University Press, 2004.

Notes and translation of (some) epigraphs

[i] "See what things are in themselves, distinguishing matter, cause, end."

[ii] The literal translation is *"To myself"*. However, the "official" Latin translation for the title ended up being the name by which these personal writings, originally in Greek, of the Roman emperor have become more well known – *"Meditations"*.

[iii] "We only listen to the questions for which we are able to find an answer."

[iv] *"The Gay Science"*.

[v] Editora Melhoramentos, 1986.

[vi]

> "The truth is that when his mind was completely gone, he had the strangest thought any lunatic in the world ever had, which was that it seemed reasonable and necessary to him, both for the sake of his honor and as a service to the nation, to become a knight errant and travel the world with his armor and his horse to seek adventures and engage in everything he had read that knights errant engaged in, righting all manner of wrongs and, by seizing the opportunity and placing himself in danger and ending those wrongs, winning eternal renown and everlasting fame. The poor man imagined himself already wearing the crown, won by the valor of his arm, of the empire of Trebizond at the very least; and so it was that with these exceedingly agreeable thoughts, and carried away by the extraordinary pleasure he took in them, he hastened to put into effect what he so fervently desired. And the first thing he did was to attempt to clean some armor that had belonged to his great-grandfathers and, stained with rust and covered with mildew, had spent many long years stored and forgotten in a corner."

(Tr. Edith Grossman)

[vii] *"Don Quixote"*. Ecco/Harper Collins, 2003.

[viii] Editora Nova Fronteira, 1988.

[ix]

> " 'It seems to me,' said Don Quixote, 'there is no human history in the world that does not have its ups and downs, especially those that deal with chivalry; they cannot be filled with nothing but successful exploits.' "

(Tr. Edith Grossman)

[x] "There are two ways to reach Despina: by ship or by camel. The city shows itself in a different way to who arrives by land or by sea. (...) Each city receives the form of the desert which it opposes; Thus do the cameleer and the sailor see Despina, city of confine between two deserts." (Tr. Pedro Lauridsen Ribeiro)

[xi] *"'The cities and the desire 3' (The invisible cities)"*.

[xii]

" 'How can that be?' responded Don Quixote. 'Is it so essential to the story to know the exact number of goats that have crossed that a mistake in the count means you cannot continue the tale?'

'No, Señor, I can't.' responded Sancho, 'because as soon as I asked your grace to tell me how many goats had crossed, and you said you didn't know, at that very moment I forgot everything I had left to say, and, by my faith, it was very interesting and pleasing.'

'Do you mean to say that the story is finished?' said Don Quixote.

'As finished as my mother.' said Sancho.

'I tell you truthfully,' responded Don Quixote, 'that you have told one of the strangest tales, stories or histories that anyone in the world ever thought of, and this manner of telling it and then stopping it is something I shall never see, and have never seen, in my life, although I expected nothing else from your intellect; but I am not surprised, for perhaps the sound of the pounding, which has not ceased, has clouded your understanding.' "

(Tr. Edith Grossman)

The engravings in the frontispieces of each Part are clippings, done by the Author, from illustrations of Gustave Doré for *"El ingenioso hidalgo don Quijote de la Mancha"* (Primero Libro), of Miguel de Cervantes Saavedra. The illustrations, originally drawn with feather tip, were wood-engraved by Héliodore Pisan, and then employed in the edition annotated and translated to French by Louis Viardot, published by Hachette in 1863 (republished in 1978). Since they have been published for the first time more than 100 years ago, these illustrations are in public domain, and can be found at Wikimedia Commons (`http://commons.wikimedia.org/wiki/Don_Quixote`) in electronic format.

Erratum for the previous versions

In the original, Portuguese version and in the previous version of the English translation of this thesis, there was a completely incorrect construction of a measure from the Lorentzian distance by means of CARATHÉODORY's procedure in Subsection 2.3.1, which (thankfully!) showed up to be irrelevant for our purposes. Thus, this part was removed and replaced by appropriate remarks and a correct alternative argument which relies only on Proposition 2.4. We starkly warn the reader and heartily apologise for such errors when reading either the original Portuguese version of the previous English version in the arXiv.

The Author
Toruń, June 2008